M. Mustafa, D. Suragan, A note on Hardy type inequalities for quaternion-valued functions, *Sib. Èlektron. Mat. Izv.*, 2021, Volume 18, Issue 1, 338–344

DOI: <https://doi.org/10.33048/semi.2021.18.023>

Use of the all-Russian mathematical portal Math-Net.Ru implies that you have read and agreed to these terms of use
<http://www.mathnet.ru/eng/agreement>

Download details:

IP: 54.198.55.26

April 26, 2023, 16:18:48

СИБИРСКИЕ ЭЛЕКТРОННЫЕ МАТЕМАТИЧЕСКИЕ ИЗВЕСТИЯ

Siberian Electronic Mathematical Reports

<http://semr.math.nsc.ru>

Том 18, №1, стр. 338–344 (2021)

УДК 517

DOI 10.33048/semi.2021.18.023

MSC 46S10, 46B85

A NOTE ON HARDY TYPE INEQUALITIES FOR QUATERNION-VALUED FUNCTIONS

M.MUSTAFA, D.SURAGAN

ABSTRACT. In this note, we present L^p -Hardy type inequalities and Hardy-Rellich type inequalities for generalized quaternion-valued functions. As consequences, uncertainty principles and embedding theorems are obtained in the quaternion context.

Keywords: Hardy inequality, Quaternion-valued function, Dirac operator, Quaternion.

1. INTRODUCTION

In the 90s, a new class of quaternion-valued functions introduced and studied intensively by several authors, for example, our motivation to write this paper is based on [2] (see also [1]). This recent theory has been growing intensively and was developed in a series of papers.

In what follows we will work in $\mathbb{H}$, the generalized skew-field of quaternions. This means each element $x' \in \mathbb{H}$ has the following representation

$$x' = x_0 + \sum_{i=1}^n e_i x_i,$$

where $1, e_1, \dots, e_n$ are the basis elements of $\mathbb{H}$. For these elements we have the multiplication rules

$$\bullet e_1^2 = \dots = e_n^2 = -1,$$

MUSTAFA, M., SURAGAN, D., A NOTE ON HARDY TYPE INEQUALITIES FOR QUATERNION-VALUED FUNCTIONS.

© 2021 SURAGAN D., MUSTAFA M.

This research is funded by the Science Committee of the Ministry of Education and Science of the Republic of Kazakhstan (Grant No. AP09057887). The first author was supported by Nazarbayev University FDCRG 021220FD3851. The second author was supported by Nazarbayev University FDCRG 240919FD3901. Durvudkhan Suragan is the corresponding author.

Received October, 25, 2019, published March, 31, 2021.

- $e_i e_j + e_j e_i = -2\delta_{ij}$, for all $i, j = 1, \dots, n$. Here δ_{ij} is the Kronecker symbol. Obviously, in the case of $n = 4$ we get the standard skew-field of the quaternion.

The conjugate element $\bar{x}'$ is given by $\bar{x}' = x_0 - \sum_{i=1}^n e_i x_i$ and we have the property

$$|x'|^2 = x' \bar{x}' = \bar{x}' x' = x_0^2 + \sum_{i=1}^n x_i^2,$$

and let us also introduce the following quasi-norm

$$|x|_q = \sqrt{\sum_{i=1}^n x_i^2}$$

on $\mathbb{H}$, where $x = \sum_{i=1}^n e_i x_i$.

For this setting one can introduce the usual Dirac operator which factorises the n -dimensional Laplace operator:

$$Du = \sum_{i=1}^n e_i \frac{\partial u}{\partial x_i},$$

and it's conjugate operator

$$\bar{D}u = - \sum_{i=1}^n e_i \frac{\partial u}{\partial x_i}.$$

By multiplying these we have

$$D\bar{D} = \bar{D}D = \Delta_n,$$

where Δ_n is the Laplacian for functions defined over domains in $\mathbb{R}^n$.

Note that for all $\gamma \in \mathbb{R}$, and $|x|_q \neq 0$, we have the identities

$$(1.1) \quad D \left(\frac{\bar{x}}{|x|_q^\gamma} \right) = \frac{|x|_q^\gamma D\bar{x} - \bar{x} D|x|_q^\gamma}{|x|_q^{2\gamma}} = \frac{n - \gamma}{|x|_q^\gamma},$$

and

$$(1.2) \quad \bar{D} \left(\frac{x}{|x|_q^\gamma} \right) = \frac{|x|_q^\gamma \bar{D}x - x \bar{D}|x|_q^\gamma}{|x|_q^{2\gamma}} = \frac{n - \gamma}{|x|_q^\gamma},$$

since

$$D\bar{x} = \sum_{i=1}^n e_i \frac{\partial \bar{x}}{\partial x_i} = -e_1^2 - \dots - e_n^2 = n,$$

and

$$\bar{D}x = - \sum_{i=1}^n e_i \frac{\partial x}{\partial x_i} = -e_1^2 - \dots - e_n^2 = n.$$

The main goal of this short note is to prove a version of the Hardy inequality for the Dirac operator D . As a consequence, we get the uncertainty principle for generalized quaternion-valued functions. We refer a recent open access book [3] for detailed discussions on the Hardy type inequalities and closely related topics. In the present note, in the proofs we use techniques from the papers [4] and [5] by M. Ruzhansky and the second author.

In Section 2, L^p -Hardy type inequalities are obtained. In turn, these imply Hardy type inequalities for generalized quaternion-valued functions. In Section 3, we present Hardy-Rellich type inequalities and introduce some embedding results.

2. L^p -HARDY TYPE INEQUALITIES

In this section, we present L^p -Hardy type inequalities for the generalized Dirac operator. As a consequence, we obtain the uncertainty principle in terms of the generalised Dirac operator. Note that here and after $\{0\}$ denotes the set of quaternions $\{x'\}$ such that $x = 0$ for $x' \in \mathbb{H}$.

Theorem 2.1. *Let $\alpha, \beta \in \mathbb{R}$. For any $u \in C_0^\infty(\mathbb{H} \setminus \{0\})$, and for all $1 < p < \infty$ we have*

$$(2.1) \quad \frac{|n - \gamma|}{p} \left\| \frac{u}{|x|_q^{\frac{\gamma}{p}}} \right\|_{L^p(\mathbb{H})}^p \leq \left\| \frac{Du}{|x|_q^\alpha} \right\|_{L^p(\mathbb{H})} \left\| \frac{|u|^{p-1}}{|x|_q^\beta} \right\|_{L^{p'}(\mathbb{H})}, \quad \frac{1}{p} + \frac{1}{p'} = 1,$$

where $\gamma = \alpha + \beta + 1$ and $|\cdot|_q$ is the quasi-norm on $\mathbb{H}$. If $\gamma \neq n$, then the constant $\frac{|n-\gamma|}{p}$ is sharp.

Note that for the quaternion case, that is, when

$$x' = x_0 + e_1x_1 + e_2x_2 + e_3x_3,$$

we have the following inequality

$$(2.2) \quad \frac{|4 - \gamma|}{p} \left\| \frac{u}{|x|_q^{\frac{\gamma}{p}}} \right\|_{L^p(\mathbb{H})}^p \leq \left\| \frac{Du}{|x|_q^\alpha} \right\|_{L^p(\mathbb{H})} \left\| \frac{|u|^{p-1}}{|x|_q^\beta} \right\|_{L^{p'}(\mathbb{H})}.$$

Proof of Theorem 2.1. The inequality is trivial when $\gamma = n$, so we assume that $\gamma \neq n$. By using identity (1.2), the divergence theorem and Schwarz's inequality we have

$$\begin{aligned} \int_{\mathbb{H}} \frac{|u|^p}{|x|_q^\gamma} dx' &= \frac{1}{n - \gamma} \int_{\mathbb{H}} |u|^p \overline{D} \left(\frac{x}{|x|_q^\gamma} \right) dx' \\ &= -\frac{1}{n - \gamma} \int_{\mathbb{H}} p |u|^{p-1} \frac{x \cdot Du}{|x|_q^\gamma} dx' \\ &\leq \left| \frac{p}{n - \gamma} \right| \int_{\mathbb{H}} \frac{|u|^{p-1}}{|x|_q^\gamma} |x \cdot Du| dx' \\ &\leq \left| \frac{p}{n - \gamma} \right| \int_{\mathbb{H}} \frac{|u|^{p-1}}{|x|_q^{\alpha+\beta}} |Du| dx' \\ &\leq \left| \frac{p}{n - \gamma} \right| \left(\int_{\mathbb{H}} \frac{|Du|^p}{|x|_q^{\alpha p}} dx' \right)^{\frac{1}{p}} \left(\int_{\mathbb{H}} \frac{|u|^{(p-1)p'}}{|x|_q^{\beta p'}} dx' \right)^{\frac{1}{p'}}. \end{aligned}$$

In the last line, we have used the Hölder inequality. Thus, we obtain the desired inequality

$$(2.3) \quad \left| \frac{n - \gamma}{p} \right| \int_{\mathbb{H}} \frac{|u|^p}{|x|_q^\gamma} dx' \leq \left(\int_{\mathbb{H}} \frac{|Du|^p}{|x|_q^{\alpha p}} dx' \right)^{\frac{1}{p}} \left(\int_{\mathbb{H}} \frac{|u|^{(p-1)p'}}{|x|_q^{\beta p'}} dx' \right)^{\frac{1}{p'}}.$$

Now it remains to show that the constant is sharp. Set

$$f(x) = \begin{cases} e^{-\frac{1}{s}|x|_q^2}, & s := \alpha - \frac{\beta}{p-1} + 1 \neq 0, \\ \frac{1}{|x|_q}, & \alpha - \frac{\beta}{p-1} + 1 = 0, \end{cases}$$

then we have

$$\frac{|Df(x)|^p}{|x|_q^{\alpha p}} = \frac{|f(x)|^p}{|x|_q^{\beta p'}}$$

which satisfies the equality condition in the Hölder inequality. It implies that the constant is sharp. $\square$

Corollary 2.2. *Let $\beta = (\alpha + 1)(p - 1)$ and $\gamma = p(\alpha + 1)$. For any $u \in C_0^\infty(\mathbb{H} \setminus \{0\})$ we have*

$$(2.4) \quad \frac{|n - p(\alpha + 1)|}{p} \left\| \frac{u}{|x|_q^{\alpha+1}} \right\|_{L^p(\mathbb{H})} \leq \left\| \frac{Du}{|x|_q^\alpha} \right\|_{L^p(\mathbb{H})},$$

where $1 < p < \infty$ and $\alpha \in \mathbb{R}$.

Note that when $\alpha = 0$ and $1 < p < n$, the inequality (2.4) gives the following L^p -Hardy inequality for the Dirac operator

$$(2.5) \quad \left\| \frac{u}{|x|_q} \right\|_{L^p(\mathbb{H})} \leq \frac{p}{n - p} \|Du\|_{L^p(\mathbb{H})}.$$

Hölder's inequality and inequality 2.5 (with $p = 2$) imply the following Heisenberg-Pauli-Weyl type uncertainly principle for quaternions.

Corollary 2.3. *For each $u \in C_0^\infty(\mathbb{H} \setminus \{0\})$ we have*

$$(2.6) \quad \|u\|_{L^2(\mathbb{H})}^2 \leq \frac{1}{2} \|Du\|_{L^2(\mathbb{H})} \| |x|_q |u| \|_{L^2(\mathbb{H})}.$$

3. HARDY-RELLICH INEQUALITY

In this section, we present the Hardy-Rellich type inequality for the Dirac operator and some of its consequences.

Theorem 3.1. *Let $\alpha, \beta \in \mathbb{R}$ such that*

$$(3.1) \quad \frac{p - n}{p - 1} \leq \alpha + \beta + 1 \leq 0,$$

then for all $1 < p < \infty$ with $\frac{1}{p} + \frac{1}{p'} = 1$, we have

$$(3.2) \quad \frac{n + \gamma(p - 1) - p}{p} \left\| \frac{Du}{|x|_q^{\frac{\gamma}{p}}} \right\|_{L^p(\mathbb{H})}^p \leq \left\| \frac{D(|Du|^{p-2} \overline{Du})}{|x|_q^\alpha} \right\|_{L^p(\mathbb{H})} \left\| \frac{Du}{|x|_q^\beta} \right\|_{L^{p'}(\mathbb{H})},$$

and for any $u \in C_0^\infty(\mathbb{H} \setminus \{0\})$, where $\gamma = \alpha + \beta + 1$.

Proof. Inequality (3.2) is trivial when $\gamma = n$, so we assume that $\gamma \neq n$. By using identity (1.1), the divergence theorem and Schwarz's inequality we have

$$\begin{aligned}
\int_{\mathbb{H}} \frac{|Du|^p}{|x|_q^\gamma} dx' &= \frac{1}{n-\gamma} \int_{\mathbb{H}} |Du|^p D \left(\frac{\bar{x}}{|x|_q^\gamma} \right) dx' \\
&= -\frac{1}{n-\gamma} \int_{\mathbb{H}} \frac{p}{2} |Du|^{p-2} \frac{\bar{x} \cdot \bar{D} |Du|^2}{|x|_q^\gamma} dx' \\
(3.3) \quad &= -\frac{p}{2(n-\gamma)} \int_{\mathbb{H}} |Du|^{p-2} \frac{\bar{x} \cdot D |Du|^2}{|x|_q^\gamma} dx'.
\end{aligned}$$

Moreover, we compute

$$\begin{aligned}
&\int_{\mathbb{H}} \frac{D \left(|Du|^{p-2} \bar{D} u \right)}{|x|_q^\gamma} \bar{x} \cdot \bar{D} u dx' \\
&= - \int_{\mathbb{H}} |Du|^{p-2} Du \cdot D \left(\frac{\bar{x} \cdot \bar{D} u}{|x|_q^\gamma} \right) dx' \\
&= - \int_{\mathbb{H}} |Du|^{p-2} \left(\frac{|Du|^2}{|x|_q^\gamma} + \frac{\bar{x} \cdot \bar{D} |Du|^2}{2|x|_q^\gamma} - \frac{\gamma |\bar{x} \cdot \bar{D} u|^2}{|x|_q^{\gamma+2}} \right) dx'.
\end{aligned}$$

It follows that

$$\begin{aligned}
&\int \frac{|Du|^{p-2}}{|x|_q^\gamma} \bar{x} \cdot \bar{D} |Du|^2 dx' \\
&= 2\gamma \int_{\mathbb{H}} |Du|^{p-2} \frac{|\bar{x} \cdot \bar{D} u|^2}{|x|_q^{\gamma+2}} dx' - 2 \int_{\mathbb{H}} \frac{|Du|^p}{|x|_q^\gamma} dx' \\
&\quad - 2 \int_{\mathbb{H}} \frac{D \left(|Du|^{p-2} \bar{D} u \right)}{|x|_q^\gamma} \bar{x} \cdot \bar{D} u dx'.
\end{aligned}$$

Combining this with (3.3), we obtain

$$\begin{aligned}
\int_{\mathbb{H}} \frac{|Du|^p}{|x|_q^\gamma} dx' &= \frac{p\gamma}{\gamma-n} \int_{\mathbb{H}} |Du|^{p-2} \frac{|\bar{x} \cdot \bar{D} u|^2}{|x|_q^{\gamma+2}} dx' \\
&\quad - \frac{p}{\gamma-n} \int_{\mathbb{H}} \frac{|Du|^p}{|x|_q^\gamma} dx' - \frac{p}{\gamma-n} \int_{\mathbb{H}} \frac{D \left(|Du|^{p-2} \bar{D} u \right)}{|x|_q^\gamma} \bar{x} \cdot \bar{D} u dx'.
\end{aligned}$$

Thus, we get

$$\begin{aligned}
&\int_{\mathbb{H}} \frac{D \left(|Du|^{p-2} \bar{D} u \right)}{|x|_q^\gamma} \bar{x} \cdot \bar{D} u dx' \\
&= \frac{n-p-\gamma}{p} \int_{\mathbb{H}} \frac{|Du|^p}{|x|_q^\gamma} dx' + \gamma \int_{\mathbb{H}} |Du|^{p-2} \frac{|\bar{x} \cdot \bar{D} u|^2}{|x|_q^{\gamma+2}} dx'.
\end{aligned}$$

Now we apply the Schwarz inequality to the second term on the right hand side since from the assumption we have $\gamma = \alpha + \beta + 1 \leq 0$, so we derive

$$\begin{aligned}
\int_{\mathbb{H}} \frac{D \left(|Du|^{p-2} \overline{Du} \right)}{|x|_q^\gamma} \cdot Du \, dx' &= \frac{n-p-\gamma}{p} \int_{\mathbb{H}} \frac{|Du|^p}{|x|_q^\gamma} \, dx' + \gamma \int_{\mathbb{H}} |Du|^{p-2} \frac{|\overline{x} \cdot \overline{Du}|^2}{|x|_q^{\gamma+2}} \, dx' \\
&\geq \frac{n-p-\gamma}{p} \int_{\mathbb{H}} \frac{|Du|^p}{|x|_q^\gamma} \, dx' + \gamma \int_{\mathbb{H}} \frac{|Du|^p}{|x|_q^\gamma} \, dx' \\
(3.4) \qquad \qquad \qquad &\geq \frac{n+\gamma(p-1)-p}{p} \int_{\mathbb{H}} \frac{|Du|^p}{|x|_q^\gamma} \, dx'
\end{aligned}$$

We also have

$$\begin{aligned}
\int_{\mathbb{H}} \frac{D \left(|Du|^{p-2} \overline{Du} \right)}{|x|_q^\gamma} \overline{x} \cdot \overline{Du} \, dx' &\leq \int_{\mathbb{H}} \frac{D \left(|Du|^{p-2} \overline{Du} \right)}{|x|_q^{\gamma-1}} |Du| \, dx' \\
(3.5) \qquad \qquad \qquad &\leq \left(\int_{\mathbb{H}} \left| \frac{D \left(|Du|^{p-2} \overline{Du} \right)}{|x|_q^\alpha} \right|^p \, dx' \right)^{\frac{1}{p}} \left(\int_{\mathbb{H}} \left| \frac{Du}{|x|_q^\beta} \right|^{p'} \, dx' \right)^{\frac{1}{p'}}
\end{aligned}$$

Here we have used Schwarz's inequality and Hölder's inequality.

Combining inequalities (3.4) and (3.5), we arrive at the desired inequality. $\square$

Let us define the following Sobolev type spaces for the quaternion-valued functions when $p = 2$:

- Let $W_\gamma^{1,2}(\mathbb{H})$ be the completion of $C_0^\infty(\mathbb{H} \setminus \{0\})$ with respect to norm

$$\|u\|_{W_\gamma^{1,2}(\mathbb{H})} = \left(\int_{\mathbb{H}} \frac{|Du|^2}{|x|_q^{2\gamma}} \, dx' \right)^{\frac{1}{2}}.$$

- Let $W_\gamma^{2,2}(\mathbb{H})$ be the completion of $C_0^\infty(\mathbb{H} \setminus \{0\})$ with respect to norm

$$\|u\|_{W_\gamma^{2,2}(\mathbb{H})} = \left(\int_{\mathbb{H}} \frac{|\Delta_n u|^2}{|x|_q^{2\gamma}} \, dx' \right)^{\frac{1}{2}}.$$

- Let $H_{\alpha,\beta}^2(\mathbb{H})$ be the completion of $C_0^\infty(\mathbb{H} \setminus \{0\})$ with respect to norm

$$\|u\|_{H_{\alpha,\beta}^2(\mathbb{H})} = \left(\int_{\mathbb{H}} \frac{|Du|^2}{|x|_q^{2\beta}} + \frac{|\Delta_n u|^2}{|x|_q^{2\alpha}} \, dx' \right)^{\frac{1}{2}}.$$

Theorem 3.2. *For $\alpha, \beta \in \mathbb{R}$ we have the following continuous embedding*

$$(3.6) \qquad H_{\alpha,\beta}^2(\mathbb{H}) \subset W_{\frac{\alpha+\beta-1}{2}}^{2,2}(\mathbb{H}),$$

for $\alpha + \beta - 1 \neq n$, and

$$(3.7) \qquad W_\alpha^{2,2}(\mathbb{H}) \subset W_{\alpha+1}^{1,2}(\mathbb{H}),$$

for $\alpha \leq \frac{n}{2} - 2$ with $\alpha \neq \frac{n}{2}$.

Proof. Since $\alpha + \beta - 1 \neq n$ from inequality (3.2) with $p = 2$ we have

$$\begin{aligned} \left\| \frac{Du}{|x|^{\frac{\alpha+\beta-1}{2}}} \right\|_{L^2(\mathbb{H})}^2 &\leq \frac{2}{|n - \alpha - \beta + 1|} \left\| \frac{\Delta_n u}{|x|^\alpha} \right\|_{L^2(\mathbb{H})} \left\| \frac{|Du|}{|x|^\beta} \right\|_{L^2(\mathbb{H})} \\ &\leq \frac{2}{|n - \alpha - \beta + 1|} \left(\int_{\mathbb{H}} \frac{|Du|^2}{|x|^{2\beta}} + \frac{|\Delta_n u|^2}{|x|^{2\alpha}} dx' \right), \end{aligned}$$

for all $u \in C_0^\infty(\mathbb{H} \setminus \{0\})$. Here we have used the Cauchy-Bunyakovsky-Schwarz inequality. This proves (3.6). Embedding (3.7) also implies from inequality (3.2) with $p = 2$, namely assuming $\beta = \alpha + 1$ with $\alpha \leq \frac{n}{2} - 2$ and $\alpha \neq \frac{n}{2}$. $\square$

REFERENCES

- [1] R. Aulaskari, J. Xiao, R. Zhao, *On subspaces and subsets of BMOA and UBC*, Analysis, **15**:2 (1995), 101–121. Zbl 0835.30027
- [2] K. Gürlebeck, U. Kähler, M.V. Shapiro, L.M. Tovar, *On Q_p -spaces of quaternion-valued functions*, Complex Variables, Theory Appl., **39**:2 (1999), 115–135. Zbl 1043.30030
- [3] M. Ruzhansky, D. Suragan, *Hardy inequalities on homogeneous groups*, Progress in Mathematics, **327**, Cham, Birkhäuser, 2019. Zbl 1428.22011
- [4] M. Ruzhansky, D. Suragan, *On horizontal Hardy, Rellich, Caffarelli-Kohn-Nirenberg and p -sub-Laplacian inequalities on stratified groups*. J. Differ. Equations, **262**:3 (2017), 1799–1821. Zbl 1358.22003
- [5] M. Ruzhansky, D. Suragan, *Hardy and Rellich inequalities, identities, and sharp remainders on homogeneous groups*. Adv. Math., **317** (2017), 799–822. Zbl 1376.22009

MANAT MUSTAFA

DEPARTMENT OF MATHEMATICS, NAZARBAYEV UNIVERSITY

53, QABANBAYBATYR AVE.,

NUR-SULTAN, 010000, KAZAKHSTAN

Email address: manat.mustafa@nu.edu.kz

DURVUDKHAN SURAGAN

DEPARTMENT OF MATHEMATICS, NAZARBAYEV UNIVERSITY

53, QABANBAYBATYR AVE.,

NUR-SULTAN, 010000, KAZAKHSTAN

Email address: durvudkhan.suragan@nu.edu.kz