Math-Net.Ru

Общероссийский математический портал

I. Jana, K. Saha, A. B. Soshnikov, Fluctuations of linear eigenvalue statistics of random band matrices, *Теория вероятн. и ее примен.*, 2015, том 60, выпуск 3, 553–592

DOI: <https://doi.org/10.4213/tvp4637>

Использование Общероссийского математического портала Math-Net.Ru подразумевает, что вы прочитали и согласны с пользовательским соглашением
<http://www.mathnet.ru/rus/agreement>

Параметры загрузки:

IP: 3.91.87.62

26 апреля 2023 г., 16:17:39

© 2015 г.

JANA I.^{*}, SAHA K.^{**}, SOSHNIKOV A.^{***}

FLUCTUATIONS OF LINEAR EIGENVALUE STATISTICS OF RANDOM BAND MATRICES

В этой статье мы изучаем флуктуацию линейных статистик собственных значений случайных ленточных матриц $M_n = W_n/\sqrt{b_n}$, где W_n есть $n \times n$ эрмитова случайная матрица с ленточной длиной b_n , то есть только диагональные и первые b_n вне-диагональные матричные элементы не равны нулю. Мы изучаем линейные статистики $\mathcal{N}(\phi) = \sum_{i=1}^n \phi(\lambda_i)$, где λ_i — собственные значения M_n и ϕ достаточно гладкая тест-функция. Мы доказываем, что $\sqrt{b_n/n}[\mathcal{N}(\phi) - \mathbb{E}\mathcal{N}(\phi)] \xrightarrow{d} N(0, V(\phi))$ при $b_n \gg \sqrt{n}$, где выражение для $V(\phi)$ даётся в Теореме 1.

Ключевые слова и фразы: случайная ленточная матрица, центральная предельная теорема, нормальное распределение, линейные статистики собственных значений, полукруговой закон Вигнера, случайные матрицы Вигнера.

1. Introduction. Random matrix theory was developed from several different sources in the early 20th century. It is used as an important mathematical tool in various fields namely, mathematics, physics, wireless communication engineering, etc. One of the earliest example of a random matrix appeared in the study of sample covariance estimation was done by John Wishart [21]. In the early 1950s, Wigner introduced random matrix ensemble to study the energy spectra of heavy atoms undergoing slow nuclear reactions.

Random matrices are also used to model wireless channels. A random matrix model of CDMA networks can be found in [17], [18].

A special kind of random matrix ensemble is a random band matrix. In 1955, Wigner studied the matrices H of the form $H = K + V$, where K is an $n \times n$ diagonal matrix consisting of $\dots, -2, -1, 0, 1, 2, \dots$, and V is

^{*}Department of Mathematics, University of California, Davis, CA; e-mail: ijana@math.ucdavis.edu

^{**}Department of Mathematics, Indian Institute of Technology, Bombay, Mumbai, India; ksaha@math.iitb.ac.in

^{***}Department of Mathematics, University of California, Davis, CA; soshniko@math.ucdavis.edu

¹⁾ Research supported by INSPIRE Fellowship, Dept. of Science and Technology, Govt. of India and VIMSS fellowship.

an $n \times n$ symmetric sign matrix having nonvanishing elements only up to a distance b_n from the main diagonal. Such a matrix H was called as bordered matrix [19], [20].

Another treatment of random band matrix was done by G. Casati et al. [4], [3] in the context of quantum chaos. They studied $n \times n$ symmetric random band matrices of bandwidth b_n , where b_n grows with n . In 1992, Molchanov et al. proved the semicircle law for random band matrices [12]. In 1991, Fyodorov and Mirlin proved that b_n^2/n is a crucial parameter for random band matrices [8], [11]. Numerical simulations show that the local eigenvalue statistics changes from Poisson to GOE or GUE as b_n changes from $b_n \ll \sqrt{n}$ to $b_n \gg \sqrt{n}$. Recently, Li and Soshnikov [9] proved the central limit theorem (CLT) for linear statistics of eigenvalues of band random matrices when the bandwidth b_n satisfies $\sqrt{n} \ll b_n \ll n$. In this article we write $\alpha_n \ll \beta_n$ if $\alpha_n/\beta_n \rightarrow 0$ as $n \rightarrow \infty$, and we write $\alpha_n = O(\beta_n)$ if $|\alpha_n/\beta_n| \leq C$ for all n for some constant $C > 0$.

In this article, we deal with the CLT for the eigenvalue statistics of band random matrices. We take the approach of M. Shcherbina in [15] to establish the CLT for band matrices with bandwidth b_n where $b_n \rightarrow \infty$ as $n \rightarrow \infty$. We give an alternative proof of Li and Soshnikov [9] result on CLT of band matrices when $\sqrt{n} \ll b_n \ll n$. We have given some simulation results in Section 5, which ensure that the CLT for band matrices will also hold if $\sqrt{n}/b_n \rightarrow 0$ and $b_n \rightarrow \infty$.

Now we define our model. Let us define the (circular) distance function $d_n: \mathbb{N} \times \mathbb{N} \rightarrow \mathbb{N}$ as

$$d_n(j, k) := \min\{|j - k|, n - |j - k|\},$$

and the index sets $I_n, I_n^+ \subset \mathbb{N} \times \mathbb{N}$, $I_1 \subset \mathbb{N}$ as

$$\begin{aligned} I_n &:= \{(j, k): d_n(j, k) \leq b_n\}, \quad I_n^+ = \{(j, k): (j, k) \in I_n, j \leq k\}, \\ I_1 &= \{1 < j \leq n: (1, j) \in I_n\}, \end{aligned} \quad (1)$$

where $\{b_n\}$ is a sequence of positive integers such that $b_n \rightarrow \infty$ as $n \rightarrow \infty$.

Define a real symmetric random band matrix $M = (m_{jk})_{n \times n}$ of bandwidth b_n as

$$m_{jk} = m_{kj} = \begin{cases} b_n^{-1/2} w_{jk} & \text{if } d_n(j, k) \leq b_n, \\ 0 & \text{otherwise,} \end{cases} \quad (2)$$

where $\{w_{ii}\}$ and $\{w_{jk}\}_{j \neq k, (j, k) \in I_n^+}$ are two sets of independent identically distributed (i.i.d.) real random variables with

$$\mathbb{E}[w_{jk}] = 0, \quad \mathbb{E}[w_{jk}^2] = \begin{cases} 1 & \text{if } j \neq k, \\ \sigma^2 & \text{if } j = k. \end{cases} \quad (3)$$

Here $\{w_{jk}\}$ may depend on n , but we suppress it when there is no confusion. Let $\lambda_1 \leq \lambda_2 \leq \dots \leq \lambda_n$ be the eigenvalues of the random band matrix M . Define the linear eigenvalue statistic of the eigenvalues of M as

$$\mathcal{N}_n(\phi) = \sum_{i=1}^n \phi(\lambda_i), \quad (4)$$

and the normalized eigenvalue statistic of the matrix M as

$$\mathcal{M}_n(\phi) = \sqrt{\frac{b_n}{n}} \mathcal{N}_n(\phi), \quad (5)$$

where ϕ is a test function.

2. Main results.

Theorem 1. *Let M be a real symmetric random band matrix as defined in (2), and b_n be a sequence of integers satisfying $\sqrt{n} \ll b_n \ll n$. Assume the following:*

(i) w_{jk} satisfies the Poincaré inequality with constant $m > 0$ not depending on j, k, n , i.e., for any continuously differentiable function f ,

$$\mathbf{D}(f(w_{jk})) \leq \frac{1}{m} \mathbb{E} \left[|f'(w_{jk})|^2 \right].$$

(ii) $\mathbb{E}[w_{jk}^4] = \mu_4$ for all $j \neq k$ and $d_n(j, k) \leq b_n$.

(iii) $\phi: \mathbb{R} \rightarrow \mathbb{R}$ is a test function in the Sobolev space H^s , i.e., $\|\phi\|_s < \infty$, where

$$\|\phi\|_s^2 = \int_{\mathbb{R}} (1 + 2|t|)^{2s} |\hat{\phi}(t)|^2 dt, \quad \hat{\phi}(t) = \frac{1}{\sqrt{2\pi}} \int_{\mathbb{R}} e^{-it\lambda} \phi(\lambda) d\lambda,$$

and $s > 5/2$.

Then the centred normalized eigenvalue statistic $\mathcal{M}^\circ(\phi) = \mathcal{M}_n(\phi) - \mathbb{E}[\mathcal{M}_n(\phi)]$ converges in distribution to the Gaussian random variable with mean zero and variance given by

$$\begin{aligned} V(\phi) &= \frac{\kappa_4}{16\pi^2} \left(\int_{-2\sqrt{2}}^{2\sqrt{2}} \frac{4 - \mu^2}{\sqrt{8 - \mu^2}} \phi(\mu) d\mu \right)^2 + \frac{\sigma^2}{16\pi^2} \left(\int_{-2\sqrt{2}}^{2\sqrt{2}} \frac{\mu \phi(\mu)}{\sqrt{8 - \mu^2}} d\mu \right)^2 \\ &\quad + \int_{-2\sqrt{2}}^{2\sqrt{2}} \int_{-2\sqrt{2}}^{2\sqrt{2}} \sqrt{(8 - x^2)(8 - y^2)} F(x, y) \\ &\quad \times \int_{-2\sqrt{2}}^{2\sqrt{2}} \int_{-2\sqrt{2}}^{2\sqrt{2}} \frac{\mu_1 \phi(\mu_1)}{(x - \mu_1) \sqrt{8 - \mu_1^2}} \frac{\mu_2 \phi(\mu_2)}{(x - \mu_2) \sqrt{8 - \mu_2^2}} d\mu_1 d\mu_2 dx dy, \end{aligned}$$

where for $x \neq y$

$$F(x, y) = 2 \int_{-\infty}^{\infty} \frac{s^3 \sin s - s \sin^3 s}{2(s^2 - \sin^2 s)^2 - (s^3 \sin s + s \sin^3 s)xy + s^2 \sin^2 s(x^2 + y^2)} ds,$$

and κ_4 is the fourth cumulant of the off-diagonal entries, i.e., $\kappa_4 = \mu_4 - 3$.

3. Proof of Theorem 1. We will follow the approach taken by M. Shcherbina in [15] for full (Wigner) matrix. This approach is based on two main ideas. The first ingredient is stated in the following proposition which gives a bound on the variance of linear eigenvalue statistics with a sufficiently smooth test function in term of the variance of the trace of the resolvent of a random matrix. For a proof of this result see [15], [13]. In what follows, we denote $X^\circ = X - \mathbb{E}[X]$ for any random variable X .

Proposition 1. *Let M be an $n \times n$ real symmetric random matrix and $\mathcal{N}_n(\phi)$ be a linear eigenvalue statistic of its eigenvalue as in (4). Then for any $s > 0$ we have*

$$\mathbf{D}[\mathcal{N}_n(\phi)] \leq C_s \|\phi\|_s^2 \int_0^\infty dy e^{-y} y^{2s-1} \int_{-\infty}^\infty \mathbf{D}[\mathrm{Tr}(G(x+iy))] dx,$$

where C_s is a constant depends only on s , and $G(z) = (M - zI)^{-1}$ is the resolvent of the matrix M .

The second ingredient of this approach is to use the martingale difference technique to provide a good bound on $\mathbf{D}(\gamma_n)$, where γ_n is the trace of the resolvent of a matrix. The following proposition gives that bound.

Proposition 2. *Consider the symmetric band matrix M defined in (2) and assume (3) is satisfied. Then for some $C > 0$ not depending on z, n we have*

$$\mathbf{D}\{\gamma_n\} \leq \frac{Cn}{b_n} (y^{-2} + y^{-4}) \left(\max \left\{ y, |x| - \frac{2}{y} \right\} \right)^{-2}, \quad (6)$$

where $\gamma_n = \mathrm{Tr}(M - zI)^{-1} = \mathrm{Tr}(G)$ and $z = x + iy$, $y > 0$.

We prove this result in the appendix section. Now we outline the proof of Theorem 1.

P r o o f o f T h e o r e m 1. By Lévy's continuity theorem, it suffices to show that if

$$Z_n(x) = \mathbb{E}[e_n(x)], \quad e_n(x) = e^{ix \cdot \mathcal{M}_n^\circ(\phi)}, \quad (7)$$

then for each $x \in \mathbb{R}$

$$\lim_{n \rightarrow \infty} Z_n(x) = \exp \left[-\frac{x^2 V(\phi)}{2} \right]$$

with $V(\phi)$ as in Theorem 1. For any test function $\phi \in H^s$, define

$$\phi_\eta = P_\eta * \phi,$$

where P_η is the Poisson kernel given by

$$P_\eta(x) = \frac{\eta}{\pi(x^2 + \eta^2)}.$$

We know that ϕ_η approximates ϕ in the H^s norm, i.e.,

$$\lim_{\eta \rightarrow 0} \|\phi - \phi_\eta\|_s \rightarrow 0. \quad (8)$$

For the moment, we denote the characteristic function defined in (7), by $Z_n(\phi)$ (to make its dependence on ϕ clear). Then we have

$$\lim_{n \rightarrow \infty} Z_n(\phi) = \lim_{\eta \downarrow 0} \lim_{n \rightarrow \infty} (Z_n(\phi) - Z_n(\phi_\eta)) + \lim_{\eta \downarrow 0} \lim_{n \rightarrow \infty} Z_n(\phi_\eta).$$

Now using the Proposition 1 and (8), we shall show that

$$\lim_{\eta \downarrow 0} \lim_{n \rightarrow \infty} (Z_n(\phi) - Z_n(\phi_\eta)) = 0 \quad (9)$$

and then $\lim_{n \rightarrow \infty} Z_n(\phi) = \lim_{\eta \downarrow 0} \lim_{n \rightarrow \infty} Z_n(\phi_\eta)$. Hence it suffices to find the limit of

$$Z_{\eta,n} := Z_n(\phi_\eta) = \mathbb{E}[e_{\eta,n}(x)] \quad (10)$$

with

$$e_{\eta,n}(x) = \exp[ix\mathcal{M}_n^\circ(\phi_\eta)]$$

as $n \rightarrow \infty$ and $\eta \downarrow 0$ uniformly in n . Proofs of (9) and (10) are given in the next two subsections and that will complete the proof of this theorem.

3.1. Proof of equation (9). First observe that

$$\begin{aligned} |Z_n(\phi) - Z_n(\phi_\eta)|^2 &\leq 2|x|^2 \mathbf{D}[\mathcal{M}_n(\phi) - \mathcal{M}_n(\phi_\eta)] \\ &\leq 2|x|^2 \frac{b_n}{n} \mathbf{D}[\mathcal{N}_n(\phi) - \mathcal{N}_n(\phi_\eta)]. \end{aligned} \quad (11)$$

Now, in view of Proposition 1, to bound $\mathbf{D}[\mathcal{N}_n(\phi) - \mathcal{N}_n(\phi_\eta)]$ we need to estimate

$$\int_{-\infty}^{\infty} \mathbf{D}(\gamma_n(x + iy)) \, dx,$$

where $\gamma_n(x + iy) = \text{Tr}(G(x + iy))$ and $G(z) = (M - zI)^{-1}$. We estimate that for $y > 0$

$$\begin{aligned} \int_{-\infty}^{\infty} \left(\max \left\{ y, |x| - \frac{2}{y} \right\} \right)^{-2} dx &\leq \int_{||x| - 2/y| < y} \frac{1}{y^2} dx \\ &\quad + \int_{||x| - 2/y| \geq y} (x - 2/y)^{-2} dx \\ &\leq \frac{10}{y} + 10y. \end{aligned}$$

Using the above estimate and (6), we have

$$\begin{aligned} \int_0^\infty dy e^{-y} y^{2s-1} \int_{-\infty}^\infty \mathbf{D}(\gamma_n) dx &\leq \frac{C'}{b_n} \int_0^\infty e^{-y} y^{2s-1} 4n \left(\frac{1}{y} + y \right) \left(\frac{1}{y^2} + \frac{1}{y^4} \right) dy \\ &= C \frac{n}{b_n} \int_0^\infty e^{-y} (2y^{2s-3-1} + y^{2s-1-1} + y^{2s-5-1}) dy \\ &= C \frac{n}{b_n} (\Gamma(2s-3) + \Gamma(2s-1) + \Gamma(2s-5)). \end{aligned} \quad (12)$$

If we take

$$s = \frac{5}{2} + \varepsilon, \quad \varepsilon > 0,$$

then $\Gamma(2s-3) = \Gamma(2+2\varepsilon)$, $\Gamma(2s-1) = \Gamma(4+2\varepsilon)$, and $\Gamma(2s-5) = \Gamma(2\varepsilon)$. By Proposition 1, and (12), we have

$$\mathbf{D}(\mathcal{N}_n(\phi) - \mathcal{N}_n(\phi_\eta)) \leq C(\varepsilon) \frac{n}{b_n} \|\phi - \phi_\eta\|_s.$$

Using the above estimate and (11), we have

$$\begin{aligned} |Z_n(\phi) - Z_n(\phi_\eta)|^2 &\leq 2|x|^2 \frac{b_n}{n} C(\varepsilon) \frac{n}{b_n} \|\phi - \phi_\eta\|_s \\ &= 2C(\varepsilon) |x|^2 \|\phi - \phi_\eta\|_s \rightarrow 0 \quad \text{as } \eta \rightarrow 0. \end{aligned}$$

The last limit follows from the equation (8). This completes the proof of (9).

3.2. Finding the limit of the characteristic function (10). We will be using the Lemma 1 and Lemma 2 from appendix in the proof of (10). Denote the averaging with respect to $\{w_{1i}; 1 \leq i \leq n\}$ by $\mathbb{E}_1$.

P r o o f o f (10). Using the dominated convergence theorem we have

$$\begin{aligned} \frac{d}{dx} Z_n(\phi_\eta) &= \frac{d}{dx} \mathbb{E}[e_{\eta,n}(x)] = \frac{d}{dx} \mathbb{E} \left[\exp \left(ix \sqrt{\frac{b_n}{n}} \mathcal{N}_n^\circ(\phi_\eta) \right) \right] \\ &= \mathbb{E} \left[i \sqrt{\frac{b_n}{n}} \mathcal{N}_n^\circ(\phi_\eta) e_{\eta,n}(x) \right]. \end{aligned}$$

Since by construction $\phi_\eta = P_\eta * \phi$, we have

$$\mathcal{N}_n^\circ(\phi_\eta) = \frac{1}{\pi} \int_{-\infty}^\infty \phi(\mu) \operatorname{Im} \gamma_n^\circ(z_\mu) d\mu, \quad \text{where } z_\mu = \mu + i\eta.$$

Hereinafter, we use the finiteness of $\int_{\mathbb{R}} |\phi(\mu)| d\mu$ for $\phi \in H^s$, $s > 1/2$, when changing the order of integration. For notational convenience, from now on

we will denote $e_{\eta,n}(x)$ by $e(x)$. Therefore,

$$\begin{aligned} \frac{d}{dx} Z_n(\phi_\eta) &= \mathbb{E} \left[i \sqrt{\frac{b_n}{n}} e(x) \frac{1}{\pi} \int_{-\infty}^{\infty} \phi(\mu) \operatorname{Im} \gamma_n^\circ(z_\mu) d\mu \right] \\ &= \frac{1}{2\pi} \sqrt{\frac{b_n}{n}} \int_{-\infty}^{\infty} \phi(\mu) \mathbb{E} [e(x) \operatorname{Tr} (G^\circ(z_\mu) - G^\circ(\bar{z}_\mu))] d\mu \\ &= \frac{1}{2\pi} \sqrt{\frac{b_n}{n}} \int_{-\infty}^{\infty} \phi(\mu) (Y_n(z_\mu, x) - Y_n(\bar{z}_\mu, x)) d\mu, \end{aligned}$$

where

$$\begin{aligned} Y_n(z, x) &= \mathbb{E} [e(x) \operatorname{Tr} (G^\circ(z))] = \mathbb{E} [e^\circ(x) \operatorname{Tr} (G(z))] = n \mathbb{E} [G_{11}(z) e^\circ(x)] \\ &= -n \mathbb{E} [(A^{-1})^\circ e_1(x)] - n \mathbb{E} [(A^{-1})^\circ (e(x) - e_1(x))], \end{aligned} \quad (13)$$

$$e_1(x) = \exp \left[ix \sqrt{\frac{b_n}{n}} (\mathcal{N}_{n-1}(\phi_\eta))^\circ \right],$$

$$(\mathcal{N}_{n-1}(\phi_\eta))^\circ = \frac{1}{\pi} \int_{-\infty}^{\infty} \phi(\mu) \operatorname{Im} (\gamma_{n-1}(z))^\circ d\mu,$$

$$\gamma_{n-1}(z) = \operatorname{Tr} G^{(1)}(z),$$

$$A(z) = z - \frac{1}{\sqrt{b_n}} w_{11} + \langle G^{(1)} m^{(1)}, m^{(1)} \rangle, \quad (14)$$

$$m^{(1)} = \frac{1}{\sqrt{b_n}} (w_{12}, w_{13}, \dots, w_{1n})^T, \quad (15)$$

$$G^{(1)}(z) = \left(G_{ij}^{(1)}(z) \right)_{i,j=2}^n = (M^{(1)} - zI)^{-1}, \quad (16)$$

and $M^{(1)}$ is the main bottom $(n-1) \times (n-1)$ minor of M . In the above notation $\langle \cdot, \cdot \rangle$ represents the inner product of two complex vectors, i.e., $\langle x, y \rangle = \bar{y}^T x$ for $x, y \in \mathbb{C}^{n-1}$. The equation (13) follows from the Schur complement lemma, which says that

$$G_{11}(z) = \frac{1}{w_{11}/\sqrt{b_n} - z - \langle G^{(1)} m^{(1)}, m^{(1)} \rangle} = -\frac{1}{A(z)}. \quad (17)$$

Now we rewrite

$$\begin{aligned} \sqrt{\frac{b_n}{n}} Y_n(z, x) &= -\sqrt{nb_n} \mathbb{E} [(A^{-1})^\circ e_1(x)] - \sqrt{nb_n} \mathbb{E} [(A^{-1})^\circ (e(x) - e_1(x))] \\ &=: T_1 + T_2. \end{aligned} \quad (18)$$

Using Taylor expansion we have

$$A^{-1} = \frac{1}{\mathbb{E}[A]} - \frac{A^\circ}{(\mathbb{E}[A])^2} + \frac{(A^\circ)^2}{(\mathbb{E}[A])^3} - \frac{(A^\circ)^3}{(\mathbb{E}[A])^4} + \frac{(A^\circ)^4}{A(\mathbb{E}[A])^4}. \quad (19)$$

Therefore, we can estimate

$$\begin{aligned}
 T_1 &= -\sqrt{nb_n} \mathbb{E} \left[(A^{-1})^\circ e_1(x) \right] \\
 &= -\sqrt{nb_n} \mathbb{E} \left[(A^{-1}) e_1^\circ(x) \right] \\
 &= -\sqrt{nb_n} \mathbb{E} \left[\left(\frac{1}{\mathbb{E}[A]} - \frac{A^\circ}{(\mathbb{E}[A])^2} + \frac{(A^\circ)^2}{(\mathbb{E}[A])^3} \right. \right. \\
 &\quad \left. \left. - \frac{(A^\circ)^3}{(\mathbb{E}[A])^4} + \frac{(A^\circ)^4}{A(\mathbb{E}[A])^4} \right) e_1^\circ(x) \right] \\
 &= \sqrt{nb_n} \mathbb{E} \left[\left(\frac{A^\circ}{(\mathbb{E}[A])^2} - \frac{(A^\circ)^2}{(\mathbb{E}[A])^3} \right) e_1^\circ(x) \right] \\
 &\quad + \sqrt{nb_n} \mathbb{E} \left[\left(\frac{(A^\circ)^3}{(\mathbb{E}[A])^4} - \frac{(A^\circ)^4}{A(\mathbb{E}[A])^4} \right) e_1^\circ(x) \right]. \tag{20}
 \end{aligned}$$

Now we shall estimate each term individually. First of all, since M is a real symmetric matrix we have

$$\|G(z)\| \leq \frac{1}{|\operatorname{Im} z|}, \tag{21}$$

and, in particular, $1/|A| \leq 1/|\operatorname{Im} z|$. It can also be checked that $1/|\mathbb{E}[A]| \leq 1/|\operatorname{Im} z|$. Hereinafter $\|X\|$ is the spectral norm of a matrix X . Using the above equation (21) and the estimates (37), (39), we have

$$\begin{aligned}
 \left| \sqrt{nb_n} \mathbb{E} \left[\frac{(A^\circ)^4}{A(\mathbb{E}[A])^4} e_1^\circ(x) \right] \right| &\leq \frac{\sqrt{nb_n}}{|\operatorname{Im} z|^5} \mathbb{E} [| (A^\circ)^4 |] \\
 &= \frac{\sqrt{nb_n}}{|\operatorname{Im} z|^5} O(b_n^{-2}) = O \left(\sqrt{\frac{n}{b_n^3}} \right) \rightarrow 0, \\
 \left| \sqrt{nb_n} \mathbb{E} \left[\frac{(A^\circ)^3}{(\mathbb{E}[A])^4} e_1^\circ(x) \right] \right| &\leq \frac{\sqrt{nb_n}}{|\operatorname{Im} z|^4} \mathbb{E} [| (A^\circ)^3 |] \\
 &= \frac{\sqrt{nb_n}}{|\operatorname{Im} z|^4} O(b_n^{-3/2}) = O \left(\sqrt{\frac{n}{b_n^2}} \right) \rightarrow 0, \\
 \left| \sqrt{nb_n} \mathbb{E} \left[\frac{(A^\circ)^2}{(\mathbb{E}[A])^3} e_1^\circ(x) \right] \right| &\leq \frac{\sqrt{nb_n}}{|\operatorname{Im} z|^3} |\mathbb{E} [e_1^\circ(x) \mathbb{E}_1 [(A^\circ)^2]]| \\
 &\leq C \sqrt{\frac{n}{b_n}} |\mathbb{E} [e_1^\circ(x) (b_n \mathbb{E}_1 (A^\circ)^2)]| \\
 &\leq C \sqrt{\frac{n}{b_n}} [\mathbf{D}(e_1^\circ(x))]^{1/2} [\mathbf{D}(b_n \mathbb{E}_1 (A^\circ)^2)]^{1/2} \\
 &\leq C \sqrt{\frac{n}{b_n}} O(b_n^{-1/2}) \\
 &= O \left(\sqrt{\frac{n}{b_n^2}} \right) \rightarrow 0 \quad \text{as } n \rightarrow \infty.
 \end{aligned}$$

Therefore, we have

$$\begin{aligned} T_1 &= \frac{\sqrt{nb_n}}{(\mathbb{E}[A])^2} \mathbb{E}[A^\circ e_1^\circ(x)] + O\left(\sqrt{\frac{n}{b_n^2}}\right) \\ &= \frac{\sqrt{nb_n}}{(\mathbb{E}[A])^2} \mathbb{E}[e_1^\circ(x) \mathbb{E}_1(A^\circ)] + O\left(\sqrt{\frac{n}{b_n^2}}\right). \end{aligned}$$

Now

$$A^\circ = -\frac{1}{\sqrt{b_n}} w_{11} + \frac{1}{b_n} \sum_{\substack{i \neq j \\ i, j \in I_1}} G_{ij}^{(1)} w_{1i} w_{1j} + \frac{1}{b_n} \sum_{i \in I_1} (G_{ii}^{(1)} w_{1i}^2 - \mathbb{E}[G_{ii}^{(1)}]),$$

where $I_1 = \{1 < j \leq n: (1, j) \in I_n\}$.

Therefore,

$$\mathbb{E}_1[A^\circ(z)] = \frac{1}{b_n} \sum_{i \in I_1} (G_{ii}^{(1)} - \mathbb{E}[G_{ii}^{(1)}])$$

and hence

$$\begin{aligned} T_1 &= \frac{\sqrt{nb_n}}{(\mathbb{E}[A])^2} \mathbb{E}[e_1^\circ(x) \mathbb{E}_1(A^\circ)] + O\left(\sqrt{\frac{n}{b_n^2}}\right) \\ &= \frac{\sqrt{nb_n}}{(\mathbb{E}[A])^2} \mathbb{E}\left[e_1^\circ(x) \frac{1}{b_n} \sum_{i \in I_1} (G_{ii}^{(1)} - \mathbb{E}[G_{ii}^{(1)}])\right] + O\left(\sqrt{\frac{n}{b_n^2}}\right) \\ &= \frac{\sqrt{nb_n}}{(\mathbb{E}[A])^2} 2 \mathbb{E}[(G_{22}^{(1)})^\circ e_1^\circ(x)] + O\left(\sqrt{\frac{n}{b_n^2}}\right) \\ &= \frac{\sqrt{nb_n}}{(\mathbb{E}[A])^2} \frac{2}{n} \mathbb{E}[\gamma_{n-1}^\circ e_1^\circ(x)] + O\left(\sqrt{\frac{n}{b_n^2}}\right) \\ &= \sqrt{\frac{b_n}{n}} \frac{2}{(\mathbb{E}[A])^2} \mathbb{E}[\gamma_{n-1}^\circ e_1(x)] + O\left(\sqrt{\frac{n}{b_n^2}}\right). \end{aligned} \tag{22}$$

Hereinafter, all bounds (implicitly) depending on z hold uniformly on the set $\{\mu + i\eta: \mu \in \mathbb{R}\}$ for any given $\eta > 0$. Now

$$\begin{aligned} &|\mathbb{E}[\gamma_{n-1}^\circ e_1(x)] - \mathbb{E}[\gamma_n^\circ e(x)]| \\ &= |\mathbb{E}[\gamma_{n-1}^\circ e_1(x)] - \mathbb{E}[\gamma_n^\circ e_1(x)] + \mathbb{E}[\gamma_n^\circ e_1(x)] - \mathbb{E}[\gamma_n^\circ e(x)]| \\ &\leq \left(\mathbb{E}[|\gamma_{n-1}^\circ - \gamma_n^\circ|^4]\right)^{1/4} + |\mathbb{E}[\gamma_n^\circ(e_1(x) - e(x))]| \\ &= O(b_n^{-1/2}) + |\mathbb{E}[\gamma_n^\circ(e_1(x) - e(x))]|. \end{aligned}$$

The last equality follows from (40). We estimate

$$\begin{aligned}
 e(x) - e_1(x) &= \exp \left[ix \sqrt{\frac{b_n}{n}} \mathcal{N}_n^\circ(\phi_\eta) \right] - \exp \left[ix \sqrt{\frac{b_n}{n}} \mathcal{N}_{n-1}^\circ(\phi_\eta) \right] \\
 &= \left(\exp \left[ix \sqrt{\frac{b_n}{n}} \mathcal{N}_n^\circ(\phi_\eta) - ix \sqrt{\frac{b_n}{n}} \mathcal{N}_{n-1}^\circ(\phi_\eta) \right] - 1 \right) e_1(x) \\
 &= ix \sqrt{\frac{b_n}{n}} (\mathcal{N}_n^\circ(\phi_\eta) - \mathcal{N}_{n-1}^\circ(\phi_\eta)) e_1(x) \\
 &\quad + \frac{b_n}{n} O \left(x^2 (\mathcal{N}_n^\circ(\phi_\eta) - \mathcal{N}_{n-1}^\circ(\phi_\eta))^2 e_1(x) \right) \\
 &= \frac{ix}{\pi} \sqrt{\frac{b_n}{n}} \int_{-\infty}^{\infty} \left[\phi(\mu) \operatorname{Im} (\gamma_n^\circ - \gamma_{n-1}^\circ) e_1(x) \right. \\
 &\quad \left. + \sqrt{\frac{b_n}{n}} \phi(\mu) O(\gamma_n^\circ - \gamma_{n-1}^\circ)^2 \right] d\mu. \tag{23}
 \end{aligned}$$

Therefore,

$$\begin{aligned}
 \mathbb{E} [\gamma_n^\circ (e(x) - e_1(x))] &= \mathbb{E} \left[\frac{ix}{\pi} \sqrt{\frac{b_n}{n}} \int_{-\infty}^{\infty} \phi(\mu) \left[\operatorname{Im} (\gamma_n^\circ - \gamma_{n-1}^\circ) e_1(x) \gamma_n^\circ \right. \right. \\
 &\quad \left. \left. + \sqrt{\frac{b_n}{n}} \gamma_n^\circ O(\gamma_n^\circ - \gamma_{n-1}^\circ)^2 \right] d\mu \right].
 \end{aligned}$$

Using estimates (6) and (40), we have

$$\begin{aligned}
 |\mathbb{E} [\operatorname{Im} (\gamma_n^\circ - \gamma_{n-1}^\circ) e_1(x) \gamma_n^\circ]| &\leq (\mathbb{E} [|\gamma_n^\circ|^2])^{1/2} \left(\mathbb{E} [|e_1(x) \operatorname{Im} (\gamma_n^\circ - \gamma_{n-1}^\circ)|^2] \right)^{1/2} \\
 &= O \left(\sqrt{\frac{n}{b_n}} \sqrt{\frac{1}{b_n}} \right).
 \end{aligned}$$

Similarly,

$$\mathbb{E} [\gamma_n^\circ O(\gamma_n^\circ - \gamma_{n-1}^\circ)^2] = O \left(\sqrt{\frac{n}{b_n}} \frac{1}{b_n} \right).$$

Therefore,

$$|\mathbb{E} [\gamma_{n-1}^\circ e_1(x)] - \mathbb{E} [\gamma_n^\circ e(x)]| = O \left(\frac{1}{\sqrt{b_n}} \right).$$

From the equation (22) and the above estimates we have

$$T_1 = \sqrt{\frac{b_n}{n}} \frac{2}{(\mathbb{E} [A])^2} \mathbb{E} [\gamma_{n-1}^\circ e_1(x)] + O \left(\frac{\sqrt{n}}{b_n} \right)$$

$$\begin{aligned}
 &= \sqrt{\frac{b_n}{n}} \frac{2}{(\mathbb{E}[A])^2} \left[\mathbb{E}[\gamma_n^\circ e(x)] + O\left(\frac{1}{\sqrt{b_n}}\right) \right] + O\left(\frac{\sqrt{n}}{b_n}\right) \\
 &= \sqrt{\frac{b_n}{n}} \frac{2}{(\mathbb{E}[A])^2} Y_n(z, x) + O\left(\frac{\sqrt{n}}{b_n}\right). \tag{24}
 \end{aligned}$$

Now consider T_2 . Using (23) and (30) we have

$$\begin{aligned}
 T_2 &= -\sqrt{nb_n} \mathbb{E}[(A^{-1})^\circ(e(x) - e_1(x))] \\
 &= -\frac{ixb_n}{\pi} \mathbb{E}\left[(A^{-1})^\circ \int_{-\infty}^{\infty} \phi(\mu) \operatorname{Im}(\gamma_n^\circ - \gamma_{n-1}^\circ) e_1(x) d\mu\right] \\
 &\quad - \frac{1}{\pi} \sqrt{\frac{b_n^3}{n}} \mathbb{E}\left[(A^{-1})^\circ \int_{-\infty}^{\infty} \phi(\mu) O(\gamma_n^\circ - \gamma_{n-1}^\circ)^2 d\mu\right] \\
 &= -\frac{ixb_n}{\pi} \int_{-\infty}^{\infty} \phi(\mu) \mathbb{E}[e_1(x)(A^{-1})^\circ \operatorname{Im}(\gamma_n^\circ - \gamma_{n-1}^\circ)] d\mu + \sqrt{\frac{b_n^3}{n}} O\left(\frac{1}{b_n}\right) \\
 &= -\frac{ixb_n}{\pi} \int_{-\infty}^{\infty} \phi(\mu) \mathbb{E}[e_1(x)(A^{-1})^\circ \operatorname{Im}(\gamma_n^\circ - \gamma_{n-1}^\circ)] d\mu + O\left(\sqrt{\frac{b_n}{n}}\right) \\
 &= -\frac{ixb_n}{\pi} \int_{-\infty}^{\infty} \phi(\mu) \mathbb{E}[e_1(x)(A^{-1})^\circ \operatorname{Im}(\gamma_n - \gamma_{n-1})^\circ] d\mu + O\left(\sqrt{\frac{b_n}{n}}\right) \\
 &= \frac{ixb_n}{\pi} \int_{-\infty}^{\infty} \phi(\mu) \mathbb{E}\left[e_1(x)(A^{-1})^\circ(z) \operatorname{Im}\left(\frac{1+B(z_\mu)}{A(z_\mu)}\right)^\circ\right] d\mu + O\left(\sqrt{\frac{b_n}{n}}\right) \\
 &= T_{21} - T_{22} + O\left(\sqrt{\frac{b_n}{n}}\right),
 \end{aligned}$$

where $B(z) = \langle G^{(1)}G^{(1)}m^{(1)}, m^{(1)} \rangle$ and

$$\begin{aligned}
 T_{21} &= \frac{xb_n}{2\pi} \int_{-\infty}^{\infty} \phi(\mu) \mathbb{E}\left[e_1(x)(A^{-1})^\circ(z) \left(\frac{1+B(z_\mu)}{A(z_\mu)}\right)^\circ\right] d\mu, \\
 T_{22} &= \frac{xb_n}{2\pi} \int_{-\infty}^{\infty} \phi(\mu) \mathbb{E}\left[e_1(x)(A^{-1})^\circ(z) \left(\frac{1+B(z_\mu)}{A(z_\mu)}\right)^\circ\right] d\mu.
 \end{aligned}$$

Using $\operatorname{Im}\langle G^{(1)}m^{(1)}, m^{(1)} \rangle = \operatorname{Im} z \langle G^{(1)*}G^{(1)}m^{(1)}, m^{(1)} \rangle$, it can be easily verified that

$$\left| \frac{B(z)}{A(z)} \right| \leq \frac{1}{|\operatorname{Im} z|}, \quad \frac{1}{|\mathbb{E}[A(z)]|} \leq \frac{1}{|\operatorname{Im} z|}, \quad \text{and} \quad |\mathbb{E}[B(z)]| \leq \frac{2}{|\operatorname{Im} z|^2}. \tag{25}$$

Applying

$$A^{-1} = \frac{1}{\mathbb{E}[A]} - \frac{A^\circ}{(\mathbb{E}[A])^2} + \frac{(A^\circ)^2}{A(\mathbb{E}[A])^2}$$

to $A^{-1}(z)$, $A^{-1}(z_\mu)$ and using (37), we get

$$\begin{aligned}
 & b_n \mathbb{E} \left[e_1(x) (A^{-1})^\circ(z) \left(\frac{1+B(z_\mu)}{A(z_\mu)} \right)^\circ \right] \\
 &= b_n \mathbb{E} \left[e_1(x) \left\{ \frac{A^\circ(z)}{\mathbb{E}^2[A(z)]} \left(-\frac{B^\circ(z_\mu)}{\mathbb{E}[A(z_\mu)]} + \frac{(1+B(z_\mu))A^\circ(z_\mu)}{\mathbb{E}^2[A(z_\mu)]} \right. \right. \right. \\
 &\quad \left. \left. \left. - \frac{\mathbb{E}[B(z_\mu)A^\circ(z_\mu)]}{\mathbb{E}^2[A(z_\mu)]} \right) \right\} \right] + O(b_n^{-1/2}) \\
 &= b_n \mathbb{E} \left[e_1(x) \left\{ \frac{A^\circ(z)}{\mathbb{E}^2[A(z)]} \left(-\frac{B^\circ(z_\mu)}{\mathbb{E}[A(z_\mu)]} + \frac{(1+\mathbb{E}[B(z_\mu)])A^\circ(z_\mu)}{\mathbb{E}^2[A(z_\mu)]} \right. \right. \right. \\
 &\quad \left. \left. \left. + \frac{(B^\circ(z_\mu)A^\circ(z_\mu))^\circ}{\mathbb{E}^2[A(z_\mu)]} \right) \right\} \right] + O(b_n^{-1/2}) \\
 &= \frac{(1+\mathbb{E}[B(z_\mu)])}{\mathbb{E}^2[A(z)] \mathbb{E}^2[A(z_\mu)]} \mathbb{E}[e_1(x)b_n A^\circ(z)A^\circ(z_\mu)] \\
 &\quad - \frac{\mathbb{E}[e_1(x)b_n A^\circ(z)B^\circ(z_\mu)]}{\mathbb{E}^2[A(z)] \mathbb{E}[A(z_\mu)]} + O(b_n^{-1/2}) \\
 &= \frac{(1+\mathbb{E}[B(z_\mu)])}{\mathbb{E}^2[A(z)] \mathbb{E}^2[A(z_\mu)]} \mathbb{E}[e_1(x) \mathbb{E}_1(b_n A^\circ(z)A^\circ(z_\mu))] \\
 &\quad - \frac{\mathbb{E}[e_1(x) \mathbb{E}_1[b_n A^\circ(z)B^\circ(z_\mu)]]}{\mathbb{E}^2[A(z)] \mathbb{E}[A(z_\mu)]} + O(b_n^{-1/2}).
 \end{aligned}$$

Using (39), from the last expression we get

$$\begin{aligned}
 & b_n \mathbb{E} \left[e_1(x) (A^{-1})^\circ(z) \left(\frac{1+B(z_\mu)}{A(z_\mu)} \right)^\circ \right] \\
 &= \frac{(1+\mathbb{E}[B(z_\mu)])}{\mathbb{E}^2[A(z)] \mathbb{E}^2[A(z_\mu)]} \mathbb{E}[e_1(x)] \mathbb{E}[b_n A^\circ(z)A^\circ(z_\mu)] \\
 &\quad - \frac{\mathbb{E}[e_1(x)] \mathbb{E}[b_n A^\circ(z)B^\circ(z_\mu)]}{\mathbb{E}^2[A(z)] \mathbb{E}[A(z_\mu)]} + O(b_n^{-1/2}). \tag{26}
 \end{aligned}$$

Define

$$\begin{aligned}
 D_n(z, z_\mu) &= \frac{(1+\mathbb{E}[B(z_\mu)]) \mathbb{E}[b_n \mathbb{E}_1\{A^\circ(z)A^\circ(z_\mu)\}]}{\mathbb{E}^2[A(z)] \mathbb{E}^2[A(z_\mu)]} \\
 &\quad - \frac{\mathbb{E}[b_n \mathbb{E}_1\{A^\circ(z)B^\circ(z_\mu)\}]}{\mathbb{E}^2[A(z)] \mathbb{E}[A(z_\mu)]}.
 \end{aligned}$$

Also, using (23) and (40), we have

$$\begin{aligned}
 \mathbb{E}[e(x)] - \mathbb{E}[e_1(x)] &= \mathbb{E} \left[\frac{ix}{\pi} \sqrt{\frac{b_n}{n}} \int_{-\infty}^{\infty} \phi(\mu) \operatorname{Im}(\gamma_n^\circ - \gamma_{n-1}^\circ) e_1(x) d\mu \right. \\
 &\quad \left. + \frac{b_n}{n} x^2 \int_{-\infty}^{\infty} \phi(\mu) O(\gamma_n^\circ - \gamma_{n-1}^\circ)^2 d\mu \right] \\
 &= O(n^{-1/2}) + O(n^{-1}).
 \end{aligned}$$

Therefore,

$$\mathbb{E}[e_1(x)] = Z_n(\phi_\eta) + O(n^{-1/2}). \quad (27)$$

Combining (18), (24), (26), and (27), we get

$$\begin{aligned} \sqrt{\frac{b_n}{n}} Y_n(z, x) &= T_1 + T_2 \\ &= \frac{2}{\mathbb{E}^2[A]} \sqrt{\frac{b_n}{n}} Y_n(z, x) + \frac{x}{2\pi} \mathbb{E}[e_1(x)] \int_{-\infty}^{\infty} [D_n(z, z_\mu) - D_n(z, \bar{z}_\mu)] \phi(\mu) d\mu \\ &\quad + O(b_n^{-1/2}) \\ &= \frac{2}{\mathbb{E}^2[A]} \sqrt{\frac{b_n}{n}} Y_n(z, x) + \frac{x}{2\pi} Z_n(\phi_\eta) \int_{-\infty}^{\infty} [D_n(z, z_\mu) - D_n(z, \bar{z}_\mu)] \phi(\mu) d\mu \\ &\quad + o(1) \\ &\approx 2f^2(z) \tilde{Y}_n(z, x) + \frac{x}{2\pi} Z_n(\phi_\eta) \int_{-\infty}^{\infty} [D_n(z, z_\mu) - D_n(z, \bar{z}_\mu)] \phi(\mu) d\mu + o(1), \end{aligned}$$

where $\tilde{Y}_n(z, x) = \sqrt{\frac{b_n}{n}} Y_n(z, x)$. Therefore,

$$\tilde{Y}_n(z, x) = Z_n(\phi_\eta) \frac{x}{2\pi} \int_{-\infty}^{\infty} (C_n(z, z_\mu) - C_n(z, \bar{z}_\mu)) \phi(\mu) d\mu + o(1)$$

uniformly in z with $\text{Im } z = \eta$, where $C_n(z, z_\mu) = D_n(z, z_\mu)/(1 - 2f^2(z))$ and $f(z)$ is given in (41). Hence

$$\begin{aligned} \frac{d}{dx} Z_n(\phi_\eta) &= \frac{1}{2\pi} \int_{-\infty}^{\infty} \phi(\mu) \left(\tilde{Y}_n(z_\mu, x) - \tilde{Y}_n(\bar{z}_\mu, x) \right) d\mu \\ &= \frac{1}{2\pi} \int_{-\infty}^{\infty} \phi(\mu_1) \left[\frac{x}{2\pi} Z_n(\phi_\eta) \int_{-\infty}^{\infty} \phi(\mu_2) (C_n(z_{\mu_1}, z_{\mu_2}) - C_n(z_{\mu_1}, \bar{z}_{\mu_2})) d\mu_2 \right. \\ &\quad \left. - \frac{x}{2\pi} Z_n(\phi_\eta) \int_{-\infty}^{\infty} \phi(\mu_2) (C_n(\bar{z}_{\mu_1}, z_{\mu_2}) - C_n(\bar{z}_{\mu_1}, \bar{z}_{\mu_2})) d\mu_2 \right] d\mu_1 + o(1) \\ &= -\frac{x}{4\pi^2} Z_n(\phi_\eta) \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} \phi(\mu_1) \phi(\mu_2) [C_n(z_{\mu_1}, \bar{z}_{\mu_2}) + C_n(\bar{z}_{\mu_1}, z_{\mu_2}) \\ &\quad - C_n(z_{\mu_1}, z_{\mu_2}) - C_n(\bar{z}_{\mu_1}, \bar{z}_{\mu_2})] d\mu_2 d\mu_1 + o(1) \\ &= -x Z_n(\phi_\eta) V_n(\phi, \eta) + o(1). \end{aligned}$$

To find the limit of $V_n(\phi, \eta)$, we shall calculate limit of $[C_n(z_{\mu_1}, \bar{z}_{\mu_2}) + C_n(\bar{z}_{\mu_1}, z_{\mu_2}) - C_n(z_{\mu_1}, z_{\mu_2}) - C_n(\bar{z}_{\mu_1}, \bar{z}_{\mu_2})]$ as $n \rightarrow \infty$. Using (34) and (35),

$$\begin{aligned} D_n(z, z_\mu) &= \frac{(1 + \mathbb{E}[B(z_\mu)]) \mathbb{E}[b_n \mathbb{E}_1\{A^\circ(z) A^\circ(z_\mu)\}]}{\mathbb{E}^2[A(z)] \mathbb{E}^2[A(z_\mu)]} \\ &\quad - \frac{\mathbb{E}[b_n \mathbb{E}_1\{A^\circ(z) B^\circ(z_\mu)\}]}{\mathbb{E}^2[A(z)] \mathbb{E}[A(z_\mu)]} \end{aligned}$$

$$\begin{aligned}
&= \frac{1 + \mathbb{E}[B(z_\mu)]}{\mathbb{E}^2[A(z)] \mathbb{E}^2[A(z_\mu)]} \\
&\quad \times \mathbb{E} \left[\frac{2}{b_n} \sum_{i,j \in I_1} G_{ij}^{(1)}(z) G_{ij}^{(1)}(z_\mu) + \sigma^2 + \frac{\kappa_4}{b_n} \sum_{i \in I_1} G_{ii}^{(1)}(z) G_{ii}^{(1)}(z_\mu) \right. \\
&\quad \left. + \frac{1}{b_n} \widetilde{\gamma_{n-1}}(z) \widetilde{\gamma_{n-1}}(z_\mu) \right] \\
&\quad - \frac{1}{\mathbb{E}^2[A(z)] \mathbb{E}[A(z_\mu)]} \frac{d}{dz_\mu} \mathbb{E} \left[\frac{2}{b_n} \sum_{i,j \in I_1} G_{ij}^{(1)}(z) G_{ij}^{(1)}(z_\mu) \right. \\
&\quad \left. + \sigma^2 + \frac{\kappa_4}{b_n} \sum_{i \in I_1} G_{ii}^{(1)}(z) G_{ii}^{(1)}(z_\mu) + \frac{1}{b_n} \widetilde{\gamma_{n-1}}(z) \widetilde{\gamma_{n-1}}(z_\mu) \right].
\end{aligned}$$

Now using (36), we get

$$\left| \mathbb{E} \left[\frac{1}{b_n} \widetilde{\gamma_{n-1}}(z) \widetilde{\gamma_{n-1}}(z_\mu) \right] \right| \leq \frac{1}{b_n} \sqrt{\mathbf{D} \sum_{i \in I_1} G_{ii}^{(1)}} \sqrt{\mathbf{D} \sum_{i \in I_1} G_{ii}^{(1)}} = O\left(\frac{1}{b_n}\right).$$

Letting $n \rightarrow \infty$, using (42) we have

$$\begin{aligned}
\lim_{n \rightarrow \infty} D_n(z, z_\mu) &= f^2(z) f^2(z_\mu) (1 + 2f'(z_\mu)) \\
&\quad \times \left[\lim_{n \rightarrow \infty} \mathbb{E}[T_n] + \sigma^2 + \kappa_4 \lim_{n \rightarrow \infty} \frac{1}{b_n} \sum_{i \in I_1} \mathbb{E} \left[G_{ii}^{(1)}(z) G_{ii}^{(1)}(z_\mu) \right] \right] \\
&\quad + f^2(z) f(z_\mu) \frac{d}{dz_\mu} \left[\lim_{n \rightarrow \infty} \mathbb{E}[T_n] + \kappa_4 \lim_{n \rightarrow \infty} \frac{1}{b_n} \sum_{i \in I_1} \mathbb{E} \left[G_{ii}^{(1)}(z) G_{ii}^{(1)}(z_\mu) \right] \right], \quad (28)
\end{aligned}$$

where

$$T_n = \frac{2}{b_n} \sum_{i,j \in I_1} G_{ij}^{(1)}(z) G_{ij}^{(1)}(z_\mu).$$

Since $\mathbf{D}(G_{ii}) = O(1/b_n)$ (see (36)), we have

$$\begin{aligned}
\lim_{n \rightarrow \infty} \frac{1}{b_n} \sum_{i \in I_1} \mathbb{E} \left[G_{ii}^{(1)}(z) G_{ii}^{(1)}(z_\mu) \right] &= \lim_{n \rightarrow \infty} \frac{1}{b_n} \sum_{i \in I_1} \mathbb{E} \left[G_{ii}^{(1)}(z) \right] \mathbb{E} \left[G_{ii}^{(1)}(z_\mu) \right] \\
&= 2f(z) f(z_\mu).
\end{aligned}$$

We shall show in the subsection 4.1 of Appendix, that

$$\lim_{n \rightarrow \infty} \mathbb{E}[T_n] = \frac{1}{4\pi^3} \int_{-2\sqrt{2}}^{2\sqrt{2}} \int_{-2\sqrt{2}}^{2\sqrt{2}} \frac{\sqrt{8-x^2} \sqrt{8-y^2}}{(x-z)(y-z_\mu)} F(x, y) \mathbf{1}_{\{x \neq y\}} dx dy, \quad (29)$$

where

$$F(x, y) = 2 \int_{-\infty}^{\infty} \frac{u - u^3}{2(1 - u^2)^2 + u^2(x^2 + y^2) - u(1 + u^2)xy} ds,$$

where $u = (\sin s)/s$. Therefore,

$$\begin{aligned} \lim_{n \rightarrow \infty} C_n(z_{\mu_1}, z_{\mu_2}) &= \frac{1}{1 - 2f^2(z_{\mu_1})} \\ &\times \left[f^2(z_{\mu_1})f^2(z_{\mu_2})(1 + 2f'(z_{\mu_2})) \lim_{n \rightarrow \infty} \mathbb{E}[T_n] + f^2(z_{\mu_1})f(z_{\mu_2}) \lim_{n \rightarrow \infty} \frac{d}{dz_{\mu_2}} \mathbb{E}[T_n] \right. \\ &\quad + \sigma^2 f^2(z_{\mu_1})f^2(z_{\mu_2})(1 + 2f'(z_{\mu_2})) \\ &\quad \left. + 2\kappa_4 \left\{ f^3(z_{\mu_1})f^3(z_{\mu_2})(1 + 2f'(z_{\mu_2})) + f^3(z_{\mu_1})f(z_{\mu_2})f'(z_{\mu_2}) \right\} \right]. \end{aligned}$$

Hence

$$\begin{aligned} V(\phi) &= \lim_{\eta \downarrow 0} \lim_{n \rightarrow \infty} V_n(\phi, \eta) \\ &= \frac{\kappa_4}{16\pi^2} \left(\int_{-2\sqrt{2}}^{2\sqrt{2}} \frac{4 - \mu^2}{\sqrt{8 - \mu^2}} \phi(\mu) d\mu \right)^2 + \frac{\sigma^2}{16\pi^2} \left(\int_{-2\sqrt{2}}^{2\sqrt{2}} \frac{\mu \phi(\mu)}{\sqrt{8 - \mu^2}} d\mu \right)^2 \\ &\quad + \int_{-2\sqrt{2}}^{2\sqrt{2}} \int_{-2\sqrt{2}}^{2\sqrt{2}} \sqrt{(8 - x^2)(8 - y^2)} F(x, y) \\ &\quad \times \int_{-2\sqrt{2}}^{2\sqrt{2}} \int_{-2\sqrt{2}}^{2\sqrt{2}} \frac{\mu_1 \phi(\mu_1)}{(x - \mu_1)\sqrt{8 - \mu_1^2}} \frac{\mu_2 \phi(\mu_2)}{(x - \mu_2)^2 \sqrt{8 - \mu_2^2}} d\mu_1 d\mu_2 dx dy. \end{aligned}$$

This completes the proof of (10) and the proof of Theorem 1.

4. Appendix.

P r o o f o f P r o p o s i t i o n 2. Denote the averaging with respect to $\{w_{ij}; 1 \leq i \leq k \text{ or } 1 \leq j \leq n\}$ by $\mathbb{E}_{\leq k}$ and the averaging with respect to $\{w_{kj}; 1 \leq j \leq n\}$ by $\mathbb{E}_k$. Using the martingale difference technique (see [6]), we have

$$\begin{aligned} \mathbf{D}\{\gamma_n\} &\leq \sum_{k=1}^n \mathbb{E} \left[|\mathbb{E}_{\leq k-1}[\gamma_n] - \mathbb{E}_{\leq k}[\gamma_n]|^2 \right] = \sum_{k=1}^n \mathbb{E} \left[\left| \mathbb{E}_{\leq k-1}[\gamma_n - \mathbb{E}_{\leq k}[\gamma_n]] \right|^2 \right] \\ &\leq \sum_{k=1}^n \mathbb{E} \left[\mathbb{E}_{\leq k-1} |\gamma_n - \mathbb{E}_k[\gamma_n]|^2 \right] = \sum_{k=1}^n \mathbb{E} \left[|\gamma_n - \mathbb{E}_k[\gamma_n]|^2 \right]. \end{aligned}$$

Note that

$$\begin{aligned} \mathbb{E} \left[|\gamma_n - \mathbb{E}_1[\gamma_n]|^2 \right] &= \mathbb{E} \left[|\text{Tr}(G) - \mathbb{E}_1[\text{Tr}(G)]|^2 \right] \\ &= \mathbb{E} \left[\left| \text{Tr}(G) - \mathbb{E}_1[\text{Tr}(G)] + \text{Tr}(G^{(1)}) - \text{Tr}(G^{(1)}) \right|^2 \right] \\ &= \mathbb{E} \left[\left| \text{Tr}(G - G^{(1)}) - \mathbb{E}_1[\text{Tr}(G - G^{(1)})] \right|^2 \right]. \end{aligned}$$

From (32) we have

$$\text{Tr}(G - G^{(1)}) = -\frac{1 + B(z)}{A(z)}, \quad (30)$$

where $A(z) = -G_{11}^{-1}$, $B(z) = \langle G^{(1)}G^{(1)}m^{(1)}, m^{(1)} \rangle$, and $G^{(1)}$ is defined in (16), and $m^{(1)} = (1/\sqrt{b_n})(w_{12}, w_{13}, \dots, w_{1n})^T$. Indeed,

$$\begin{aligned} \mathbb{E} \left[|\gamma_n - \mathbb{E}_1[\gamma_n]|^2 \right] &\leq \mathbb{E} \left[\left| \frac{1+B(z)}{A(z)} - \mathbb{E}_1 \left[\frac{1+B(z)}{A(z)} \right] \right|^2 \right] \\ &\leq 2 \mathbb{E} \left[\left| \frac{1}{A(z)} - \mathbb{E}_1 \left[\frac{1}{A(z)} \right] \right|^2 \right] + 2 \mathbb{E} \left[\left| \frac{B(z)}{A(z)} - \mathbb{E}_1 \left[\frac{B(z)}{A(z)} \right] \right|^2 \right]. \end{aligned}$$

Now, by (21) and (25),

$$\begin{aligned} \mathbb{E}_1 \left[\left| \frac{B(z)}{A(z)} - \mathbb{E}_1 \left[\frac{B(z)}{A(z)} \right] \right|^2 \right] &\leq \mathbb{E}_1 \left[\left| \frac{B(z)}{A(z)} - \frac{\mathbb{E}_1[B(z)]}{\mathbb{E}_1[A(z)]} \right|^2 \right] \\ &\leq \mathbb{E}_1 \left[\left| \frac{B_1^\circ}{\mathbb{E}_1[A]} - \frac{A_1^\circ}{\mathbb{E}_1[A]} \frac{B}{A} \right|^2 \right] \\ &\leq 2 \mathbb{E}_1 \left[\left| \frac{B_1^\circ}{\mathbb{E}_1[A]} \right|^2 \right] + \frac{2}{|\operatorname{Im} z|^2} \mathbb{E}_1 \left[\left| \frac{A_1^\circ}{\mathbb{E}_1[A]} \right|^2 \right], \end{aligned}$$

where $A_1^\circ = A - \mathbb{E}_1[A]$. So it is enough to estimate $\mathbb{E}_1[|A_1^\circ/\mathbb{E}_1[A]|^2]$ and $\mathbb{E}_1[|B_1^\circ/\mathbb{E}_1[A]|^2]$. Note that

$$\begin{aligned} A &= z - \frac{1}{\sqrt{b_n}} w_{11} + \langle G^{(1)}m^{(1)}, m^{(1)} \rangle, \\ A_1^\circ &= -\frac{1}{\sqrt{b_n}} w_{11} + \frac{1}{b_n} \sum_{\substack{i \neq j \\ i, j \in I_1}} G_{ij}^{(1)} w_{1i} w_{1j} + \frac{1}{b_n} \sum_{i \in I_1} G_{ii}^{(1)} (w_{1i}^2)^\circ. \end{aligned}$$

Therefore,

$$\begin{aligned} \mathbb{E}_1 \left[|A_1^\circ|^2 \right] &= \mathbb{E}_1 \left[\frac{1}{b_n} w_{11}^2 + \frac{1}{b_n^2} \sum_{\substack{i \neq j \\ i, j \in I_1}} G_{ij}^{(1)} w_{1i} w_{1j} \sum_{\substack{k \neq l \\ k, l \in I_1}} \overline{G_{kl}^{(1)}} w_{1k} w_{1l} \right. \\ &\quad \left. + \frac{1}{b_n^2} \sum_{i \in I_1} G_{ii}^{(1)} (w_{1i}^2)^\circ \sum_{l \in I_1} \overline{G_{ll}^{(1)}} (w_{1l}^2)^\circ \right] \\ &= \frac{\sigma^2}{b_n} + \frac{2}{b_n^2} \sum_{\substack{i \neq j \\ i, j \in I_1}} |G_{ij}^{(1)}|^2 + \frac{\mu_4 - 1}{b_n^2} \sum_{i \in I_1} |G_{ii}^{(1)}|^2 \\ &\leq \frac{\sigma^2}{b_n} + \frac{2}{b_n^2} \frac{2b_n}{|\operatorname{Im} z|^2} + \frac{\mu_4 - 1}{b_n^2} \frac{2b_n}{|\operatorname{Im} z|^2} \\ &\leq \frac{1}{b_n} \left(\sigma^2 + \frac{2 + 2\mu_4}{|\operatorname{Im} z|^2} \right). \end{aligned} \tag{31}$$

Now we want to estimate $\mathbb{E}_1[|B_1^\circ|^2]$, where $B = \langle G^{(1)}G^{(1)}m^{(1)}, m^{(1)} \rangle = \langle H^{(1)}m^{(1)}, m^{(1)} \rangle$, and $B_1^\circ = B - \mathbb{E}_1[B]$. Therefore,

$$\mathbb{E}_1[B] = \frac{1}{b_n} \sum_{i \in I_1} H_{ii}^{(1)} = \frac{1}{b_n} \sum_{i \in I_1} \sum_{j=2}^n \left(G_{ij}^{(1)} \right)^2,$$

and

$$B_1^\circ = \frac{1}{b_n} \sum_{\substack{i \neq j \\ i, j \in I_1}} H_{ij}^{(1)} w_{1i} w_{1j} + \frac{1}{b_n} \sum_{i \in I_1} H_{ii}^{(1)} (w_{1i}^2)^\circ.$$

Let us call $C_0 = \mathbb{E}[(w_{1i}^2)^\circ]^2$. Then

$$\begin{aligned} \mathbb{E}_1[|B_1^\circ|^2] &= \frac{1}{b_n^2} \sum_{\substack{i \neq j \\ i, j \in I_1}} |H_{ij}^{(1)}|^2 + \frac{C_0}{b_n^2} \sum_{i \in I_1} |H_{ii}^{(1)}|^2 \\ &= \frac{1}{b_n^2} \sum_{\substack{i \neq j \\ i, j \in I_1}} \left| \sum_{k=2}^n G_{ik}^{(1)} G_{kj}^{(1)} \right|^2 + \frac{C_0}{b_n^2} \sum_{i \in I_1} \left| \sum_{k=2}^n G_{ik}^{(1)} G_{ki}^{(1)} \right|^2 \\ &\leq \frac{1}{b_n^2} \sum_{i \in I_1} \|(G^{(1)})^2\|^2 + \frac{C_0}{b_n^2} 2b_n \|(G^{(1)})^2\|^2 \\ &= \frac{2}{b_n} \frac{1}{|\operatorname{Im} z|^4} + \frac{2C_0}{b_n} \frac{1}{|\operatorname{Im} z|^4} \\ &= \frac{2(1+C_0)}{b_n |\operatorname{Im} z|^4}. \end{aligned}$$

We also have

$$\mathbb{E} \left[\left| \frac{1}{A(z)} - \mathbb{E}_1 \left[\frac{1}{A(z)} \right] \right|^2 \right] \leq \frac{1}{|\operatorname{Im} z|^2} \mathbb{E} \left[\left| \frac{A_1^\circ}{\mathbb{E}_1[A]} \right|^2 \right].$$

Note that $\mathbb{E}_1[A] = z + (1/b_n) \sum_{i \in I_1} G_{ii}^{(1)}$. Since $\operatorname{Im} G_{ii}^{(1)} > 0$, we have $|\mathbb{E}_1[A]| \geq |\operatorname{Im} A| \geq y$. Also we know that $|G_{ii}^{(1)}| \leq 1/|\operatorname{Im} z| = 1/y$. Therefore, $|\mathbb{E}_1[A]| \geq |x| - 2/y$. Combining these we have

$$|\mathbb{E}_1[A]| > \max \left\{ y, |x| - \frac{2}{y} \right\}.$$

Therefore,

$$\begin{aligned} \mathbb{E} \left[|\gamma_n - \mathbb{E}_1[\gamma_n]|^2 \right] &\leq C_1 \frac{2(1+C_0)}{b_n |\operatorname{Im} z|^4} |\mathbb{E}_1[A]|^{-2} + \frac{C_2}{b_n} \left(\sigma^2 + \frac{2+2\mu_4}{|\operatorname{Im} z|^2} \right) \frac{|\mathbb{E}_1[A]|^{-2}}{|\operatorname{Im} z|^2} \\ &\leq \frac{C}{b_n} \left(\frac{1}{|\operatorname{Im} z|^2} + \frac{1}{|\operatorname{Im} z|^4} \right) |\mathbb{E}_1[A]|^{-2}, \end{aligned}$$

for some $C_1, C_2, C > 0$ not depending on z, n . This implies

$$\mathbf{D}(\gamma_n) \leq \frac{Cn}{b_n} \left(\frac{1}{|\operatorname{Im} z|^2} + \frac{1}{|\operatorname{Im} z|^4} \right) \left(\max \left\{ y, |x| - \frac{2}{y} \right\} \right)^{-2}.$$

This completes the proof of Proposition 2.

Now we proceed to the proofs of the asymptotic estimates. All the asymptotic estimates listed in Lemma 1 and Lemma 2 hold uniformly in the set $\{z \in \mathbb{C}: |\operatorname{Im} z| \geq \eta\}$ for any given $\eta > 0$.

Lemma 1. Let M be an $n \times n$ symmetric band matrix as defined in (2) which satisfies (3) and $\mathbb{E}[|w_{ij}|^8]$ is uniformly bounded. Then

(i)

$$G_{ii}^{(1)} - G_{ii} = \frac{1}{A(z)} (G^{(1)} m^{(1)})_i^2 = \frac{1}{A(z)} \left(\frac{1}{\sqrt{b_n}} \sum_{j \in I_1} G_{ij}^{(1)} w_{1j} \right)^2, \quad (32)$$

where $2 \leq i \leq n$, $A(z)$, $m^{(1)}$, and $G^{(1)}$ are as defined in (14), (15), and (16).

(ii) $|\mathbb{E}[G_{ii}^{(1)}(z)] - \mathbb{E}[G_{ii}(z)]| = O(1/b_n)$.

(iii)

$$\begin{aligned} \mathbb{E}[|G_{12}|^2] &= O\left(\frac{1}{b_n}\right) \frac{1}{|\operatorname{Im} z|^6}, \quad \mathbb{E}[|G_{12}|^4] = O\left(\frac{1}{b_n^2}\right) \frac{1}{|\operatorname{Im} z|^{12}}, \\ \mathbb{E}[|G_{12}|^8] &= O\left(\frac{1}{b_n^4}\right) \frac{1}{|\operatorname{Im} z|^{24}}. \end{aligned} \quad (33)$$

(iv) Let us denote the averaging with respect to $\{w_{1i}\}_{1 \leq i \leq n}$ by $\mathbb{E}_1$. Then

$$\begin{aligned} b_n \mathbb{E}_1[A^\circ(z_1)A^\circ(z_2)] &= \sigma^2 + \frac{2}{b_n} \sum_{i,j \in I_1} G_{ij}^{(1)}(z_1)G_{ij}^{(1)}(z_2) \\ &\quad + \frac{\kappa_4}{b_n} \sum_{i \in I_1} G_{ii}^{(1)}(z_1)G_{ii}^{(1)}(z_2) \\ &\quad + \frac{1}{b_n} \widetilde{\gamma_{n-1}}(z_1) \widetilde{\gamma_{n-1}}(z_2), \end{aligned} \quad (34)$$

where $\widetilde{\gamma_{n-1}}(z) = \sum_{i \in I_1} (G_{ii}^{(1)} - \mathbb{E}[G_{ii}^{(1)}(z)])$ and $I_1 = \{1 < i \leq n: (1, i) \in I_n\}$.

(v)

$$\begin{aligned} \mathbb{E}_1[A^\circ(z_1)B^\circ(z_2)] &= \frac{d}{dz_2} \mathbb{E}_1[A^\circ(z_1)A^\circ(z_2)], \quad \text{where} \\ B(z_2) &= \left\langle G^{(1)}(z_2)G^{(1)}(z_2)m^{(1)}, m^{(1)} \right\rangle. \end{aligned} \quad (35)$$

Lemma 2. Let M be an $n \times n$ symmetric band matrix as defined in (2) which satisfies (3). Also assume that the probability distribution of w_{jk} satisfies the Poincaré inequality with some uniform constant m which does not depend on n, j, k . Then

(i)

$$\mathbf{D}\left(\sum_{(1,i) \in I_n} G_{ii}\right) = O(1) \quad \text{and} \quad \mathbf{D}(G_{11}(z)) = O\left(\frac{1}{b_n}\right). \quad (36)$$

(ii)

$$\mathbb{E}[|A^\circ|^4] = O\left(\frac{1}{b_n^2}\right), \quad \mathbb{E}[|A^\circ|^3] = O\left(\frac{1}{b_n^{3/2}}\right), \quad (37)$$

$$\mathbb{E}[|B^\circ|^4] = O\left(\frac{1}{b_n^2}\right). \quad (38)$$

(iii)

$$\begin{aligned} \mathbf{D} \{b_n \mathbb{E}_1 [A^\circ(z_1)A^\circ(z_2)]\} &= O\left(\frac{1}{b_n}\right) \quad \text{and} \\ \mathbf{D} \{b_n \mathbb{E}_1 [A^\circ(z_1)B^\circ(z_2)]\} &= O\left(\frac{1}{b_n}\right). \end{aligned} \quad (39)$$

(iv)

$$\mathbb{E} \left[|\gamma_{n-1}^\circ(z) - \gamma_n^\circ(z)|^4 \right] = O\left(\frac{1}{b_n^2}\right) \quad \text{and} \quad \mathbb{E} \left[|\gamma_n^\circ|^4 \right] = O\left(\frac{n^2}{b_n^2}\right). \quad (40)$$

(v)

$$\frac{1}{n} \mathbb{E} [\text{Tr } G(z)] = f(z) + O\left(\frac{1}{|\text{Im } z|^6 b_n}\right), \quad \text{where } f(z) = \frac{1}{4} \left(-z + \sqrt{z^2 - 8} \right). \quad (41)$$

(vi)

$$(\mathbb{E} [A(z)])^{-1} = -f(z) + O(b_n^{-1}) \quad \text{and} \quad \mathbb{E} [B(z)] = 2f'(z) + O(b_n^{-1}). \quad (42)$$

P r o o f o f L e m m a 1. **P r o o f** of (i). Suppose $(X_1, X_2, \dots, X_n)$ is a n dimensional normal random vector with a positive definite covariance matrix A^{-1} and a mean $A^{-1}\underline{h}$, where $\underline{h} \in \mathbb{R}^n$. Then we have

$$\int \exp \left[-\frac{1}{2} \langle A\underline{x}, \underline{x} \rangle + \langle \underline{h}, \underline{x} \rangle \right] d\underline{x} = (2\pi)^{n/2} |\det A|^{-1/2} \exp \left[\frac{1}{2} \langle A^{-1}\underline{h}, \underline{h} \rangle \right], \quad (43)$$

$$\frac{\int x_i x_j \exp \left[-\frac{1}{2} \langle A\underline{x}, \underline{x} \rangle + \langle \underline{h}, \underline{x} \rangle \right] d\underline{x}}{\int \exp \left[-\frac{1}{2} \langle A\underline{x}, \underline{x} \rangle + \langle \underline{h}, \underline{x} \rangle \right] d\underline{x}} = (A^{-1})_{ij} + (A^{-1}\underline{h})_i (A^{-1}\underline{h})_j, \quad (44)$$

where $\underline{x} = (x_1, x_2, \dots, x_n)^T$. In particular, for $\underline{h} = 0$,

$$(A^{-1})_{ij} = \frac{\int x_i x_j \exp[-\langle A\underline{x}, \underline{x} \rangle / 2] d\underline{x}}{\int \exp[-\langle A\underline{x}, \underline{x} \rangle / 2] d\underline{x}}. \quad (45)$$

Now doing the integrations in (45) with respect to all variables except x_1 , and using (43) we get

$$\begin{aligned} \int \exp \left[-\frac{1}{2} \langle A\underline{x}, \underline{x} \rangle \right] d\underline{x} &= \int \exp \left[-\frac{a_{11}x_1^2}{2} \right] \\ &\quad \times \int \exp \left[-\frac{1}{2} \langle A_1\underline{x}^{(1)}, \underline{x}^{(1)} \rangle - \langle x_1 \underline{a}_1, \underline{x}^{(1)} \rangle \right] d\underline{x} \\ &= \frac{(2\pi)^{(n-1)/2}}{|\det A_1|^{1/2}} \int \exp \left[-\frac{x_1^2}{2} (a_{11} - \langle A_1^{-1} \underline{a}_1, \underline{a}_1 \rangle) \right] dx_1, \end{aligned}$$

where $\underline{x}^{(1)} = (x_2, x_3, \dots, x_n)^T$, $\underline{a}_1 = (a_{12}, a_{13}, \dots, a_{1n})^T$ and $A_1 = ((A_1)_{ij})_{i,j=2}^n$ is the $(n-1) \times (n-1)$ matrix obtained from A after removing first row and first column, and for $i, j \neq 1$, using (44) and (43) we get

$$\begin{aligned} & \int x_i x_j \exp \left[-\frac{1}{2} \langle A \underline{x}, \underline{x} \rangle \right] d\underline{x} \\ &= \int \exp \left[-\frac{a_{11} x_1^2}{2} \right] \int x_i x_j \exp \left[-\frac{1}{2} \langle A_1 \underline{x}^{(1)}, \underline{x}^{(1)} \rangle - \langle x_1 \underline{a}_1, \underline{x}^{(1)} \rangle \right] d\underline{x}^{(1)} dx_1 \\ &= \int \exp \left[-\frac{a_{11} x_1^2}{2} \right] [(A_1^{-1})_{ij} + x_1^2 (A_1^{-1} \underline{a}_1)_i (A_1^{-1} \underline{a}_1)_j] \\ & \quad \times \int \exp \left[-\frac{1}{2} \langle A_1 \underline{x}^{(1)}, \underline{x}^{(1)} \rangle - \langle x_1 \underline{a}_1, \underline{x}^{(1)} \rangle \right] d\underline{x}^{(1)} dx_1 \\ &= \frac{(2\pi)^{(n-1)/2}}{|\det A_1|^{1/2}} \int [(A_1^{-1})_{ij} + x_1^2 (A_1^{-1} \underline{a}_1)_i (A_1^{-1} \underline{a}_1)_j] \\ & \quad \times \exp \left[-\frac{x_1^2}{2} (a_{11} - \langle A_1^{-1} \underline{a}_1, \underline{a}_1 \rangle) \right] dx_1. \end{aligned}$$

Therefore, from (45) we get

$$\begin{aligned} (A^{-1})_{ij} &= (A_1^{-1})_{ij} + (A_1^{-1} \underline{a}_1)_i (A_1^{-1} \underline{a}_1)_j \\ & \quad \times \frac{\int x_1^2 \exp[-x_1^2 (a_{11} - \langle A_1^{-1} \underline{a}_1, \underline{a}_1 \rangle)/2] dx_1}{\int \exp[-x_1^2 (a_{11} - \langle A_1^{-1} \underline{a}_1, \underline{a}_1 \rangle)/2] dx_1} \\ &= (A_1^{-1})_{ij} + \frac{(A_1^{-1} \underline{a}_1)_i (A_1^{-1} \underline{a}_1)_j}{a_{11} - \langle A_1^{-1} \underline{a}_1, \underline{a}_1 \rangle}. \end{aligned}$$

Applying the above formula for $A = (M - zI)$, where $z \in \mathbb{R}$, $|z| > \|M\|$, we obtain

$$G_{ij} = G_{ij}^{(1)} + \frac{(G^{(1)} m^{(1)})_i (G^{(1)} m^{(1)})_j}{w_{11}/\sqrt{b_n} - z - \langle G^{(1)} m^{(1)}, m^{(1)} \rangle}, \quad i, j \geq 2,$$

where $m^{(1)}$, $G^{(1)}$ are as defined in (15), (16), respectively. From the above formula we obtain

$$G_{ii} - G_{ii}^{(1)} = -\frac{(G^{(1)} m^{(1)})_i^2}{A(z)}, \quad \text{for all } 2 \leq i \leq n,$$

where $A(z)$ is as defined in (14). The above is true for all $z \in \mathbb{R}$ such that $|z| > \|M\|$. By analytic continuity one can extend it to the whole complex plane. This completes the proof.

P r o o f o f (i i). Recall $I_1 = \{1 < i \leq n: (1, i) \in I_n\}$. Now using (32) and (21) we have

$$\left| \mathbb{E} [G_{ii}^{(1)}(z)] - \mathbb{E} [G_{ii}(z)] \right| = \left| \mathbb{E} \left[\frac{1}{A} \left(G^{(1)} m^{(1)} \right)_i^2 \right] \right|$$

$$\begin{aligned}
 &= \left| \mathbb{E} \left[\frac{1}{A} \left(\frac{1}{\sqrt{b_n}} \sum_{j \in I_1} G_{ij}^{(1)} w_{1j} \right)^2 \right] \right| \\
 &\leq \frac{1}{b_n} \frac{1}{|\operatorname{Im} z|} \mathbb{E} \left[\left| \sum_{j \in I_1} G_{ij}^{(1)} w_{1j} \right|^2 \right] \\
 &\leq \frac{1}{b_n |\operatorname{Im} z|} \mathbb{E} \left[\sum_{j \in I_1} |G_{ij}^{(1)}|^2 w_{1j}^2 + \sum_{j_1 \neq j_2 \in I_1} G_{ij_1}^{(1)} \overline{G_{ij_2}^{(1)}} w_{1j_1} w_{1j_2} \right] \\
 &= \frac{1}{b_n |\operatorname{Im} z|} \mathbb{E} \mathbb{E}_1 \left[\sum_{j \in I_1} |G_{ij}^{(1)}|^2 w_{1j}^2 + \sum_{j_1 \neq j_2 \in I_1} G_{ij_1}^{(1)} \overline{G_{ij_2}^{(1)}} w_{1j_1} w_{1j_2} \right] \\
 &= \frac{1}{b_n |\operatorname{Im} z|} \mathbb{E} \left[\sum_{j \in I_1} |G_{1j}^{(1)}|^2 \right] \\
 &\leq \frac{1}{b_n |\operatorname{Im} z|} \mathbb{E} \|G^{(1)}\|^2 \leq \frac{1}{b_n |\operatorname{Im} z|^3}.
 \end{aligned}$$

P r o o f o f (i i i). Using the resolvent formula given in [7], we have

$$G_{12} = -G_{22} G_{11}^{(2)} K_{12}^{(12)},$$

where $G^{(2)}$ is the resolvent of the $(n-1) \times (n-1)$ minor obtained by removing the k th row and k th column from the matrix M ,

$$K_{12}^{(12)} = m_{12} - m_{(1)} G^{(12)} m_{(2)},$$

$$m_{(1)} = \frac{1}{\sqrt{b_n}} (w_{13}, w_{14}, \dots, w_{1n}), \quad m_{(2)} = \frac{1}{\sqrt{b_n}} (w_{23}, w_{24}, \dots, w_{2n})^T,$$

$G^{(ij)} = (M^{(ij)} - zI)^{-1}$, and $M^{(ij)}$ is $(n-2) \times (n-2)$ matrix obtained from M after removing i th and j th rows and columns. Therefore,

$$\begin{aligned}
 \mathbb{E} [|G_{12}|^2] &= \mathbb{E} \left[\left| G_{22} G_{11}^{(2)} K_{12}^{(12)} \right|^2 \right] \\
 &\leq \frac{1}{|\operatorname{Im} z|^2} \frac{1}{|\operatorname{Im} z|^2} \mathbb{E} \left[\left| m_{12} - m_{(1)} G^{(12)} m_{(2)} \right|^2 \right] \\
 &= \frac{1}{|\operatorname{Im} z|^4} \mathbb{E} \left[\left| \frac{w_{12}}{\sqrt{b_n}} - \frac{1}{b_n} \sum_{\substack{(1,i), (2,j) \in I_n \\ i,j \neq 1,2}} G_{ij}^{(12)} w_{1i} w_{2j} \right|^2 \right] \\
 &\leq \frac{1}{|\operatorname{Im} z|^4} \mathbb{E} \mathbb{E}_{\leq 2} \left[\frac{w_{12}^2}{b_n} + \frac{1}{b_n^2} \sum_{\substack{(1,i), (2,j) \in I_n \\ i,j \neq 1,2}} |G_{ij}^{(12)}|^2 w_{1i}^2 w_{2j}^2 \right] \\
 &\leq \frac{1}{|\operatorname{Im} z|^4} \mathbb{E} \left[\frac{1}{b_n} + \frac{1}{b_n^2} \sum_{i,j} |G_{ij}^{(12)}|^2 \mathbb{E}_{\leq 2} [w_{1i}^2] \mathbb{E}_{\leq 2} [w_{2j}^2] \right] \\
 &\leq \frac{1}{|\operatorname{Im} z|^4} \mathbb{E} \left[\frac{1}{b_n} + \frac{1}{b_n^2} \frac{b_n}{|\operatorname{Im} z|^2} \right] = O\left(\frac{1}{b_n}\right) \frac{1}{|\operatorname{Im} z|^6},
 \end{aligned}$$

where $E_{\leq 2}$ is the averaging with respect to the first two rows and columns. Similarly, we can prove that

$$\mathbb{E}[|G_{12}|^4] = O\left(\frac{1}{b_n^2}\right) \frac{1}{|\operatorname{Im} z|^{12}}, \quad \text{and} \quad \mathbb{E}[|G_{12}|^8] = O\left(\frac{1}{b_n^4}\right) \frac{1}{|\operatorname{Im} z|^{24}}.$$

P r o o f o f (i v). We know that

$$\begin{aligned} A(z_1) &= z_1 - \frac{w_{11}}{\sqrt{b_n}} + \left\langle G^{(1)} m^{(1)}, m^{(1)} \right\rangle, \\ A^\circ(z_1) &= -\frac{w_{11}}{\sqrt{b_n}} + \frac{1}{b_n} \sum_{i \neq j \in I_1} G_{ij}^{(1)} w_{1i} w_{1j} + \frac{1}{b_n} \sum_{i \in I_1} G_{ii}^{(1)} w_{1i}^2 - \frac{1}{b_n} \sum_{i \in I_1} \mathbb{E}[G_{ii}^{(1)}]. \end{aligned}$$

Now we can estimate

$$\begin{aligned} & b_n \mathbb{E}_1[A^\circ(z_1)A^\circ(z_2)] \\ &= \sigma^2 + \frac{1}{b_n} \mathbb{E}_1 \left[\sum_{\substack{i_1 \neq j_1 \in I_1 \\ i_2 \neq j_2 \in I_1}} G_{i_1 j_1}^{(1)}(z_1) w_{1i_1} w_{1j_1} G_{i_2 j_2}^{(1)}(z_2) w_{1i_2} w_{1j_2} \right] \\ &+ \frac{1}{b_n} \mathbb{E}_1 \left[\sum_{i,j \in I_1} G_{ii}^{(1)}(z_1) G_{jj}^{(1)}(z_2) w_{1i}^2 w_{1j}^2 \right] \\ &- \frac{1}{b_n} \mathbb{E} \left[\sum_{i \in I_1} G_{ii}^{(1)}(z_2) \right] \mathbb{E}_1 \left[\sum_{i \in I_1} G_{ii}^{(1)}(z_1) w_{1i}^2 \right] \\ &- \frac{1}{b_n} \mathbb{E} \left[\sum_{i \in I_1} G_{ii}^{(1)}(z_1) \right] \mathbb{E}_1 \left[\sum_{i \in I_1} G_{ii}^{(1)}(z_2) w_{1i}^2 \right] \\ &+ \frac{1}{b_n} \mathbb{E} \left[\sum_{i \in I_1} G_{ii}^{(1)}(z_1) \right] \mathbb{E} \left[\sum_{i \in I_1} G_{ii}^{(1)}(z_2) \right] \\ &= \sigma^2 + \frac{2}{b_n} \sum_{i \neq j \in I_1} G_{ij}^{(1)}(z_1) G_{ij}^{(1)}(z_2) + \frac{1}{b_n} \sum_{i \neq j \in I_1} G_{ii}^{(1)}(z_1) G_{jj}^{(1)}(z_2) \\ &+ \frac{\mu_4}{b_n} \sum_{i \in I_1} G_{ii}^{(1)}(z_1) G_{ii}^{(1)}(z_2) + \frac{1}{b_n} \widetilde{\gamma_{n-1}}(z_1) \widetilde{\gamma_{n-1}}(z_2) \\ &- \frac{1}{b_n} \left(\sum_{i \in I_1} G_{ii}^{(1)}(z_1) \right) \left(\sum_{i \in I_1} G_{ii}^{(1)}(z_2) \right) \\ &= \sigma^2 + \frac{2}{b_n} \sum_{i,j \in I_1} G_{ij}^{(1)}(z_1) G_{ij}^{(1)}(z_2) + \frac{\mu_4}{b_n} \sum_{i \in I_1} G_{ii}^{(1)}(z_1) G_{ii}^{(1)}(z_2) \\ &- \frac{3}{b_n} \sum_{i \in I_1} G_{ii}^{(1)}(z_1) G_{ii}^{(1)}(z_2) + \frac{1}{b_n} \widetilde{\gamma_{n-1}}(z_1) \widetilde{\gamma_{n-1}}(z_2) \\ &= \sigma^2 + \frac{2}{b_n} \sum_{i,j \in I_1} G_{ij}^{(1)}(z_1) G_{ij}^{(1)}(z_2) + \frac{\kappa_4}{b_n} \sum_{i \in I_1} G_{ii}^{(1)}(z_1) G_{ii}^{(1)}(z_2) \\ &+ \frac{1}{b_n} \widetilde{\gamma_{n-1}}(z_1) \widetilde{\gamma_{n-1}}(z_2), \end{aligned}$$

where $\kappa_4 = \mu_4 - 3$.

P r o o f o f (v). Observe that

$$\begin{aligned} B(z_2) &= \left\langle G^{(1)} G^{(1)} m^{(1)}, m^{(1)} \right\rangle = \frac{1}{b_n} \sum_{i,j \in I_1} \left(G^{(1)} G^{(1)} \right)_{ij} w_{1i} w_{1j} \\ &= \frac{1}{b_n} \sum_{i,j \in I_1} \sum_{k=2}^n G_{ik}^{(1)} G_{kj}^{(1)} w_{1i} w_{1j}, \end{aligned}$$

and

$$\frac{d}{dz_2} G_{ij}^{(1)}(z_2) = \left(G^{(1)}(z_2) G^{(1)}(z_2) \right)_{ij} = \sum_{k=2}^n G_{ik}^{(1)}(z_2) G_{kj}^{(1)}(z_2).$$

Now, proceed as in (iv) and use the above facts to prove the result. Here we skip the details.

P r o o f o f L e m m a 2. P r o o f o f (i). Since w_{jk} satisfies the Poincaré inequality with constant m and the Poincaré inequality tensorises, the joint distribution of $\{w_{jk}\}_{(j,k) \in I_n^+}$ on $\mathbb{R}^{n(b_n+1)}$ satisfies the Poincaré inequality with same constant m . Therefore, we have

$$\mathbf{D} \left(\Phi(\{w_{jk}\}_{(j,k) \in I_n^+}) \right) \leq \frac{1}{m} \sum_{(j,k) \in I_n^+} \mathbb{E} \left[\left| \frac{\partial \Phi}{\partial w_{jk}} \right|^2 \right],$$

for any continuously differentiable function Φ . Therefore,

$$\begin{aligned} \mathbf{D} \left(\sum_{(1,i) \in I_n} G_{ii} \right) &\leq \frac{1}{m} \sum_{(j,k) \in I_n^+} \mathbb{E} \left[\left| \frac{\partial}{\partial w_{jk}} \sum_{(1,i) \in I_n} G_{ii} \right|^2 \right] \\ &\leq \frac{4}{mb_n} \sum_{(j,k) \in I_n^+} \mathbb{E} \left[\left| \sum_{(1,i) \in I_n} G_{ij} G_{ki} \right|^2 \right] \\ &= \frac{4}{mb_n} \sum_{(j,k) \in I_n^+} \mathbb{E} [|\alpha_{kj}|^2] \quad \text{where } \alpha_{kj} = \sum_{(1,i) \in I_n} G_{ki} G_{ij} \\ &\leq \frac{4}{mb_n} \sum_{j,k=1}^n \mathbb{E} [|\alpha_{kj}|^2] \\ &= \frac{4}{mb_n} \mathbb{E} [\|VV^T\|_{Fb}^2] \\ &= \frac{4}{mb_n} \mathbb{E} \left[\sum_{i=1}^n |\beta_i|^2 \right], \end{aligned} \tag{46}$$

where

$$V = \begin{bmatrix} G_{11} & G_{12} & \cdots & G_{1k_n} & 0 & \cdots & 0 \\ G_{21} & G_{22} & \cdots & G_{2k_n} & 0 & \cdots & 0 \\ & & & \vdots & & & \\ G_{n1} & G_{n2} & \cdots & G_{nk_n} & 0 & \cdots & 0 \end{bmatrix}_{n \times n}$$

and $\|\cdot\|_{Fb}$ stands for the Frobenius norm, and β_i s are the eigenvalues of VV^T . Here, we denote the set $\{i: (1, i) \in I_n\}$ by $\{1, 2, \dots, k_n\}$. Observe that $k_n = 2b_n + 1$. Since $\text{rank}(VV^T) \leq k_n = O(b_n)$, we have $\#\{i: \beta_i \neq 0\} \leq k_n = O(b_n)$. Also we know that $\|V\| \leq \|G\|$. Therefore,

$$|\beta_i|^2 \leq \|VV^T\|^2 \leq \|G\|^4 \leq \frac{1}{|\text{Im } z|^4}.$$

Consequently, we have

$$\mathbf{D}\left(\sum_{(1,i) \in I_n} G_{ii}\right) \leq \frac{4}{mb_n} \mathbb{E}\left[\sum_{i=1}^n |\beta_i|^2\right] \leq \frac{4}{mb_n} \frac{O(b_n)}{|\text{Im } z|^4} = O(1). \quad (47)$$

This completes proof of first part of (36).

Recall the definition of A from (14), $A = z - b_n^{-1/2}w_{11} + (G^{(1)}m^{(1)}, m^{(1)})$. Then

$$\begin{aligned} A^\circ &= A - \mathbb{E}[A] \\ &= -\frac{1}{\sqrt{b_n}}w_{11} + \frac{1}{b_n} \sum_{\substack{i \neq j \\ i, j \in I_1}} G_{ij}^{(1)} w_{1i} w_{1j} + \frac{1}{b_n} \sum_{i \in I_1} \left(G_{ii}^{(1)} w_{1i}^2 - \mathbb{E}[G_{ii}^{(1)}]\right), \end{aligned}$$

Consider

$$\begin{aligned} A_1^\circ &= A - \mathbb{E}_1[A] \\ &= -\frac{1}{\sqrt{b_n}}w_{11} + \frac{1}{b_n} \sum_{\substack{i \neq j \\ i, j \in I_1}} G_{ij}^{(1)} w_{1i} w_{1j} + \frac{1}{b_n} \sum_{i \in I_1} \left(G_{ii}^{(1)} w_{1i}^2 - G_{ii}^{(1)}\right). \end{aligned} \quad (48)$$

So we have

$$A^\circ - A_1^\circ = \frac{1}{b_n} \sum_{i \in I_1} \left(G_{ii}^{(1)} - \mathbb{E}[G_{ii}^{(1)}]\right) =: \frac{1}{b_n} \widetilde{\gamma_{n-1}}. \quad (49)$$

Hence

$$\mathbb{E}[|A^\circ|^2] = \mathbb{E}[|A_1^\circ + b_n^{-1} \widetilde{\gamma_{n-1}}|^2] \leq 2 \left[\mathbb{E}[|A_1^\circ|^2] + \frac{1}{b_n^2} \mathbb{E}[|\widetilde{\gamma_{n-1}}|^2] \right].$$

From (31), we know that $\mathbb{E}[|A_1^\circ|^2] = O(1/b_n)$ and from (47), We have $\mathbb{E}[|\widetilde{\gamma_{n-1}}|^2] = O(1)$. Combining these two facts and using (17), we have

$$\mathbf{D}(G_{11}(z)) = \mathbb{E} \left| \frac{1}{A} - \mathbb{E} \frac{1}{A} \right|^2 \leq \mathbb{E} \left| \frac{1}{A} - \frac{1}{\mathbb{E}A} \right|^2 = \mathbb{E} \left| \frac{A^\circ}{A \mathbb{E}A} \right|^2 = O\left(\frac{1}{b_n}\right).$$

This completes the proof of second part.

P r o o f o f (i i). Proof of (37). Recall from (48)

$$A_1^\circ = -\frac{w_{11}}{\sqrt{b_n}} + \frac{1}{b_n} \sum_{i \neq j \in I_1} G_{ij}^{(1)} w_{1i} w_{1j} + \frac{1}{b_n} \sum_{i \in I_1} G_{ii}^{(1)} (w_{1i}^2)^\circ =: T_1 + T_2 + T_3. \quad (50)$$

We have $\mathbb{E}[|T_1|^4] = O(1/b_n^2)$. Now

$$\mathbb{E}[|T_2|^4] = \frac{1}{b_n^4} \mathbb{E} \left[\sum_{i \neq j, k \neq l, p \neq q, s \neq t \in I_1} G_{ij}^{(1)} \overline{G_{kl}^{(1)}} G_{pq}^{(1)} \overline{G_{st}^{(1)}} w_{1i} w_{1j} w_{1k} w_{1l} w_{1p} w_{1q} w_{1s} w_{1t} \right].$$

We use the similar technique as the moment method in the proof of the semicircle law. In the above sum of expectations, we have nonzero terms if the indices of w_{1m} 's match in a certain way. Nonzero contribution to $\mathbb{E}[|T_2|^4]$ come from the two types of matches.

Fig. 1. Type I matching.

Fig. 2. Type II matching.

Type I. Contribution from this kind of matching is

$$\begin{aligned} & \frac{1}{b_n^4} \mathbb{E} \left[\sum_{i \neq j, p \neq k \in I_1} |G_{ij}^{(1)}|^2 |G_{pq}^{(1)}|^2 w_{1i}^2 w_{1j}^2 w_{1p}^2 w_{1q}^2 \right] \\ &= \frac{1}{b_n^4} \mathbb{E} \mathbb{E}_1 \left[\sum_{i \neq j, p \neq k \in I_1} |G_{ij}^{(1)}|^2 |G_{pq}^{(1)}|^2 w_{1i}^2 w_{1j}^2 w_{1p}^2 w_{1q}^2 \right] \\ &= \frac{1}{b_n^4} \sum_{i \neq j} \sum_{p \neq q} \mathbb{E} \left[|G_{ij}^{(1)}|^2 |G_{pq}^{(1)}|^2 \right] \\ &\leq \frac{1}{b_n^4} \sum_{i \neq j} \sum_{p \neq q} \sqrt{\mathbb{E} \left[|G_{ij}^{(1)}|^4 \right] \mathbb{E} \left[|G_{pq}^{(1)}|^4 \right]} \\ &= \frac{1}{b_n^4} \sum_{i \neq j} \sum_{p \neq q} O\left(\frac{1}{b_n^2}\right) \quad (\text{using (33)}) \\ &= O\left(\frac{1}{b_n^2}\right). \end{aligned}$$

Type II. Similarly, contribution from the type II matching is

$$\frac{1}{b_n^4} \mathbb{E} \left[\sum_{i \neq j} \sum_{q \neq l \in I_1} G_{ij}^{(1)} \overline{G_{il}^{(1)}} G_{jq}^{(1)} \overline{G_{ql}^{(1)}} w_{1i}^2 w_{1j}^2 w_{1q}^2 w_{1l}^2 \right] = O\left(\frac{1}{b_n^2}\right).$$

Similarly, $\mathbb{E}[|T_3|^4] = O(1/b_n^2)$. Hence

$$\mathbb{E}[|A_1^\circ|^4] = O\left(\frac{1}{b_n^2}\right). \quad (51)$$

Using Lemma 4.4.3. from [1] with the help of the Poincaré inequality, we have $\mathbb{E}[|\widetilde{\gamma_{n-1}}|^4] \leq C \|\nabla \widetilde{\gamma_{n-1}}\|_2^4$, where C is a constant depends only on the constant m of the Poincaré inequality. Following the arguments given at the right side of (46) onward and (47), one can show that $\|\nabla \widetilde{\gamma_{n-1}}\|_2 \leq C/|\operatorname{Im} z|^4$, where C depends only on m . Hence $\mathbb{E}[|\widetilde{\gamma_{n-1}}|^4] = O(1)$. Consequently, using relation (49) and (51), we have $\mathbb{E}[|A^\circ|^4] = O(1/b_n^2)$. Then $\mathbb{E}[|A^\circ|^3] \leq (\mathbb{E}[|A^\circ|^4])^{3/4} = O(1/b_n^{3/2})$.

P r o o f of (38). First we write B as

$$B = \left\langle G^{(1)} G^{(1)} m^{(1)}, m^{(1)} \right\rangle = \left\langle H^{(1)} m^{(1)}, m^{(1)} \right\rangle = \frac{1}{b_n} \sum_{i,j \in I_1} H_{ij}^{(1)} w_{1i} w_{1j},$$

where $H^{(1)} = G^{(1)} G^{(1)}$. Define

$$B_1^\circ := \frac{1}{b_n} \sum_{i \neq j \in I_1} H_{ij}^{(1)} w_{1i} w_{1j} + \frac{1}{b_n} \sum_{i \in I_1} H_{ii}^{(1)} (w_{1i}^2)^\circ.$$

Then we can write

$$\begin{aligned} B^\circ &= B - \mathbb{E}[B] \\ &= \frac{1}{b_n} \sum_{i \neq j \in I_1} H_{ij}^{(1)} w_{1i} w_{1j} + \frac{1}{b_n} \sum_{i \in I_1} \left[H_{ii}^{(1)} w_{1i}^2 - \mathbb{E}[H_{ii}^{(1)}] \right] \\ &= B_1^\circ + \frac{1}{b_n} \sum_{i \in I_1} \left(H_{ii}^{(1)} - \mathbb{E}[H_{ii}^{(1)}] \right) \\ &= B_1^\circ + \frac{1}{b_n} \overline{\gamma_{n-1}}, \end{aligned}$$

where

$$\begin{aligned} \overline{\gamma_{n-1}}(z) &= \sum_{i \in I_1} \left(H_{ii}^{(1)} - \mathbb{E}[H_{ii}^{(1)}] \right) = \sum_{i \in I_1} \sum_{j=2}^n \left(G_{ij}^{(1)} G_{ji}^{(1)} - \mathbb{E}[G_{ij}^{(1)} G_{ji}^{(1)}] \right) \\ &= \frac{d}{dz} \widetilde{\gamma_{n-1}}(z). \end{aligned}$$

Proceeding as in the estimate of $\mathbb{E}[|A_1^\circ|^4]$, we can show

$$\mathbb{E}[|B_1^\circ|^4] = O\left(\frac{1}{b_n^2}\right). \quad (52)$$

We have shown that $\mathbb{E}[|\widetilde{\gamma_{n-1}}(z)|^4] = O(1)$. Using this fact and Cauchy's theorem we have $\mathbb{E}[|\overline{\gamma_{n-1}}(z)|^4] = O(1)$. Hence we have the result.

P r o o f of (i i i).

$$\begin{aligned} \mathbf{D}\left\{b_n \mathbb{E}_1[A^\circ(z_1) A^\circ(z_2)]\right\} &= \mathbf{D}(T_1) + \mathbf{D}(T_2) + \mathbf{D}(T_3) \\ &\quad + 2 \operatorname{Cov}(T_1, T_2) + 2 \operatorname{Cov}(T_2, T_3) + 2 \operatorname{Cov}(T_3, T_1), \end{aligned}$$

where

$$T_1 = \frac{2}{b_n} \sum_{i,j \in I_1} G_{ij}^{(1)}(z_1) G_{ij}^{(1)}(z_2), \quad T_2 = \frac{\kappa_4}{b_n} \sum_{i \in I_1} G_{ii}^{(1)}(z_1) G_{ii}^{(1)}(z_2),$$

$$T_3 = \frac{1}{b_n} \widetilde{\gamma_{n-1}}(z_1) \widetilde{\gamma_{n-1}}(z_2).$$

Now,

$$\mathbf{D}(T_2) = \frac{\kappa_4^2}{b_n^2} \mathbf{D} \left\{ \sum_{i \in I_1} G_{ii}^{(1)}(z_1) G_{ii}^{(1)}(z_2) \right\}$$

and

$$\begin{aligned} \mathbf{D} \left\{ G_{ii}^{(1)}(z_1) G_{ii}^{(1)}(z_2) \right\} &= \mathbb{E} \left| G_{ii}^{(1)}(z_1) G_{ii}^{(1)}(z_2) - \mathbb{E} [G_{ii}^{(1)}(z_1) G_{ii}^{(1)}(z_2)] \right|^2 \\ &\leq \frac{2}{|\operatorname{Im} z_1|^2} \mathbf{D} \left(G_{ii}^{(1)}(z_2) \right) + \frac{2}{|\operatorname{Im} z_2|^2} \mathbf{D} \left(G_{ii}^{(1)}(z_1) \right) \\ &= \left(\frac{1}{|\operatorname{Im} z_1|^2} + \frac{1}{|\operatorname{Im} z_2|^2} \right) O \left(\frac{1}{b_n} \right). \end{aligned}$$

Therefore,

$$\mathbf{D}(T_2) \leq \frac{\kappa_4^2}{b_n^2} \left(b_n O \left(\frac{1}{b_n} \right) + b_n^2 O \left(\frac{1}{b_n} \right) \right) = O \left(\frac{1}{b_n} \right).$$

Now

$$\begin{aligned} \mathbf{D}(T_3) &\leq \frac{1}{b_n^2} \mathbf{D} \left(\widetilde{\gamma_{n-1}}(z_1) \widetilde{\gamma_{n-1}}(z_2) \right) \\ &\leq \frac{1}{b_n^2} \mathbb{E} [|\widetilde{\gamma_{n-1}}(z_1)|^2 |\widetilde{\gamma_{n-1}}(z_2)|^2] \\ &\leq \frac{1}{b_n^2} \sqrt{\mathbb{E} [|\widetilde{\gamma_{n-1}}(z_1)|^4]} \sqrt{\mathbb{E} [|\widetilde{\gamma_{n-1}}(z_2)|^4]} = \frac{1}{b_n^2} O(1). \end{aligned}$$

Last equality holds, since $\mathbb{E} [|\widetilde{\gamma_{n-1}}(z_1)|^4] = O(1)$. And finally

$$\mathbf{D}(T_1) = \frac{4}{b_n^2} \mathbf{D} \left(\sum_{i,j \in I_1} G_{ij}^{(1)}(z_1) G_{ij}^{(1)}(z_2) \right).$$

Now using the Poincaré inequality

$$\begin{aligned} &\mathbf{D} \left(\sum_{i,j \in I_1} G_{ij}^{(1)}(z_1) G_{ij}^{(1)}(z_2) \right) \\ &\leq \frac{1}{m} \sum_{(s,t) \in I_n^+} \mathbb{E} \left[\left| \frac{\partial}{\partial w_{st}} \sum_{i,j \in I_1} G_{ij}^{(1)}(z_1) G_{ij}^{(1)}(z_2) \right|^2 \right] \\ &\leq \frac{1}{mb_n} \sum_{(s,t) \in I_n^+} \mathbb{E} \left[\left| \sum_{i,j \in I_1} G_{is}^{(1)}(z_1) G_{tj}^{(1)}(z_1) G_{ij}^{(1)}(z_2) \right|^2 \right] \end{aligned}$$

$$\begin{aligned}
& + G_{ij}^{(1)}(z_1) G_{is}^{(1)}(z_2) G_{tj}^{(1)}(z_2) \Big|^2 \Big] \\
& \leq \frac{2}{mb_n} \sum_{(s,t) \in I_n^+} \mathbb{E} \left[\left| \sum_{i,j \in I_1} G_{is}^{(1)}(z_1) G_{tj}^{(1)}(z_1) G_{ij}^{(1)}(z_2) \right|^2 \right] \\
& \quad + \frac{2}{mb_n} \sum_{(s,t) \in I_n^+} \mathbb{E} \left[\left| \sum_{i,j \in I_1} G_{ij}^{(1)}(z_1) G_{is}^{(1)}(z_2) G_{tj}^{(1)}(z_2) \right|^2 \right] \\
& =: I_1 + I_2.
\end{aligned}$$

We estimate

$$\begin{aligned}
I_1 &= \frac{2}{mb_n} \sum_{(s,t) \in I_n^+} \mathbb{E} \left[\left| \sum_{i,j \in I_1} G_{is}^{(1)}(z_1) G_{tj}^{(1)}(z_1) G_{ij}^{(1)}(z_2) \right|^2 \right] \\
&= \frac{2}{mb_n} \sum_{(s,t) \in I_n^+} \mathbb{E} \left[\left| \sum_{i \in I_1} G_{is}^{(1)}(z_1) G_{ti}^{(1)}(z_1, z_2) \right|^2 \right] \\
&= \frac{2}{mb_n} \sum_{(s,t) \in I_n^+} \mathbb{E} \left[\left| G_{st}^{(1)}(z_1, z_2, z_1) \right|^2 \right] \\
&\leq \frac{2}{mb_n} \mathbb{E} \left[\sum_{s,t=1}^n \left| G_{st}^{(1)}(z_1, z_2, z_1) \right|^2 \right] = \frac{2}{mb_n} \mathbb{E} [\|A\|_{Fb}^2] \\
&= \frac{2}{mb_n} \mathbb{E} \left[\sum_{i=1}^n \beta_i^2 \right] \leq \frac{C(z_1, z_2)}{mb_n} O(b_n) = O(1),
\end{aligned}$$

where $\|\cdot\|_{Fb}$ is the Frobenius norm, β_i are the eigenvalues of VV^* , and V is the following matrix:

$$\begin{aligned}
V_{n \times n} &= \begin{bmatrix} G_{11}^{(1)}(z_1) & G_{12}^{(1)}(z_1) & \cdots & G_{1k_n}^{(1)}(z_1) \\ G_{21}^{(1)}(z_1) & G_{22}^{(1)}(z_1) & \cdots & G_{2k_n}^{(1)}(z_1) \\ \vdots & \vdots & & \vdots \\ G_{n1}^{(1)}(z_1) & G_{n2}^{(1)}(z_1) & \cdots & G_{nk_n}^{(1)}(z_1) \end{bmatrix}_{n \times k_n} \\
&\times \begin{bmatrix} G_{11}^{(1)}(z_2) & G_{12}^{(1)}(z_2) & \cdots & G_{1k_n}^{(1)}(z_2) \\ G_{21}^{(1)}(z_2) & G_{22}^{(1)}(z_2) & \cdots & G_{2k_n}^{(1)}(z_2) \\ \vdots & \vdots & & \vdots \\ G_{k_n1}^{(1)}(z_2) & G_{k_n2}^{(1)}(z_2) & \cdots & G_{k_nk_n}^{(1)}(z_2) \end{bmatrix}_{k_n \times k_n} \\
&\times \begin{bmatrix} G_{11}^{(1)}(z_1) & G_{12}^{(1)}(z_1) & \cdots & G_{1n}^{(1)}(z_1) \\ G_{21}^{(1)}(z_1) & G_{22}^{(1)}(z_1) & \cdots & G_{2n}^{(1)}(z_1) \\ \vdots & \vdots & & \vdots \\ G_{k_n1}^{(1)}(z_1) & G_{k_n2}^{(1)}(z_1) & \cdots & G_{k_nn}^{(1)}(z_1) \end{bmatrix}_{k_n \times n}.
\end{aligned}$$

Here we denoted the elements of set I_1 as $I_1 = \{1, 2, \dots, k_n\}$. Observe that $k_n = 2b_n$. Rank of $V \leq k_n = O(b_n)$. This implies

$$\sum_{i=1}^n \beta_i^2 \leq k_n C(z_1, z_2) = O(b_n) C(z_1, z_2).$$

Therefore, $\mathbf{D}(T_1) = O(1/b_n^2)$, and hence $\mathbf{D}\{b_n \mathbb{E}_1[A^\circ(z_1)A^\circ(z_2)]\} = O(1/b_n)$.

Second part of (iii) follows from the following two facts with the help of Cauchy's theorem:

$$\begin{aligned} b_n \mathbb{E}_1[A^\circ(z_1)B^\circ(z_2)] &= b_n \frac{d}{dz_2} \mathbb{E}_1[A^\circ(z_1)A^\circ(z_2)], \\ \mathbf{D}\{b_n \mathbb{E}_1\{A^\circ(z_1)A^\circ(z_2)\}\} &= O\left(\frac{1}{b_n}\right). \end{aligned}$$

Here we skip the details.

P r o o f o f (i v). Using (25) and (30), and proceeding as in the proof of Proposition 2,

$$\begin{aligned} \mathbb{E}\left[|\gamma_{n-1}^\circ(z) - \gamma_n^\circ(z)|^4\right] &= \mathbb{E}\left[\left|\left(\mathrm{Tr} G^{(1)}(z) - \mathbb{E}[\mathrm{Tr} G^{(1)}(z)]\right) \right. \right. \\ &\quad \left. \left. - \left(\mathrm{Tr} G(z) - \mathbb{E}[\mathrm{Tr} G(z)]\right)\right|^4\right] \\ &= \mathbb{E}\left[\left|\frac{1+B(z)}{A(z)} - \mathbb{E}\left[\frac{1+B(z)}{A(z)}\right]\right|^4\right] \\ &\leq \frac{C}{|\mathrm{Im} z|^8} \left(\mathbb{E}[|A^\circ|^4] + \mathbb{E}[|B^\circ|^4] + \mathbb{E}[|A^\circ|^4]\right) \\ &= O\left(\frac{1}{b_n^2}\right). \end{aligned}$$

The last equality follows from the estimates (37) and (38).

Using martingale differences as in the proof of Proposition 2,

$$\mathbb{E}[|\gamma_n^\circ|^4] \leq Cn \sum_{k=1}^n \mathbb{E}[|\gamma_n - \mathbb{E}_k[\gamma_n]|^4].$$

Consider for $k = 1$, others will be similar.

$$\begin{aligned} \mathbb{E}\left[|\gamma_n - \mathbb{E}_1[\gamma_n]|^4\right] &= \mathbb{E}\left[\left|\mathrm{Tr}(G - G^{(1)}) - \mathbb{E}_1[\mathrm{Tr}(G - G^{(1)})]\right|^4\right] \\ &= \mathbb{E}\left[\left|\frac{1+B(z)}{A(z)} - \mathbb{E}_1\left[\frac{1+B(z)}{A(z)}\right]\right|^4\right] \\ &\leq C_1(z) \mathbb{E}[|A_1^\circ|^4] + C_2(z) \mathbb{E}[|B_1^\circ|^4] = O\left(\frac{1}{b_n^2}\right). \end{aligned}$$

The last equality follows from (51) and (52). Hence we have the result.

P r o o f o f (v). Using resolvent identity

$$(X_2 - zI)^{-1} = (X_1 - zI)^{-1} + (X_1 - zI)^{-1}(X_1 - X_2)(X_2 - zI)^{-1},$$

we have

$$zG_{11}(z) = -1 + \sum_{(1,k) \in I_n} m_{1k} G_{k1}, \quad (53)$$

where I_n is defined in (1) and m_{ij} 's are defined in (2). Now to analyse the terms $\mathbb{E}[m_{1k} G_{k1}]$, we use the following (see, e.g., [10]): Given ξ , a real-valued random variable with $p+2$ finite moments, and ϕ , a function from $\mathbb{C}$ to $\mathbb{R}$ with $p+1$ continuous and bounded derivatives then:

$$\mathbb{E}[\xi \phi(\xi)] = \sum_{a=0}^p \frac{\kappa_{a+1}}{a!} \mathbb{E}[\phi^{(a)}(\xi)] + \epsilon_{p+1}, \quad (54)$$

where κ_a is the a -th cumulant of ξ , $|\epsilon_{p+1}| \leq C \sup_t |\phi^{(p+1)}(t)| \mathbb{E}[|\xi|^{p+2}]$ and C depends only on p . Since $f_n(z) = (1/n) \mathbb{E}[\text{Tr } G(z)] = \mathbb{E}[G_{11}(z)]$, using (53) and (54) we get

$$\begin{aligned} z f_n(z) &= -1 + \sum_{(1,k) \in I_n} \mathbb{E}[m_{1k} G_{k1}] \\ &= -1 - \sum_{k \in I_1} \frac{1}{b_n} \mathbb{E}[G_{k1}^2 + G_{kk} G_{11}] + r_n, \end{aligned} \quad (55)$$

where r_n contains the third cumulant term corresponding to $p=2$ in (54) for $k \neq 1$, and the error terms due to the truncation of the decoupling formula (54) at $p=2$ for $k \neq 1$ and at $p=0$ for $k=1$. We write (55)

$$\begin{aligned} z f_n(z) &= -1 - \frac{1}{b_n} \mathbb{E}[G_{11}] \mathbb{E}\left[\sum_{k \in I_1} G_{kk}\right] - \frac{1}{b_n} \text{Cov}\left(G_{11}, \sum_{k \in I_1} G_{kk}\right) \\ &\quad - \frac{1}{b_n} \mathbb{E}\left[\sum_{k \in I_1} G_{k1}^2\right] + r_n \\ &= -1 - f_n(z)(2f_n(z)) - \frac{1}{b_n} \text{Cov}\left(G_{11}, \sum_{k \in I_1} G_{kk}\right) \\ &\quad - \frac{1}{b_n} \mathbb{E}\left[\sum_{k \in I_1} G_{k1}^2\right] + r_n. \end{aligned}$$

Now, by the Cauchy–Schwarz inequality and (47) we get

$$\begin{aligned} \frac{1}{b_n} \left| \text{Cov}\left(G_{11}, \sum_{k \in I_1} G_{kk}\right) \right| &\leq \frac{1}{b_n} \sqrt{\mathbf{D}(G_{11})} \sqrt{\mathbf{D}\left(\sum_{k \in I_1} G_{kk}\right)} \\ &\leq \frac{1}{b_n} 2 |\text{Im } z|^{-1} \sqrt{O(|\text{Im } z|^{-4})} \\ &= O\left(\frac{1}{b_n |\text{Im } z|^3}\right). \end{aligned}$$

Also notice that

$$\frac{1}{b_n} \left| \mathbb{E} \left(\sum_{k \in I_1} G_{k1}^2 \right) \right| \leq \frac{1}{b_n} |\operatorname{Im} z|^{-2}.$$

We claim $r_n = O(1/(b_n |\operatorname{Im} z|^4))$. To prove this, observe that the third cumulant term gives

$$\frac{\kappa_3}{2b_n^{3/2}} \mathbb{E} \left[\sum_{k \in I_1} 2(G_{1k})^3 + 6G_{11}G_{1k}G_{kk} \right]. \quad (56)$$

Since $\sum_{k \in I_1} |G_{1k}|^2 \leq \|G\|^2 \leq |\operatorname{Im} z|^{-2}$ and $|G_{ij}| \leq |\operatorname{Im} z|^{-1}$, we conclude that the third cumulant term contributes $O(1/(b_n |\operatorname{Im} z|^3))$ to r_n . In a similar manner, the error due to truncation of decoupling formula (54) at $p = 2$ is $O(1/(b_n |\operatorname{Im} z|^4))$. Similarly, the error term due to truncation of decoupling formula at $p = 0$ for $k = 1$ is $O(1/(b_n |\operatorname{Im} z|^2))$. Thus the claim is proved. Hence

$$zf_n(z) = -1 - 2f_n^2(z) + O\left(\frac{1}{b_n |\operatorname{Im} z|^4}\right) \quad \text{for } z \in \mathbb{C} \setminus \mathbb{R}.$$

Now following similar argument given in the proof of (3.1) in [14], one can show that

$$|f_n(z) - f(z)| \leq O\left(\frac{1}{b_n |\operatorname{Im} z|^6}\right),$$

where $f(z) = (-z + \sqrt{z^2 - 8})/4$.

P r o o f o f (v i). Recall $A(z) = z - b_n^{-1/2}w_{11} + b_n^{-1} \sum_{i,j \in I_1} G_{ij}^{(1)} w_{1i} w_{1j}$. Now using (41) with G replaced by $G^{(1)}$, we have

$$\begin{aligned} (\mathbb{E}[A(z)])^{-1} &= \frac{1}{z + b_n^{-1} \sum_{j \in I_1} \mathbb{E}[G_{jj}^{(1)}]} = \frac{1}{z + 2f_n(z)} \\ &= (z + 2f(z))^{-1} + O(b_n^{-1}) = -f(z) + O(b_n^{-1}). \end{aligned}$$

Hence $(\mathbb{E}[A(z)])^{-1} = -f(z) + O(b_n^{-1})$. To prove the second part, observe that

$$\begin{aligned} \mathbb{E}[B(z)] &= \frac{1}{b_n} \mathbb{E} \left[\sum_{i,j \in I_1} (G^{(1)} G^{(1)})_{ij} w_{1i} w_{1j} \right] = \frac{1}{b_n} \mathbb{E} \left[\sum_{i \in I_1} (G^{(1)} G^{(1)})_{ii} \right] \\ &= \frac{1}{b_n} \mathbb{E} \left[\sum_{i \in I_1} \sum_{k=2}^n G_{ik}^{(1)} G_{ki}^{(1)} \right] = \frac{1}{b_n} \sum_{i \in I_1} \frac{d}{dz} G_{ii}^{(1)}. \end{aligned}$$

Again using (41) and Cauchy's integral formula, we have

$$\mathbb{E}[B(z)] = \frac{d}{dz} (2f_n(z)) = 2f'(z) + O(b_n^{-1}).$$

This completes the proof of Lemma 2.

4.1. Proof of (29).

P r o o f. We have to find the limit of

$$\mathbb{E}[T_n] = \frac{2}{b_n} \mathbb{E} \left[\sum_{i,j \in I_1} G_{ij}^{(1)}(z) G_{ij}^{(1)}(z_\mu) \right]$$

as $n \rightarrow \infty$, where $I_1 = \{2 \leq i \leq n: (1, i) \in I_n\}$. Let $f, g \in C_b(\mathbb{R})$. Define a bilinear form on $C_b(\mathbb{R})$ as

$$\langle f, g \rangle_n = \frac{1}{b_n} \mathbb{E} \left[\sum_{i,j \in I_1} f(M)_{ij} g(M)_{ji} \right]. \quad (57)$$

Then $\mathbb{E}[T_n] = \langle h(M), h_\mu(M) \rangle_n$, where $h(x) = (x - z)^{-1}$ and $h_\mu(x) = (x - z_\mu)^{-1}$.

Lemma 3. For $f, g \in C_b(\mathbb{R})$ the limit $\langle f, g \rangle = \lim_{n \rightarrow \infty} \langle f, g \rangle_n$ exists.

P r o o f. The idea of the proof is similar to the proof of Lemma 3.11 of [9]. First we prove this result for monomials. Although monomials are unbounded, still (57) makes sense for all n , since all moments of the entries of M are finite. Consider $f(x) = x^l$, and $g(x) = x^m$ where $l, m \in \mathbb{N}$. Then

$$\langle x^l, x^m \rangle_n = \frac{1}{b_n^{1+(l+m)/2}} \sum_{\substack{(i_0, i_1), (i_1, i_2), \dots, (i_{l+m-1}, i_0) \in I_n \\ i_0, i_l \in I_1}} \mathbb{E} [w_{i_0 i_1} w_{i_1 i_2} \cdots w_{i_{l+m-1} i_0}].$$

If $(l + m)$ is odd, then $\langle x^l, x^m \rangle_n \rightarrow 0$ using independence of matrix entries and $\mathbb{E}[w_{ij}] = 0$, and order counting of independent vertices. The argument is similar to the combinatorial argument given in the proof of Wigner semi-circular law (see [1]). We leave it for the reader.

Now we assume $l + m$ is even. Then

$$\begin{aligned} \langle x^l, x^m \rangle_n &= \frac{1}{b_n^{1+(l+m)/2}} \sum_{\substack{(i_0, i_1), (i_1, i_2), \dots, (i_{l+m-1}, i_0) \in I_n \\ i_0, i_l \in I_1}} \mathbb{E} [w_{i_0 i_1} w_{i_1 i_2} \cdots w_{i_{l+m-1} i_0}] \\ &= \frac{1}{b_n^{1+(l+m)/2}} \\ &\quad \times \sum_{\substack{(i_0, i_1), (i_1, i_2), \dots, (i_{l+m-1}, i_0) \in I_n \\ i_0, i_l \in I_1}} \mathbb{E} [w_{1 i_0} w_{i_0 i_1} w_{i_1 i_2} \cdots w_{i_{l+m-1} i_0} w_{i_0 1}] + O(b_n^{-1}) \\ &= \frac{1}{b_n^{1+(l+m)/2}} \\ &\quad \times \sum_{\substack{(i_0, i_1), (i_1, i_2), \dots, (i_{l+m-1}, i_0) \in I_n \\ (1, i_0), (1, i_l) \in I_n}} \mathbb{E} [w_{1 i_0} w_{i_0 i_1} w_{i_1 i_2} \cdots w_{i_{l+m-1} i_0} w_{i_0 1}] + O(b_n^{-1}). \end{aligned} \quad (58)$$

The second last equality in (58) holds due to order calculation of independent vertices and independence of matrix entries. Now define for $k = 1, 2, \dots, l + m$,

$$\begin{aligned} x_k &= \begin{cases} i_k - i_{k-1} & \text{if } |i_k - i_{k-1}| \leq b_n, \\ (i_k - i_{k-1}) - n & \text{if } i_k - i_{k-1} > b_n, \\ n + (i_k - i_{k-1}) & \text{if } i_k - i_{k-1} < -b_n \end{cases} \quad \text{with } i_{l+m} = i_0, \\ x_0 &= \begin{cases} i_0 - 1 & \text{if } |i_0 - 1| \leq b_n, \\ (i_0 - 1) - n & \text{if } i_0 - 1 > b_n, \end{cases} \\ x_{l+m+1} &= \begin{cases} 1 - i_0 & \text{if } |1 - i_0| \leq b_n, \\ n + (1 - i_0) & \text{if } 1 - i_0 < -b_n. \end{cases} \end{aligned}$$

Note, $x_0 = -x_{l+m+1}$. Since l, m are fixed and $b_n \rightarrow \infty$, for large n the restrictions $\{(i_0, i_1), (i_1, i_2), \dots, (i_{l+m-1}, i_0) \in I_n \text{ and } (1, i_0), (1, i_l) \in I_n\}$ are equivalent to $\{|x_0|, |x_1|, \dots, |x_{l+m}| \leq b_n, x_0 + x_1 + \dots + x_{l+m} + x_{l+m+1} = 0 \text{ and } |x_0 + x_1 + \dots + x_l| \leq b_n\}$. Also observe that $x_0 + x_1 + \dots + x_{l+m} + x_{l+m+1} = 0$ is same as $x_1 + \dots + x_{l+m} = 0$ since $x_0 = -x_{l+m+1}$. Therefore for large n

$$\begin{aligned} \langle x^l, x^m \rangle_n &= \frac{1}{b_n^{1+(l+m)/2}} \\ &\sum_{\substack{x_1 + \dots + x_{l+m} = 0 \\ |x_i| \leq b_n, 0 \leq i \leq l+m, |x_0 + x_1 + \dots + x_l| \leq b_n}} \mathbb{E} [w_{1i_0} w_{i_0 i_1} w_{i_1 i_2} \dots w_{i_{l+m-1} i_0} w_{i_0 1}] + O(b_n^{-1}). \end{aligned}$$

Without loss of generality, we assume that $l \leq m$. Each $\{i_0, i_1, i_2, \dots, i_{l+m-1}, i_0\}$ is a closed path such that distance between the end points of each edge is bounded by b_n . As in the proof of Wigner semicircular law only the paths whose edges are pair matched contributes to the limit, here also, only such paths contribute to the limit. And contribution of each path is $\mathbb{E}\{w_{1i_0} w_{i_0 i_1} \dots w_{i_{l+m-1} i_0} w_{i_0 1}\} = 1$ since $\mathbb{E}\{w_{ij}^2\} = 1$. Each such path corresponds to a Dyck path of length $(l + m)$. Recall that a Dyck path $(S(0), S(1), \dots, S(l + m))$ of length $(l + m)$ satisfies (see [1])

$$\begin{aligned} S(0) = S(l + m) &= 0, \quad S(1), S(2), \dots, S(l + m - 1) \geq 0, \\ |S(i + 1) - S(i)| &= 1, \quad \text{for } i = 0, 1, \dots, l + m - 1. \end{aligned}$$

Specifically, $S(t + 1) - S(t) = 1$ if the non oriented edge (i_t, i_{t+1}) appears in $\{i_0, i_1, \dots, i_{l+m-1}, i_0\}$ for the first time and $S(t + 1) - S(t) = -1$ if the edge (i_t, i_{t+1}) appears in $\{i_0, i_1, \dots, i_{l+m-1}, i_0\}$ for the second time.

Here each Dyck path does not give equal contribution to the limit due to the condition that $(1, i_l) \in I_n$ and in terms of x_i , which is same as $|x_0 + x_1 + \dots + x_l| \leq b_n$. We have to take into account this condition. Suppose $S(l) = k$, $0 \leq k \leq l$. Then during the first l steps of the path $\{i_0, i_1, \dots, i_{l+m-1}, i_0\}$, k edges appear only once and $(l - k)/2$ edges appear twice. The edges appearing

twice, the corresponding two number x_i have same absolute value but with different sign. We rename the remaining k numbers x_i which appear only once as $y_1, y_2, \dots, y_k$ (according to their order of appearance) and x_0 as y_0 . So the condition $|x_0 + x_1 + \dots + x_l| \leq b_n$ reduces to $|y_0 + y_1 + \dots + y_k| \leq b_n$. Therefore,

$$\begin{aligned} \langle x^l, x^m \rangle_n &= \frac{1}{b_n^{1+(l+m)/2}} \sum_{k=0}^l \#\{\text{Dyck path of length } l+m \text{ with } S(l)=k\} \\ &\quad \times \#\left\{ |y_0| \leq b_n, |y_1| \leq b_n, \dots, |y_k| \leq b_n, \dots, |y_{l+m}| \leq b_n, \right. \\ &\quad \left. |y_0 + y_1 + \dots + y_k| \leq b_n \right\} + O(b_n^{-1}) \end{aligned}$$

and

$$\begin{aligned} \langle x^l, x^m \rangle &= \lim_{n \rightarrow \infty} \langle x^l, x^m \rangle_n \\ &= (\sqrt{2})^{l+m+2} \sum_{k=0}^l \#\{\text{Dyck path of length } l+m \text{ with } S(l)=k\} \\ &\quad \times \text{Vol} \left\{ |t_0| \leq \frac{1}{2}, |t_1| \leq \frac{1}{2}, \dots, |t_{(l+m)/2}| \leq \frac{1}{2}, |t_0 + t_1 + \dots + t_k| \leq \frac{1}{2} \right\} \\ &= (\sqrt{2})^{l+m+2} \sum_{k=0}^l \#\{\text{Dyck path of length } l+m \text{ with } S(l)=k\} \\ &\quad \times P \left(|T_0 + T_1 + \dots + T_k| \leq \frac{1}{2} \right), \end{aligned}$$

where $T_0, T_1, \dots, T_{(l+m)/2}$ are independent random variables uniformly distributed on $[-1/2, 1/2]$. Let $S_{k+1} = T_0 + T_1 + \dots + T_k$. Then

$$\mathbb{E}[e^{ixS_{k+1}}] = (\mathbb{E}[e^{ixT_0}])^{k+1} = \left(\frac{\sin x/2}{x/2} \right)^{k+1}.$$

Using inversion formula, the density of S_{k+1} is given by

$$f_{k+1}(s) = \frac{1}{2\pi} \int_{-\infty}^{\infty} e^{-ixs} \left(\frac{\sin x/2}{x/2} \right)^{k+1} dx.$$

Now

$$\begin{aligned} \gamma_{k+1} &:= P \left(|S_{k+1}| \leq \frac{1}{2} \right) = \int_{-1/2}^{1/2} f_{k+1}(s) ds \\ &= \frac{1}{2\pi} \int_{-\infty}^{\infty} \left(\frac{\sin x/2}{x/2} \right)^{k+2} dx = f_{k+2}(0), \end{aligned}$$

using [5] we get exact formula of γ_{k+1} :

$$\gamma_{k+1} = \begin{cases} \frac{1}{(k+1)!} \sum_{s=0}^{(k+1)/2} (-1)^s \binom{k+2}{s} \left(\frac{k+1}{2} - s + \frac{1}{2} \right)^{k+1} & \text{if } k+1 \text{ even,} \\ \frac{1}{(k+1)!} \sum_{s=0}^{k/2} (-1)^s \binom{k+2}{s} \left(\frac{k+1}{2} - s + \frac{1}{2} \right)^{k+1} & \text{if } k+1 \text{ odd.} \end{cases} \quad (59)$$

The number of Dyck path of length $l+m$ with $S(l) = k$ is

$$\begin{aligned} & \left[\binom{l}{\frac{l-k}{2}} - \binom{l}{\frac{l-k-2}{2}} \right] \times \left[\binom{m}{\frac{m-k}{2}} - \binom{m}{\frac{m-k-2}{2}} \right] \\ &= \frac{(k+1)^2}{(l+1)(m+1)} \binom{l+1}{\frac{l+k+2}{2}} \binom{m+1}{\frac{m+k+2}{2}}. \end{aligned} \quad (60)$$

Hence from (59) and (60), we get

$$\langle x^l, x^m \rangle = (\sqrt{2})^{l+m+2} C_{l,m},$$

where $C_{l,m} = 0$ if $(l+m)$ is odd and

$$\begin{aligned} C_{l,m} &= \sum_{k=0}^l \frac{(k+1)^2}{(l+1)(m+1)} \binom{l+1}{\frac{l+k+2}{2}} \binom{m+1}{\frac{m+k+2}{2}} \gamma_{k+1} \\ &= \begin{cases} \sum_{k=0}^{l/2} \frac{(2k+1)^2}{(l+1)(m+1)} \binom{l+1}{\frac{l-2k}{2}} \binom{m+1}{\frac{m-2k}{2}} \gamma_{2k+1} & \text{if } l \text{ even,} \\ \sum_{k=0}^{(l-1)/2} \frac{(2k+2)^2}{(l+1)(m+1)} \binom{l+1}{\frac{l-2k-1}{2}} \binom{m+1}{\frac{m-2k-1}{2}} \gamma_{2k+2} & \text{if } l \text{ odd,} \end{cases} \end{aligned}$$

if $(l+m)$ is even and $l \leq m$, otherwise, $C_{l,m} = C_{m,l}$. If f, g are polynomials, $f(x) = \sum_{i=0}^p a_i x^i$, $g(x) = \sum_{i=0}^q b_i x^i$, then by linearity

$$\langle f, g \rangle = \sum_{i=0}^p \sum_{j=0}^q a_i b_j (\sqrt{2})^{i+j+2} C_{i,j}. \quad (61)$$

For general bounded continuous functions f, g , to show that $\langle f, g \rangle$ exists we have to use the Stone–Weierstrass theorem to approximate f, g by appropriate polynomial and then (61). The argument is similar to the argument given in the proof of Lemma 3.11 of [9]. We skip the details.

In the next lemma we diagonalize the bilinear form $\langle f, g \rangle$.

Lemma 4. *Let $\{U_n(x)\}_{n \geq 0}$ be the rescaled Chebyshev polynomial of the second kind on $[-2\sqrt{2}, 2\sqrt{2}]$,*

$$U_n(x) = \sum_{k=0}^{\lfloor n/2 \rfloor} (-1)^k \binom{n-k}{k} \left(\frac{x}{\sqrt{2}} \right)^{n-2k}.$$

Then $\{U_n(x)\}$ are orthogonal with respect to the bilinear form (61), that is,

$$\langle U_n, U_m \rangle = 2\delta_{nm}\gamma_{n+1}, \quad (62)$$

where γ_{n+1} is defined in (59).

P r o o f. The proof of this lemma is similar to the proof of Lemma 3.12 of [9]. For sake of completeness we outline it here. Since $\langle x^l, x^m \rangle = 0$ if $l + m$ is odd, from linearity $\langle U_l, U_m \rangle = 0$ if $l + m$ is odd. We are left to compute $\langle U_{2n}, U_{2m} \rangle$ and $\langle U_{2n+1}, U_{2m+1} \rangle$. We first compute $\langle x^{2l}, U_{2n} \rangle$ and $\langle x^{2l+1}, U_{2n+1} \rangle$ for $l = 0, 1, \dots, n$:

$$\begin{aligned} \langle x^{2l}, U_{2n} \rangle &= (\sqrt{2})^{2l+2} \sum_{k=0}^n (-1)^k \binom{2n-k}{k} C_{2l, 2n-2k} \\ &= (\sqrt{2})^{2l+2} \left[\sum_{k=0}^{n-l} (-1)^k \binom{2n-k}{k} \sum_{t=0}^l \frac{(2t+1)^2}{(2l+1)(2n-2k+1)} \right. \\ &\quad \times \binom{2l+1}{l-t} \binom{2n-2k+1}{n-k-t} \gamma_{2t+1} \\ &\quad + \sum_{k=n-l+1}^n (-1)^k \binom{2n-k}{k} \sum_{t=0}^{n-k} \frac{(2t+1)^2}{(2l+1)(2n-2k+1)} \\ &\quad \times \left. \binom{2l+1}{l-t} \binom{2n-2k+1}{n-k-t} \gamma_{2t+1} \right] \\ &= (\sqrt{2})^{2l+2} \sum_{t=0}^l \frac{(2t+1)^2}{2l+1} \binom{2l+1}{l-t} \\ &\quad \times \left[\sum_{k=0}^{n-t} \frac{(-1)^k (2n-k)!}{k! (n-k-t)! (n-k+t+1)!} \right] \gamma_{2t+1} \\ &= (\sqrt{2})^{2l+2} \sum_{t=0}^l \frac{(2t+1)^2}{2l+1} \binom{2l+1}{l-t} G_1(n, t) \gamma_{2t+1}, \end{aligned}$$

where

$$G_1(n, t) = \sum_{k=0}^{n-t} \frac{(-1)^k (2n-k)!}{k! (n-k-t)! (n-k+t+1)!}.$$

Similarly,

$$\begin{aligned} \langle x^{2l+1}, U_{2n+1} \rangle &= (\sqrt{2})^{2l+3} \sum_{t=0}^l \frac{(2t+2)^2}{2l+2} \binom{2l+2}{l-t} \\ &\quad \times \left[\sum_{k=0}^{n-t} \frac{(-1)^k (2n+1-k)!}{k! (n-k-t)! (n-k+t+2)!} \right] \gamma_{2t+2} \\ &= (\sqrt{2})^{2l+3} \sum_{t=0}^l \frac{(2t+2)^2}{2l+2} \binom{2l+2}{l-t} G_2(n, t) \gamma_{2t+2}, \end{aligned}$$

where

$$G_2(n, t) = \sum_{k=0}^{n-t} \frac{(-1)^k (2n+1-k)!}{k! (n-k-t)! (n-k+t+2)!}.$$

$G_1(n, t)$ and $G_2(n, t)$ can be written in terms of hypergeometric function as follows:

$$G_1(n, t) = \frac{(2n)!}{(n-t)! (n+t+1)!} {}_2F_1 \left(\begin{matrix} -(n-t), -(n+t+1) \\ -2n \end{matrix}; 1 \right),$$

$$G_2(n, t) = \frac{(2n+1)!}{(n-t)! (n+t+2)!} {}_2F_1 \left(\begin{matrix} -(n-t), -(n+t+2) \\ -2n-1 \end{matrix}; 1 \right),$$

where ${}_2F_1$ is a hypergeometric function. By the Chu–Vandermonde identity (see [2]), we have

$${}_2F_1 \left(\begin{matrix} -(n-t), -(n+t+1) \\ -2n \end{matrix}; 1 \right) = \frac{(-n+t+1)_{n-t}}{(-2n)_{n-t}},$$

$${}_2F_1 \left(\begin{matrix} -(n-t), -(n+t+2) \\ -2n-1 \end{matrix}; 1 \right) = \frac{(-n+t+1)_{n-t}}{(-2n-1)_{n-t}},$$

where $(a)_n = a(a+1) \cdots (a+n-1)$. Since

$$(-n+t+1)_{n-t} = \begin{cases} 0 & \text{if } t = 0, 1, \dots, n-1, \\ 1 & \text{if } t = n, \end{cases}$$

we have $G_1(n, t) = 0$, $G_2(n, t) = 0$ for $t = 0, 1, \dots, n-1$ and $G_1(n, n) = 1/(2n+1)$, $G_2(n, n) = 1/(2n+2)$. Therefore, $\langle x^{2l}, U_{2n} \rangle = 0$ for $0 \leq l \leq n-1$ and $\langle x^{2n}, U_{2n} \rangle = (\sqrt{2})^{2n+2} \gamma_{2n+1}$. Similarly, $\langle x^{2l+1}, U_{2n+1} \rangle = 0$ for $0 \leq l \leq n-1$ and $\langle x^{2n+1}, U_{2n+1} \rangle = (\sqrt{2})^{2n+3} \gamma_{2n+2}$. Therefore,

$$\langle U_{2n}, U_{2n} \rangle = 2\gamma_{2n+1} \text{ and } \langle U_{2n+1}, U_{2n+1} \rangle = 2\gamma_{2n+2}.$$

This completes the proof of the lemma.

Now we complete the proof of (29). For $f, g \in C_b(\mathbb{R})$, if

$$f_k = \frac{1}{4\pi} \int_{-2\sqrt{2}}^{2\sqrt{2}} f(x) U_k(x) \sqrt{8-x^2} dx,$$

$$g_k = \frac{1}{4\pi} \int_{-2\sqrt{2}}^{2\sqrt{2}} g(x) U_k(x) \sqrt{8-x^2} dx,$$

then

$$\langle f, g \rangle = \sum_{k=0}^{\infty} f_k g_k 2\gamma_{k+1} \quad (63)$$

$$\begin{aligned}
&= \frac{1}{8\pi^3} \int_{-2\sqrt{2}}^{2\sqrt{2}} \int_{-2\sqrt{2}}^{2\sqrt{2}} f(x)g(y) \sqrt{8-x^2} \sqrt{8-y^2} \\
&\quad \times \left[\pi \sum_{k=0}^{\infty} U_k(x)U_k(y)\gamma_{k+1} \right] dx dy \\
&= \frac{1}{8\pi^3} \int_{-2\sqrt{2}}^{2\sqrt{2}} \int_{-2\sqrt{2}}^{2\sqrt{2}} f(x)g(y) \sqrt{8-x^2} \sqrt{8-y^2} F(x,y) dx dy,
\end{aligned}$$

where

$$\begin{aligned}
F(x,y) &= \pi \sum_{k=0}^{\infty} U_k(x)U_k(y)\gamma_{k+1} \\
&= 2 \int_{-\infty}^{\infty} \frac{z - z^3}{2(1 - z^2)^2 + z^2(x^2 + y^2) - z(1 + z^2)xy} ds \quad (64)
\end{aligned}$$

with $z = (\sin s)/s$. Now (63) holds due to (62) and orthogonality of Chebyshev polynomial with respect to the Wigner semicircular law, that is,

$$\int_{-2\sqrt{2}}^{2\sqrt{2}} U_n(x)U_m(x) \frac{1}{4\pi} \sqrt{8-x^2} dx = \delta_{mn}.$$

And (64) is a straightforward consequence of the Fourier analysis using the following fact

$$U_n(x) = \frac{\sin[(n+1)\theta]}{\sin \theta}, \quad x = 2\sqrt{2} \cos \theta.$$

This completes the proof of (29).

Recent development. Recently, after submission of our paper, M. Shcherbina [16] improved our result by removing the restriction $b_n \gg \sqrt{n}$ and proved it for all b_n which satisfies $b_n \rightarrow \infty$ and $b_n/n \rightarrow 0$ as $n \rightarrow \infty$.

Acknowledgment. We thank the referee for her/his constructive comments which have led to a significant improvement in presentation.

5. Some MATLAB simulation results. Here is what we found in MATLAB simulations.

a) $n = 2000$, $b_n = n^{0.2}$. Fourth moment/(variance) $^2 = 2.92$.

b) $n = 2000$, $b_n = n^{0.4}$. Fourth moment/(variance) $^2 = 2.71$.

Fig. 3. The eigenvalue statistics was sampled 400 times. The test function was

$$\phi(x) = \sqrt{16 - x^2}.$$

In the following example we had taken a different test function.

Fig. 4. The eigenvalue statistics was sampled 400 times. The test function was

$$\phi(x) = e^{-x^2}.$$

REFERENCES

1. Anderson G. W., Guionnet A., Zeitouni O. An introduction to random matrices. Cambridge: Cambridge Univ. Press, 2010, 492 p.
2. Andrews G. E., Askey R., Roy R. Special functions. Encyclopedia of Mathematics and its Applications. Cambridge: Cambridge Univ. Press, 1999, 664 p.

3. *Casati G., Izrailev F., Molinari L.* Scaling properties of the eigenvalue spacing distribution for band random matrices. — J. Phys. A: Mathematical and General, 1991, v. 24, № 20, p. 4755.
4. *Casati G., Molinari L., Izrailev F.* Scaling properties of band random matrices. — Phys. Rev. Lett., 1990, v. 64, № 16, p. 1851–1854.
5. *Крамер Г.* Математические методы статистики. М.: Мир, 1975, 648 с.
6. *Dharmadhikari S., Fabian V., Jogdeo K.* Bounds on the moments of martingales. — Ann. Math. Statist., 1968, v. 39, № 6, p. 1719–1723.
7. *Эрдёш Л.* Универсальность случайных матриц Вигнера: обзор последних результатов. — Успехи матем. наук., т. 66, № 3, с. 67–198.
8. *Fyodorov Y. V., Mirlin A. D.* Scaling properties of localization in random band matrices: a σ -model approach. — Phys. Rev. Lett., 1991, v. 67, № 18, p. 2405.
9. *Li L., Soshnikov A.* Central limit theorem for linear statistics of eigenvalues of band random matrices. — Random Matrices Theory Appl., 2013, v. 2, № 4, p. 50.
10. *Lytova A., Pastur L.* Central limit theorem for linear eigenvalue statistics of random matrices with independent entries. — Ann. Probab., 2009, v. 37, № 5, p. 1778–1840.
11. *Mirlin A. D., Fyodorov Y. V., Dittes F.-M., Quezada J., Seligman T. H.* Transition from localized to extended eigenstates in the ensemble of power-law random banded matrices. — Phys. Rev. E, 1996, v. 54, № 4, p. 3221.
12. *Molchanov S. A., Pastur L. A., Khorunzhi A.* Limiting eigenvalue distribution for band random matrices. — Theoret. Math. Phys., 1992, v. 90, № 2, p. 108–118.
13. *O'Rourke S., Renfrew D., Soshnikov A.* On fluctuations of matrix entries of regular functions of wigner matrices with non-identically distributed entries. — J. Theoret. Probab., 2013, v. 26, № 3, p. 750–780.
14. *Pizzo A., Renfrew D., Soshnikov A.* Fluctuations of matrix entries of regular functions of Wigner matrices. — J. Statist. Phys., 2012, v. 146, № 3, p. 550–591.
15. *Shcherbina M.* Central limit theorem for linear eigenvalue statistics of the wigner and sample covariance random matrices. — J. Math. Phys. Anal. Geom., 2011, v. 7, № 2, p. 176–192.
16. *Shcherbina M.* On fluctuations of eigenvalues of random band matrices. — J. Stat. Phys., 2015, v. 4, № 1, p. 73–90.
17. *Tulino A. M., Verdú S.* Random Matrix Theory and Wireless Communications, Vol. 1. Princeton: Now Publishers Inc., 2004, 182 p.
18. *Verdú S., Shamai S.* Spectral efficiency of CDMA with random spreading. — IEEE Trans. Inform. Theory, 1999, v. 45, № 2, p. 622–640.
19. *Wigner E. P.* Characteristic vectors of bordered matrices with infinite dimensions. — Ann. Math., 1955, v. 62, № 3, p. 548–564.
20. *Wigner E. P.* Characteristics vectors of bordered matrices with infinite dimensions. II. — Ann. Math., 1957, v. 65, № 2, p. 203–207.
21. *Wishart J.* The generalised product moment distribution in samples from a normal multivariate population. — Biometrika, 1928, v. 20A, № 1–2, p. 32–52.

Поступила в редакцию
4.II.2015