

The quadruple bubble in the plane





E. Paolini

(Università di Pisa)

~~Moscow~~, 17 settembre 2020

Zoom

Joint works with: Tamagnini (2018) and Tortorelli (2020)



Existence of optimal clusters:

We say that \mathbf{E} is a minimal cluster of prescribed areas $\mathbf{a} \in \mathbb{R}_+^N$ if $P(\mathbf{E})$ is minimum among all clusters with $\mathbf{m}(\mathbf{E}) = \mathbf{a}$.



Such minimal cluster exist for any $\mathbf{a} \in \mathbb{R}_+^N$ (Almgren, Morgan)

$n > 3$


$n = 2$

1: i
2: c
 ∞ :
3: t



↑





Pressure formula

$$P(\underline{E}) \quad m(\underline{E}) = (|E_1|, |E_2|, \dots, |E_n|)$$

Pressures $\underline{p} = (p_1, \dots, p_n)$ are Lagrange multipliers associated to the area constraint:

$$\left[\frac{d}{dt} P(\underline{E}(t)) \right]_{t=t_0} = \underline{p} \cdot \left[\frac{d}{dt} \underline{m}(\underline{E}(t)) \right]_{t=t_0}$$

—○—


$$\underline{p} = (p_1, \dots, p_n)$$

Applying the previous formula to a rescaling of a stationary cluster \underline{E} we obtain

$$P(\underline{E}) = 2 \underline{p} \cdot \underline{m}(\underline{E})$$

$$\underline{m}(\underline{E}) = (1, 1, 1, 1)$$

The double bubble



Main difficulty: prove that minimizers are connected.

Tool: rotate a subcluster along an edge (two connected components can share at most one edge)

Main idea: enlarge class of minimizers including clusters which enclose more area than prescribed. This enables to easily prove that E_0 is connected (no internal camera)

Weak minimizers

$$\underline{m}(\epsilon) \leq \underline{a}$$


Weak minimizers

We say that \mathbf{E} is a weak minimizer with prescribed areas $\mathbf{a} \in \mathbb{R}_+^N$ if $P(\mathbf{E})$ is minimum among all sets with $m(\mathbf{E}) \geq \mathbf{a}$.
Notation $\mathbf{E} \in \mathcal{M}^*(\mathbf{a})$.



Properties of weak minimizers:

- the external region E_0 is connected; ←
- no negative pressures: $\mathbf{p} \geq \mathbf{0}$; ←
- if $|E_i| > a_i$ then $p_i = 0$; ←
- if the total number of connected components is $M \leq 6$ then the weak minimizer is a (strong) minimizer.



Möbius transforms / monotonicity

Möbius transformations preserve stationarity of clusters!



Passing through Möbius transformations it is easy to inflate/deflate triangular components of a cluster.







Monotonicity: when a triangular component is inflated (resp. deflated) both the area and the radius of each edge increases (resp. decreases).




Uniqueness: in double-bubbles and triple-bubbles there is a one-to-one correspondence $\mathbf{p} \mapsto \mathbf{a}$ between pressures and areas. (Montesinos)





Four equal areas



with Andrea Tamagnini (unifi): Minimal clusters of four planar regions with the same area (ESAIM COCV 2018)



with Vincenzo M. Tortorelli (unipi): The quadruple planar bubble enclosing equal areas is symmetric (Calc. Var. 2020)

Variational tools

Variation I. Remove a component C with an edge of length ℓ and replace it by a disk of equal area. Then $\ell \leq 2\sqrt{\pi}\sqrt{|C|}$.



Variation II. Remove a component C of the region E_i with n edges and rescale the resulting cluster to recover the prescribed areas. Then

$$|C| \geq \frac{16\pi|E_i|^2}{n^2 P^2(\mathbf{E})} \left(1 - \frac{16\pi|E_i|}{n^2 P^2(\mathbf{E})} \right).$$



Variation III. Remove a component C with n edges and recover the measure by enlarging an external edge ℓ of the same region E_i . Then

$$p_i \geq \frac{2\sqrt{\pi}}{n\sqrt{|C|}} - \frac{2}{\ell}.$$

A priori estimate on the number of components

Let M be the number of connected components of a weak minimal N -cluster $\mathbf{E} \in \mathcal{M}^*(\mathbf{a})$ with $N \geq 3$. Then

$$M \leq \frac{9}{20} N^2 \frac{\|\mathbf{a}\|_{\frac{1}{2}}}{\|\mathbf{a}\|_{-1}}.$$

$$\|\mathbf{a}\|_p = \left(\sum_{j=1}^N |a_j|^p \right)^{1/p}$$

Isoperimetric inequality

$$P(\mathbf{E}) = \frac{1}{2} \sum_{i=0}^N P(E_i) \geq \sqrt{\pi} \left[\sqrt{\sum_{i=1}^N |E_i|} + \sum_{i=1}^N \sqrt{|E_i|} \right]$$

—○—

If $E_i = E'_i \cup E''_i$ is disconnected:

$$P(E_i) \geq 2\sqrt{\pi} \left(\sqrt{|E'_i|} + \sqrt{|E''_i|} \right)$$

—○—

big / small components...










$$\mathbf{E} \in \mathcal{M}^*(1, 1, 1, 1)$$

Explicit computation on well chosen competitor:



$$P(\mathbf{E}) \leq 11.1962$$




- Minimal clusters have at most $4 + 2$ components (big + small).
- Exclude there can be two small components.
- Reduce to 9 possible topologies for clusters with $4 + 1$ components.
- Exclude them: minimal clusters have 4 connected components.
- Two topologies: sandwich, flower.
- Exclude flower.



Flower / Sandwich



the Sandwich is symmetric



the Sandwich is symmetric

Let $F_k = T(E_k)$ with $T(z) = \frac{1}{z}$, $\det DT(z) = \frac{1}{z^4}$

$$\begin{aligned} |E_1| - |E_2| &= \iint_{F_1} \frac{1}{|x + iy|^4} dx dy - \iint_{F_2} \frac{1}{|x + iy|^4} dx dy \\ &= \int_0^{\frac{3}{2}\pi} \int_{r_1(t, \theta)}^{r_2(t, \theta)} \left[\frac{1}{|1 + re^{i(\theta+t)}|^4} - \frac{1}{|1 + re^{i(\theta-t)}|^4} \right] r dr dt. \end{aligned}$$

but

$$|1 + re^{i\alpha}|^2 = 1 + 2r \cos \alpha + r^2$$

and for $t \in (0, \frac{2}{3}\pi]$ and $\theta \in (0, \frac{\pi}{3})$

$$\cos(\theta + t) < \cos(\theta - t).$$