

Chernoff approximations as a method for finding the resolvent of a linear operator and solving a linear ODE with variable coefficients

Ivan D. Remizov^{1,2}

¹ Laboratory of Topological Methods in Dynamics, NRU HSE

25/12 Bol. Pecherskaya Ulitsa, Room 224, Nizhny Novgorod, 603155, Russia

² Institute for Information Transmission Problems of the Russian Academy of Sciences

Bolshoy Karetny per. 19, build.1, Moscow 127051 Russia

ivremizov@yandex.ru

Abstract. The Chernoff approximation method is a powerful and flexible tool of functional analysis, which allows in many cases to express $\exp(tL)$ in terms of variable coefficients of a linear differential operator L . In this paper, we prove a theorem that allows us to apply this method to find the resolvent of L . Our theorem states that the Laplace transforms of Chernoff approximations of a C_0 -semigroup converge to the resolvent of the generator of this semigroup. We demonstrate the proposed method on a second-order differential operator with variable coefficients. As a consequence, we obtain a new representation of the solution of a nonhomogeneous linear ordinary differential equation of the second order in terms of functions that are coefficients of this equation, playing the role of parameters of the problem. For the Chernoff function, based on the shift operator, we give an estimate for the rate of convergence of approximations to the solution.

Keywords: operator semigroups, resolvent operator, linear ODE with variable coefficients, representation of the solution, Chernoff approximations

MSC2020: 47A10, 47D06, 34A05

Черновские аппроксимации как метод нахождения резольвенты линейного оператора и решения линейного ОДУ с переменными коэффициентами

И. Д. Ремизов^{1,2}

¹Лаборатория топологических методов в динамике НИУ ВШЭ

603155, Россия, Нижний Новгород, Большая Печерская, 25/12

²Институт проблем передачи информации имени А. А. Харкевича РАН

127051, г. Москва, Большой Каретный переулок, д.19 стр. 1.

ivremizov@yandex.ru

Аннотация. Метод черновских аппроксимаций является мощным и гибким инструментом функционального анализа, позволяющим во многих случаях выразить $\exp(tL)$ через переменные коэффициенты линейного дифференциального оператора L . В данной работе доказывается теорема, позволяющая применять этот метод для нахождения резольвенты оператора L . Наша теорема утверждает, что преобразования Лапласа аппроксимаций Чернова C_0 -полугруппы сходятся к резольвенте генератора этой полугруппы. Мы демонстрируем предложенный метод на дифференциальном операторе второго порядка с переменными коэффициентами. В качестве следствия мы получаем новое представление решения неоднородного линейного обыкновенного дифференциального уравнения второго порядка в терминах функций, являющихся коэффициентами этого уравнения, играющих роль параметров задачи. Для функции Чернова на основе оператора сдвига мы даем оценку скорости сходимости приближений к решению.

Ключевые слова: полугруппы операторов, резольвента оператора, линейное ОДУ с переменными коэффициентами, представление решения, черновские аппроксимации

УДК 517.986.7+517.988.8+517.926.4

1 Introduction

The straightforward way to find the resolvent $(\lambda I - L)^{-1}$ of the linear operator $(L, D(L))$ in the Banach space \mathcal{F} is to study (for given $\lambda \in \mathbb{C}$) if the equation $\lambda f - Lf = g$ has for each $g \in \mathcal{F}$ a unique solution $f \in D(L)$. If we know that $(L, D(L))$ is closed then this is enough, but if not then we must also check that f depends on g continuously. In case of success we have $f = (\lambda I - L)^{-1}g$. For example, if \mathcal{F} is one of the subsets of space of all functions $f: \mathbb{R} \rightarrow \mathbb{C}$, and $Lf = af'' + bf' + cf$ is a differential operator defined by constant coefficients $a, b, c \in \mathbb{C}$, then this task is not difficult. Indeed $\lambda f - Lf = g$ is a second order differential equation $\lambda f(x) - af''(x) - bf'(x) - cf(x) = g(x)$, $x \in \mathbb{R}$, with only one variable coefficient g , and there are standard formulas in ODE books to solve this equation using the variation of parameters method. But if a, b, c are not constants, but functions that depend on x then the situation becomes much worse because there are no standard formulas for solution of equation $\lambda f(x) - a(x)f''(x) - b(x)f'(x) - c(x)f(x) = g(x)$, $x \in \mathbb{R}$.

However, if $(L, D(L))$ holds one additional property, then we can use a non-straightforward method of finding the resolvent, this method is the main result of the paper. The property is that $(L, D(L))$ is the generator of a C_0 -semigroup, which informally means that exponent e^{tL} exists as a linear bounded operator, and depends on $t \in [0, +\infty)$ continuously in some sense. If $(L, D(L))$ is a linear bounded operator with $D(L) = \mathcal{F}$ then this condition holds automatically, but if $(L, D(L))$ is not bounded then there is a beautiful theory around it, see e.g. [1, 2, 3, 4, 5]. In most interesting cases (e.g. if L is a differential operator with variable coefficients) it is difficult to calculate e^{tL} directly because power series $e^{tL} = \sum_{n=0}^{\infty} (tL)^n/n!$ is useless in case of unbounded operator L . Yet it is possible to find e^{tL} approximately using the Chernoff theorem [6], this theorem will be discussed below. After that, having e^{tL} we calculate the resolvent via well known [3] formula $(\lambda I - L)^{-1}f = \int_0^{\infty} e^{-\lambda t} e^{tL} f dt$. The method may look complicated because of the many steps it has (find Chernoff function for L , find e^{tL} via Chernoff approximations, find $(\lambda I - L)^{-1}$ via e^{tL}), however all these steps are supported by known methods, and so our proposed method may still be much simpler than considering the equation $\lambda f - Lf = g$.

Brief history and overview of the results obtained up to 2017 in constructing Chernoff approximations of e^{tL} for several classes of operators L can be found in [7]. Several papers on the topic showing the diversity of cases studied are [8, 9, 10, 11, 12, 13], see also [14, 15, 16]. Speed of convergence of Chernoff approximations were studied in [17, 18, 19, 20, 21, 22, 23], see also references therein. See also paper [24] which numerically investigates the efficiency of the Monte Carlo method based on the application of the Chernoff theorem, and paper [25] that mathematically substantiate such an approach to the Chernoff approximations.

Summing up, we can say that the method of Chernoff approximation is an extremely effective tool for expressing e^{tL} in terms of variable coefficients of operator L . The present paper shows that this method can be also be used for expressing $(\lambda I - L)^{-1}$ in terms of variable coefficients of operator L , and for finding the solution of the corresponding differential equation $\lambda f - Lf = g$. As an example we consider second order linear ODE with variable coefficients.

Limits of multiple integral as multiplicity tends to infinity (such expressions are called Feynman formulas [8]) are one of the ways (which actually was used originally by Richard Feynman [26, 27]) to define Feynman path integral [28, 29]. So in theorem 5 a solution of an ODE is in the first time in history of science represented via Feynman formula, which can also be interpreted as Feynman integral if one wishes to do. Such theoretical step is novel and may lead to some unexpected applications in the future.

2 Preliminaries: operator semigroups and their Chernoff approximations

Let us recall some relevant definitions and facts of C_0 -semigroup theory following [3].

Definition 1. Let \mathcal{F} be a Banach space over the field \mathbb{R} or \mathbb{C} . Let $\mathcal{L}(\mathcal{F})$ be the set of all bounded linear operators in \mathcal{F} . Suppose we have a mapping $V: [0, +\infty) \rightarrow \mathcal{L}(\mathcal{F})$, i.e. $V(t)$ is a bounded linear operator $V(t): \mathcal{F} \rightarrow \mathcal{F}$ for each $t \geq 0$. The mapping V , or equivalently the family $(V(t))_{t \geq 0}$, is called a *strongly continuous one-parameter semigroup of linear bounded operators* (or just a *C_0 -semigroup*) iff it satisfies the following three conditions:

- 1) $V(0)$ is the identity operator I , i.e. $V(0)\varphi = \varphi$ for each $\varphi \in \mathcal{F}$;
- 2) V maps the addition of numbers in $[0, +\infty)$ into the composition of operators in $\mathcal{L}(\mathcal{F})$, i.e. for all $t \geq 0$ and all $s \geq 0$ we have $V(t+s) = V(t) \circ V(s)$, where for each $\varphi \in \mathcal{F}$ the notation $(A \circ B)(\varphi) = A(B(\varphi)) = AB\varphi$ is used;
- 3) V is continuous with respect to the strong operator topology in $\mathcal{L}(\mathcal{F})$, i.e. for all $\varphi \in \mathcal{F}$ vector $V(t)\varphi$ depends on t continuously, i.e. the function $[0, +\infty) \ni t \mapsto V(t)\varphi \in \mathcal{F}$ is continuous.

Remark 1. The definition of a *C_0 -group* $(V(t))_{t \in \mathbb{R}}$ is obtained by substituting $[0, +\infty)$ with \mathbb{R} in the definition above.

Definition 2. Let $(V(t))_{t \geq 0}$ be a C_0 -semigroup in Banach space \mathcal{F} . Its *infinitesimal generator* (or just *generator*) is defined as the operator $L: D(L) \rightarrow \mathcal{F}$ with the domain

$$D(L) = \left\{ \varphi \in \mathcal{F} : \text{there exists a limit } \lim_{t \rightarrow +0} \frac{V(t)\varphi - \varphi}{t} \right\} \subset \mathcal{F},$$

and

$$L\varphi = \lim_{t \rightarrow +0} \frac{V(t)\varphi - \varphi}{t}.$$

Remark 2. It is known [3] that for each C_0 -semigroup $(V(t))_{t \geq 0}$ in Banach space \mathcal{F} , the set $D(L)$ is a dense linear subspace of \mathcal{F} . Moreover, $(L, D(L))$ is a closed linear operator that uniquely defines the semigroup $(V(t))_{t \geq 0}$. Also for each C_0 -semigroup $(V(t))_{t \geq 0}$ in Banach space there exist constants $M \geq 1$ and $\omega \in \mathbb{R}$ such that $\|V(t)\| \leq Me^{\omega t}$ for all $t \geq 0$.

Remark 3. Very often the notation $V(t) = e^{tL}$ is used. This is a good notation for several reasons. First, properties $e^{0 \cdot L} = I$ and $e^{(t+s)L} = e^{tL}e^{sL}$ are consistent with the case when L is a number or a matrix. Second, if L is a bounded linear operator then the exponent can be defined via the standard power series $e^{tL} = \sum_{n=0}^{\infty} (tL)^n/n!$ that converges in the operator norm. Finally, in general case we have $e^{tL}f = (I + tL)f + o(t)$ for all $f \in D(L)$ which again retains properties of the exponent in number and matrix case.

Now we are ready to state the Chernoff theorem. From several options (see [3, 5, 6, 29]), we choose the one given in [5] (in equivalent formulation):

Theorem 1 (P. R. CHERNOFF (1968), cf. [3, 5, 6, 29]). *Suppose that the following three conditions are met:*

1. *C_0 -semigroup $(e^{tL})_{t \geq 0}$ with generator $(L, D(L))$ in Banach space \mathcal{F} is given, such that for some $w \geq 0$ the inequality $\|e^{tL}\| \leq e^{wt}$ holds for all $t \geq 0$.*
2. *There exists a strongly continuous mapping $S: [0, +\infty) \rightarrow \mathcal{L}(\mathcal{F})$ such that $S(0) = I$ and the inequality $\|S(t)\| \leq e^{wt}$ holds for all $t \geq 0$.*

3. There exists a dense linear subspace $D \subset \mathcal{F}$ such that for all $f \in D$ there exists a limit $S'(0)f := \lim_{t \rightarrow +0} (S(t)f - f)/t$. Moreover, $S'(0)$ on D has a closure that coincides with the generator $(L, D(L))$.

Then the following statement holds:

- (C) for every $f \in \mathcal{F}$, as $n \rightarrow \infty$ we have $S(t/n)^n f \rightarrow e^{tL}f$ locally uniformly with respect to $t \geq 0$, i.e. for each $T > 0$ and each $f \in \mathcal{F}$ we have

$$\lim_{n \rightarrow \infty} \sup_{t \in [0, T]} \|S(t/n)^n f - e^{tL}f\| = 0.$$

Above $S(t/n)^n = \underbrace{S(t/n) \circ \dots \circ S(t/n)}_n$ is the composition of n copies of linear bounded operator $S(t/n)$ defined everywhere on \mathcal{F} .

Definition 3. Let C_0 -semigroup $(e^{tL})_{t \geq 0}$ with generator L in Banach space \mathcal{F} be given. The mapping $S: [0, +\infty) \rightarrow \mathcal{L}(\mathcal{F})$ is called a *Chernoff function* for operator L iff it satisfies the condition (C) of Chernoff theorem 1. In this case expressions $S(t/n)^n$ are called *Chernoff approximations* to the semigroup e^{tL} .

There is also a variant of the Chernoff theorem that allows us to prove the existence of the semigroup.

Theorem 2. (Chernoff-type theorem, corollary 5.3 from theorem 5.2 in [3]) Let \mathcal{F} be a Banach space, and $\mathcal{L}(\mathcal{F})$ be the space of all linear bounded operators on \mathcal{F} . Suppose there is a function $S: [0, +\infty) \rightarrow \mathcal{L}(\mathcal{F})$, meeting the condition $S(0) = I$, where I is the identity operator. Suppose there are numbers $M \geq 1$ and $\omega \in \mathbb{R}$ such that $\|S(t)^k\| \leq Me^{k\omega t}$ for every $t \geq 0$ and every $k \in \mathbb{N}$. Suppose the limit $\lim_{t \rightarrow +0} \frac{S(t)\varphi - \varphi}{t} =: L\varphi$ exists for every $\varphi \in \mathcal{D} \subset \mathcal{F}$, where \mathcal{D} is a dense subspace of \mathcal{F} . Suppose there is a number $\lambda_0 > \omega$ such that $(\lambda_0 I - L)(\mathcal{D})$ is a dense subspace of \mathcal{F} .

Then the closure \bar{L} of the operator L is a generator of a strongly continuous semigroup of operators $(e^{\bar{L}t})_{t \geq 0}$ given by the formula $e^{\bar{L}t}\varphi = \lim_{n \rightarrow \infty} S(t/n)^n \varphi$ where the limit exists for every $\varphi \in \mathcal{F}$ and is uniform with respect to $t \in [0, T]$ for every $T > 0$. Moreover $(e^{\bar{L}t})_{t \geq 0}$ satisfies the estimate $\|e^{\bar{L}t}\| \leq Me^{\omega t}$ for every $t \geq 0$.

Remark 4. There are several theorems that help to find out if some linear operator (or the closure of it) generates a C_0 -semigroup. Most general are Hille-Yosida theorem and Feller-Miyadera-Phillips theorem. Unfortunately both heavily use properties of the resolvent which is a kind of circulus vitiosus because we start all this activity to find the resolvent but need its properties to do it.

However there are several results that only use properties of the operator to prove that it generates a C_0 -semigroup:

- Stone's generation theorem: if A is a self-adjoint operator in Hilbert space, then iA generates a C_0 -semigroup; even more, this is a C_0 -group of unitary operators.
- Lumer-Phillips theorem: linear, closed, densely defined in Banach space, dissipative operator A with a property that $A - \lambda_0 I$ is surjective for some $\lambda_0 > 0$, generates a C_0 -semigroup of contraction operators.
- A. Yu. Neklyudov's inversion of Chernoff's theorem [31].

Remark 5. Chernoff's theorem is a deep result of functional analysis and is designed for dealing with infinite-dimensional spaces \mathcal{F} . However, it can be illustrated in one-dimensional setting in two ways, which helps to build intuition. First, one-dimensional version of Chernoff's theorem, when $\mathcal{F} = \mathcal{L}(\mathcal{F}) = \mathbb{R}$, says that if $s: [0, +\infty) \rightarrow \mathbb{R}$, $l \in \mathbb{R}$ and $s(t) = 1 + tl + o(t)$ as $t \rightarrow 0$, then $\lim_{n \rightarrow \infty} (s(t/n))^n = e^{tl}$, which is a simple fact of calculus. Second, one can see that Chernoff's theorem is an infinite-dimensional analogue of the forward Euler method for solving ordinary differential equations numerically.

3 Main result

Theorem 3. Let \mathcal{F} be real or complex Banach space, and let $\mathcal{L}(\mathcal{F})$ be the set of all linear bounded operators in \mathcal{F} . Suppose that linear operator $L: \mathcal{F} \supset D(L) \rightarrow \mathcal{F}$ generates C_0 -semigroup $(e^{tL})_{t \geq 0}$ satisfying for some constants $M \geq 1$ and $\omega \geq 0$ inequality $\|e^{tL}\| \leq Me^{\omega t}$ for all $t \geq 0$. Suppose that function $S: [0, +\infty) \rightarrow \mathcal{L}(\mathcal{F})$ is given and $\|S(t)^k\| \leq Me^{\omega tk}$ for all $t \geq 0$ and all $k = 1, 2, 3, \dots$. Let us denote the resolvent of $(L, D(L))$ by the symbol $R_\lambda = (\lambda I - L)^{-1}$ for all $\lambda \in \rho(L)$. Suppose that the number $\lambda \in \mathbb{C}$ is given and $\operatorname{Re} \lambda > \omega$. Then $\lambda \in \rho(L)$ and:

1. If for all $T > 0$ we have $\lim_{n \rightarrow \infty} \sup_{t \in [0, T]} \|e^{tL}f - (S(t/n))^n f\| = 0$ for all $f \in \mathcal{F}$, then for all $f \in \mathcal{F}$ we have

$$\lim_{n \rightarrow \infty} \left\| R_\lambda f - \int_0^\infty e^{-\lambda t} (S(t/n))^n f dt \right\| = 0. \quad (1)$$

2. If for all $T > 0$ we have $\lim_{n \rightarrow \infty} \sup_{t \in [0, T]} \|e^{tL} - (S(t/n))^n\| = 0$, then we have

$$\lim_{n \rightarrow \infty} \left\| R_\lambda - \int_0^\infty e^{-\lambda t} (S(t/n))^n dt \right\| = 0. \quad (2)$$

Доказательство. Integrals in (1) and (2) can be understood as improper Riemann integrals because for each n integrands are continuous functions of t . Moreover $\|e^{-\lambda t} (S(t/n))^n\| \leq e^{-\operatorname{Re} \lambda t} M e^{(\omega - \operatorname{Re} \lambda)t} = M e^{t(\omega - \operatorname{Re} \lambda)}$ with $\omega - \operatorname{Re} \lambda < 0$, so both integrals converge. Theorem II.1.10 from [3] says that $\lambda \in \rho(L)$ and $R_\lambda f = \int_0^\infty e^{-\lambda t} e^{tL} f dt$ for each $f \in \mathcal{F}$ with $\|R_\lambda\| \leq M/(\operatorname{Re} \lambda - \omega)$.

Let us prove item 1. Suppose $\varepsilon > 0$ and $f \in \mathcal{F}$ are given. Let us prove that there exists $n_0 \in \mathbb{N}$ such that for all $n > n_0$ the estimate $\|R_\lambda f - \int_0^\infty e^{-\lambda t} (S(t/n))^n f dt\| < \varepsilon$ holds.

Integral in (1) and $\int_0^\infty e^{-\lambda t} e^{tL} f dt$ both converge hence the integral on the right hand side of the equality

$$R_\lambda f - \int_0^\infty e^{-\lambda t} (S(t/n))^n f dt = \int_0^\infty e^{-\lambda t} (e^{tL} f - (S(t/n))^n f) dt$$

converges. Moreover, it convergences uniformly in $n \in \mathbb{N}$ due to the estimate

$$\|e^{-\lambda t} (e^{tL} - (S(t/n))^n)\| \leq e^{-\operatorname{Re} \lambda t} (\|e^{tL}\| + \|(S(t/n))^n\|) \leq e^{-\operatorname{Re} \lambda t} (M e^{\omega t} + M e^{(\omega - \operatorname{Re} \lambda)t}) \leq 2M e^{t(\omega - \operatorname{Re} \lambda)}$$

with $\omega - \operatorname{Re} \lambda < 0$.

Let us use the so-called $\varepsilon/2$ -method to prove that $\int_0^\infty < \varepsilon$, using representation $\int_0^\infty = \int_0^T + \int_T^\infty$. First, using ε , we find such $T > 0$ that $\int_T^\infty \leq \varepsilon/2$ for all n , and then for this T we find such n_0 that for all $n > n_0$ we have $\int_0^T < \varepsilon/2$. This will give us $\int_0^\infty < \varepsilon$. Indeed, we have

$$\left\| \int_0^\infty e^{-\lambda t} (e^{tL} f - (S(t/n))^n f) dt \right\| \leq \int_0^\infty 2M e^{t(\omega - \operatorname{Re} \lambda)} dt = 2M \frac{1}{\operatorname{Re} \lambda - \omega} e^{T(\omega - \operatorname{Re} \lambda)} \leq \varepsilon/2$$

for all $n \in \mathbb{N}$ and all T satisfying inequality $T \geq \max \left(0, \frac{1}{\operatorname{Re} \lambda - \omega} \ln \frac{4M}{(\operatorname{Re} \lambda - \omega)\varepsilon} \right)$. Suppose that such number T is selected.

Next, thanks to the conditions of the theorem we have $\lim_{n \rightarrow \infty} \sup_{t \in [0, T]} \|e^{tL} f - (S(t/n))^n f\| = 0$, so there exists n_0 such that for all $n > n_0$, we have

$$\begin{aligned} \left\| \int_0^T e^{-\lambda t} (e^{tL} f - (S(t/n))^n f) dt \right\| &\leq \int_0^T e^{-t \operatorname{Re} \lambda} \|e^{tL} f - (S(t/n))^n f\| dt \leq \\ &\leq T \max_{t \in [0, T]} e^{-t \operatorname{Re} \lambda} \sup_{t \in [0, T]} \|e^{tL} f - (S(t/n))^n f\| < \varepsilon/2. \end{aligned}$$

So we proved that for arbitrary $\varepsilon > 0$ there exists n_0 such that for all $n > n_0$ we have

$$\left\| R_\lambda f - \int_0^\infty e^{-\lambda t} (S(t/n))^n f dt \right\| < \varepsilon.$$

Item 1 is proved. Proof of item 2 is obtained by deleting f from the proof of item 1. \square

4 Corollary: Feynman formula for the resolvent

Assumption 1. Consider functions $a, b, c: \mathbb{R} \rightarrow \mathbb{R}$ on \mathbb{R} . Assume that $a(x) > 0$ for all $x \in \mathbb{R}$. Assume that there exists such $\beta \in (0, 1]$ that function c is bounded and Hölder continuous with Hölder exponent β , and functions $a, x \mapsto 1/a(x), b$ are bounded and Hölder continuous with Hölder exponent β with derivatives of order one and two.

Consider Banach space $C_0(\mathbb{R}, \mathbb{R})$ of all continuous functions $f: \mathbb{R} \rightarrow \mathbb{R}$ vanishing at infinity (i.e. $\lim_{\|x\| \rightarrow \infty} f(x) = 0$), with the uniform norm $\|f\| = \sup_{x \in \mathbb{R}} |f(x)|$. Consider space $C_c^2(\mathbb{R}, \mathbb{R})$ of all compactly-supported functions that are continuous with derivatives of order one and two; note that this space is a dense linear subspace in $C_0(\mathbb{R}, \mathbb{R})$.

For all $f \in C_c^2(\mathbb{R}, \mathbb{R})$ define linear operator H via the formula

$$(Hf)(x) = a(x)f''(x) + b(x)f'(x) + c(x)f(x). \quad (3)$$

Assume that the closure $(\overline{H}, D(\overline{H}))$ of operator $H: D(H) = C_c^2(\mathbb{R}, \mathbb{R}) \rightarrow C_0(\mathbb{R}, \mathbb{R})$ exists and generates a C_0 -semigroup $(e^{t\overline{H}})_{t \geq 0}$ in $C_0(\mathbb{R}, \mathbb{R})$; this assumption is fulfilled, e.g. if $a(x) \geq a_0$ for some constant $a_0 > 0$, and b, c only satisfy already mentioned conditions. Now operator $(\overline{H}, D(\overline{H}))$ is well defined.

Remark 6. Following [30], let us consider operator-valued functions S_1, S_2, S_3, S that are defined on $[0, \infty)$ and take values in $\mathcal{L}(C_0(\mathbb{R}, \mathbb{R}))$. For all $x, y \in \mathbb{R}, t > 0$ define

$$\begin{aligned} (S_1(t)f)(x) &= \frac{1}{\sqrt{4\pi ta(x)}} \int_{\mathbb{R}} \exp\left(\frac{-(x-y)^2}{4ta(x)}\right) f(y) dy, \\ (S_2(t)f)(x) &= \frac{1}{\sqrt{4\pi ta(x)}} \int_{\mathbb{R}} \exp\left(\frac{-(x-y)^2}{4ta(x)} - \frac{b(x)(x-y)}{2a(x)}\right) f(y) dy, \\ (S_3(t)f)(x) &= \exp\left(t\left(c(x) - \frac{b(x)^2}{4a(x)}\right)\right) f(x), \quad S(t) = S_3(t)S_2(t). \end{aligned} \quad (4)$$

It is proved in [30] that for all $f \in C_c^2(\mathbb{R}, \mathbb{R})$ and $t \rightarrow 0$ we have

$$(S_1(t)f)(x) = f(x) + ta(x)f''(x) + o(t),$$

$$(S(t)f)(x) = f(x) + t[a(x)f''(x) + b(x)f'(x) + c(x)f(x)] + o(t),$$

and inequality $\|S(t)\| \leq e^{wt}$ holds for all $t \geq 0$ with $w = \max(0, \sup_{x \in \mathbb{R}} C(x))$. Operator \overline{H} generates a C_0 -semigroup, so all conditions of the Chernoff theorem are fulfilled. Hence

$$(e^{t\overline{H}}f)(x) = \left(\lim_{n \rightarrow \infty} S(t/n)^n f \right)(x), \quad \text{for all } t > 0, x \in \mathbb{R}, f \in C_0(\mathbb{R}, \mathbb{R})$$

locally uniformly in t , i.e. for all $T > 0$ we have $\lim_{n \rightarrow \infty} \sup_{t \in [0, T]} \|e^{t\overline{H}}f - (S(t/n))^n f\| = 0$.

The expression for $S(t/n)^n f$ can be rewritten as follows:

$$\begin{aligned} (e^{t\overline{H}}f)(x_0) &= \lim_{n \rightarrow \infty} \underbrace{\int_{\mathbb{R}} \cdots \int_{\mathbb{R}}}_n \exp \left(\frac{t}{n} \sum_{j=1}^n \left(c(x_{j-1}) - \frac{b(x_{j-1})^2}{4a(x_{j-1})} \right) \right) \exp \left(\sum_{j=1}^n \frac{b(x_{j-1})(x_j - x_{j-1})}{2a(x_{j-1})} \right) \times \\ &\quad \times \left(\frac{\sqrt{n}}{\sqrt{4\pi t}} \right)^n \left(\prod_{j=0}^{n-1} a(x_j) \right)^{-1/2} \exp \left(-\frac{n}{4t} \sum_{j=0}^{n-1} \frac{(x_j - x_{j+1})^2}{a(x_j)} \right) f(x_n) dx_1 \dots dx_n. \end{aligned}$$

Now we can apply theorem 3 and obtain a formula for the resolvent of \overline{H} .

Theorem 4. *Under notation and assumptions from assumption 1 and remark 6, the resolvent $R_\lambda = (\lambda I - \overline{H})^{-1}$ is given for all $\lambda \in \mathbb{C}$ satisfying $\operatorname{Re} \lambda > w$, all $g \in C_0(\mathbb{R}, \mathbb{R})$, all $x_0 \in \mathbb{R}$ by the following formula:*

$$\begin{aligned} (R_\lambda g)(x_0) &= \lim_{n \rightarrow \infty} \int_0^\infty e^{-\lambda t} \left[\underbrace{\int_{\mathbb{R}} \cdots \int_{\mathbb{R}}}_n \exp \left(\frac{t}{n} \sum_{j=1}^n \left(c(x_{j-1}) - \frac{b(x_{j-1})^2}{4a(x_{j-1})} \right) \right) \exp \left(\sum_{j=1}^n \frac{b(x_{j-1})(x_j - x_{j-1})}{2a(x_{j-1})} \right) \times \right. \\ &\quad \times \left. \left(\frac{\sqrt{n}}{\sqrt{4\pi t}} \right)^n \left(\prod_{j=0}^{n-1} a(x_j) \right)^{-1/2} \exp \left(-\frac{n}{4t} \sum_{j=0}^{n-1} \frac{(x_j - x_{j+1})^2}{a(x_j)} \right) g(x_n) dx_1 \dots dx_n \right] dt, \end{aligned}$$

where the limit $\lim_{n \rightarrow \infty}$ exists uniformly in $x_0 \in \mathbb{R}$.

Доказательство. Let us check conditions of theorem 3. Set $\mathcal{F} = C_0(\mathbb{R}, \mathbb{R})$ with the uniform norm $\|f\| = \sup_{x \in \mathbb{R}} f(x)$. Consider $L = \overline{H}$ defined via (3), $D(L) = D(\overline{H})$, $\omega = w$, $M = 1$. Consider $S(t)$ defined in (4). We already have $\|S(t)\| \leq e^{wt}$, which implies $\|S(t)^k\| \leq e^{wt k}$. Condition $\lim_{n \rightarrow \infty} \sup_{t \in [0, T]} \|e^{t\overline{H}}f - (S(t/n))^n f\| = 0$ for all $f \in \mathcal{F}$ and all $T > 0$ is true thanks to [30], where the Chernoff theorem is used. The proof is finished due to item 1 of theorem 3. \square

Remark 7. Note that for an arbitrary linear operator $(L, D(L))$ the resolvent $(\lambda I - L)^{-1}$ and the semigroup e^{tL} , if they exist, are defined by $(L, D(L))$ uniquely. Meanwhile, there are many Chernoff functions for the same operator $(L, D(L))$, so there are many Chernoff approximations for $(\lambda I - L)^{-1}$ and e^{tL} . This gives us some freedom in constructing such approximations. The representation for the resolvent proposed in theorem 4 is only one of the representations that can be obtained via the Chernoff theorem.

5 Corollary: representing solution of ODEs via Feynman formula

There is no standard well known method of expressing the solution of ODE $-a(x)f''(x) - b(x)f'(x) - c(x)f(x) + \lambda f(x) = g(x)$, $x \in \mathbb{R}$ in terms of variable coefficients a, b, c, g and constant λ . Meanwhile, our method gives a formula for the solution, because $f = (\lambda I - L)^{-1}g$ for L given by $(Lf)(x) = a(x)f''(x) + b(x)f'(x) + c(x)f(x)$. We will rewrite $\lambda f - Lf = g$ as $Lf - \lambda f = -g$ because it is easier to follow the idea. Please allow us to make the statement of the theorem a bit wordy to keep it self-contained.

Theorem 5. Consider second order ordinary differential equation for function $f: \mathbb{R} \rightarrow \mathbb{R}$

$$a(x)f''(x) + b(x)f'(x) + (c(x) - \lambda)f(x) = -g(x) \text{ for all } x \in \mathbb{R}, \quad (5)$$

where functions $a, b, c, g: \mathbb{R} \rightarrow \mathbb{R}$ are known parameters and number $\lambda \in \mathbb{C}$ is also a known parameter. Assume that there exists constant $a_0 > 0$ such that $a(x) > a_0$ for all $x \in \mathbb{R}$. Assume that there exists $\beta \in (0, 1]$ such that function c is bounded and Hölder continuous with Hölder exponent β , and functions $a, x \mapsto 1/a(x), b$ are bounded and Hölder continuous with Hölder exponent β with derivatives of order one and two. Assume that function g is continuous and vanishes at infinity. Assume that $\mathbb{R} \ni \lambda > \max(0, \sup_{x \in \mathbb{R}} c(x))$.

Then for (5) there exists a unique continuous and vanishing at infinity solution f given for all $x_0 \in \mathbb{R}$ by the formula

$$f(x_0) = \lim_{n \rightarrow \infty} \int_0^\infty e^{-\lambda t} \left[\underbrace{\int_{\mathbb{R}} \cdots \int_{\mathbb{R}}}_n \exp \left(\frac{t}{n} \sum_{j=1}^n \left(c(x_{j-1}) - \frac{b(x_{j-1})^2}{4a(x_{j-1})} \right) \right) \exp \left(\sum_{j=1}^n \frac{b(x_{j-1})(x_j - x_{j-1})}{2a(x_{j-1})} \right) \times \right. \\ \left. \times \left(\frac{\sqrt{n}}{\sqrt{4\pi t}} \right)^n \left(\prod_{j=0}^{n-1} a(x_j) \right)^{-1/2} \exp \left(-\frac{n}{4t} \sum_{j=0}^{n-1} \frac{(x_j - x_{j+1})^2}{a(x_j)} \right) g(x_n) dx_1 \dots dx_n \right] dt,$$

where the limit $\lim_{n \rightarrow \infty}$ exists uniformly in $x_0 \in \mathbb{R}$.

Доказательство. In theorem 4 set $f = R_\lambda g$. □

Remark 8. This reasoning also works in the multi-dimensional situation for $x \in \mathbb{R}^d$, where we have an elliptic PDE instead of ODE.

6 Corollary: translation-based formula as a method of solving ODEs

Another Chernoff approximations for the same semigroup are known, these approximations do not involve multiple integrals but use multiple shifts instead [12]. For these approximations error bounds are known. Rate of convergence of Chernoff approximations is given in [19] for the general case of arbitrary semigroup, and also in this particular case of translation-based approximations [18] for the semigroup that is discussed in the next theorem. The word "translation" is used because for $a(x) = a_0 \equiv \text{const}$ operator $f \mapsto [x \mapsto f(x + 2\sqrt{a(x)t})]$ is indeed a translation (shift) of f by the value $2\sqrt{a_0 t}$.

Let us use symbol $UC_b(\mathbb{R})$ to denote Banach space of all bounded and uniformly continuous functions $f: \mathbb{R} \rightarrow \mathbb{R}$ with the uniform norm $\|f\| = \sup_{x \in \mathbb{R}} |f(x)|$. Let us use symbol $C_b^\infty(\mathbb{R})$ for the subspace of $UC_b(\mathbb{R})$ consisting of all infinitely differentiable functions that are bounded and have bounded derivatives of all orders.

Theorem 6. Suppose that functions $a, b, c \in UC_b(\mathbb{R})$ are bounded with their derivatives up to order 3, and there exists such a constant $a_0 > 0$ that estimate $\inf_{x \in \mathbb{R}} a(x) \geq a_0 > 0$ is satisfied for all $x \in \mathbb{R}$. For each function $\phi \in C_b^\infty(\mathbb{R}) = D(A)$ define $A\phi = a\phi'' + b\phi' + c\phi$. For each $t \geq 0$, each $x \in \mathbb{R}$ and each $f \in UC_b(\mathbb{R})$ define

$$(S(t)f)(x) = \frac{1}{4}f\left(x + 2\sqrt{a(x)t}\right) + \frac{1}{4}f\left(x - 2\sqrt{a(x)t}\right) + \frac{1}{2}f(x + 2b(x)t) + tc(x)f(x). \quad (6)$$

Assume also that $\mathbb{R} \ni \lambda > \sup_{x \in \mathbb{R}} |c(x)| = \|c\|$. Then:

1. Closure \bar{A} of operator A generates a C_0 -semigroup in $UC_b(\mathbb{R})$.
2. For each $g \in UC_b(\mathbb{R})$ the solution $f: \mathbb{R} \rightarrow \mathbb{R}$ of the equation

$$a(x)f''(x) + b(x)f'(x) + (c(x) - \lambda)f(x) = -g(x) \text{ for all } x \in \mathbb{R}, \quad (7)$$

exists, is unique in $UC_b(\mathbb{R})$ and is given for all $x \in \mathbb{R}$ by the formula

$$f(x) = \int_0^\infty e^{-\lambda t} \left(e^{\bar{A}} g \right) (x) dt = \lim_{n \rightarrow \infty} \int_0^\infty e^{-\lambda t} ((S(t/n))^n g) (x) dt, \quad (8)$$

where $S(t/n)$ is obtained by substitution of t with t/n in (6), and $(S(t/n))^n$ is the composition of n copies of linear bounded operator $S(t/n)$.

Suppose additionally that function g is bounded with derivatives up to order 5. Then:

3. There exist nonnegative constants C_0, C_1, \dots, C_4 such that for all $t > 0$ and all $n \in \mathbb{N}$ the following inequality holds:

$$\|S(t/n)^n g - e^{t\bar{A}} g\| \leq \frac{t^2 e^{\|c\|t}}{n} (C_0 \|g\| + C_1 \|g'\| + C_2 \|g''\| + C_3 \|g'''\| + C_4 \|g^{(IV)}\|). \quad (9)$$

4. Error bound in (8) for all $n \in \mathbb{N}$ is given by inequality

$$\sup_{x \in \mathbb{R}} \left| f(x) - \int_0^\infty e^{-\lambda t} ((S(t/n))^n g) (x) dt \right| \leq \frac{2C_g}{n \cdot (\lambda - \|c\|)^3},$$

where $C_g = C_0 \|g\| + C_1 \|g'\| + C_2 \|g''\| + C_3 \|g'''\| + C_4 \|g^{(IV)}\|$.

5. Integral in item 2 can be calculated over $[0, T]$ instead of $[0, \infty)$ with controlled level of error. This means that for each $\varepsilon > 0$ there exists $T = \max \left(0, \frac{1}{\lambda - \|c\|} \ln \frac{2}{(\lambda - \|c\|)\varepsilon} \right)$ such that for all $n \in \mathbb{N}$ we have

$$\sup_{x \in \mathbb{R}} \left| f(x) - \int_0^T e^{-\lambda t} ((S(t/n))^n g) (x) dt \right| \leq \frac{2C_g}{n \cdot (\lambda - \|c\|)^3} + \varepsilon.$$

Доказательство. Item 1 follows from theorem 4.2 in [18]. Item 2 is a particular case of the main result of the paper, theorem 3. Item 3 follows is example 4.2 in [18]. Item 4 follows from items 2 and 3 with simple estimate

$$\begin{aligned} & \left\| \int_0^\infty e^{-\lambda t} \left(e^{\bar{A}} g \right) (x) dt - \int_0^\infty e^{-\lambda t} ((S(t/n))^n g) (x) dt \right\| \leq \\ & \leq \int_0^\infty e^{-\lambda t} \left\| e^{\bar{A}} g - (S(t/n))^n g \right\| dt \leq \int_0^\infty e^{(\|c\| - \lambda)t} \frac{t^2}{n} C_g dt = \frac{2C_g}{n \cdot (\lambda - \|c\|)^3}. \end{aligned}$$

Item 5 (by repeating the reasoning in the first part of the proof of theorem 3) follows from item 4 and the well known fact that the semigroup $(e^{t\bar{A}})_{t \geq 0}$ is a quasi-contraction, i.e. in estimate for norm $\|e^{t\bar{A}}\| \leq M e^{\omega t}$ it is possible to set $M = 1$, $\omega = \|c\|$. \square

Remark 9. Independently of Chernoff function used (is it based on integral operators as in theorem 5 or on translation operators as in theorem 6), Chernoff approximations are allowing to calculate value of the solution in only one point of the domain of solution (in one point $x \in \mathbb{R}$ in our examples). Meanwhile methods based on a computational grid calculate values of the solution in all points of the computational grid. Moreover, values of Chernoff approximations at different points of the domain can be calculated in parallel, using multi-core processors and GPU which is an advantage of this approach.

Acknowledgments. Author is thankful to Oleg Galkin, Denis Mineev and Polina Panteleeva for comments on the manuscript. The work (except theorem 5) was supported by the Russian Science Foundation (project 23-71-30008). Theorem 5 was obtained in IITP (Dobrushin's Math. Lab.).

Список литературы

- [1] W. Arendt. Semigroups and evolution equations: Functional calculus, regularity and kernel estimates. Handbook of Differential Equations: Evolutionary Equations, vol 1, 2002, pp. 1-85
- [2] W. Arendt, C.J.K. Batty, M. Hieber, F. Neubrander. Cauchy Problems. Vector-valued Laplace Transforms and Cauchy Problems. Monographs in Mathematics, vol 96. Birkhäuser, Basel, 2001.
- [3] K.-J. Engel, R. Nagel. One-Parameter Semigroups for Linear Evolution Equations. — Springer, 2000.
- [4] J.A. Goldstein. Semigroups of Linear Operators and Applications. — Dover Publications, New York, 2017.
- [5] V.I. Bogachev, O.G. Smolyanov. Real and Functional Analysis. — Springer, 2020.
- [6] P.R. Chernoff. Note on product formulas for operator semigroups. // J. Funct. Anal. 2:2 (1968), 238–242.
- [7] Ya.A. Butko. The method of Chernoff approximation. Springer Proceedings in Mathematics and Statistics. Volume 325. — Springer, Cham, 2020. Pp. 19–46.
- [8] I.D. Remizov. Feynman and Quasi-Feynman Formulas for Evolution Equations// Doklady Mathematics, 96:2 (2017), 433-437
- [9] I.D. Remizov. Formulas that represent Cauchy problem solution for momentum and position Schrödinger equation. — Potential Analysis, 52 (2020), 339-370
- [10] I.D. Remizov. New Method for Constructing Chernoff Functions// Differential Equations, 53:4 (2017), 566-570
- [11] I.D. Remizov. Quasi-Feynman formulas — a method of obtaining the evolution operator for the Schrödinger equation. // J. Funct. Anal. 270:12 (2016), 4540–4557.
- [12] I.D. Remizov. Approximations to the solution of Cauchy problem for a linear evolution equation via the space shift operator (second-order equation example). // Applied Mathematics and Computation. 328 (2018), 243–246.
- [13] I.D. Remizov. Solution-giving formula to Cauchy problem for multidimensional parabolic equation with variable coefficients. // Journal of Mathematical Physics. 60:7 (2019), 071505.
- [14] Yu. N. Orlov, V. Zh. Sakbaev, O. G. Smolyanov. Feynman Formulas and the Law of Large Numbers for Random One-Parameter Semigroups// Proc. Steklov Inst. Math., 306 (2019), 196-211.
- [15] V. Zh. Sakbaev, N. V. Tsoy. Analogue of Chernoff theorem for cylindrical pseudomeasures// Lobachevskii J. Math., 41:12 (2020), 2369-2382
- [16] R. Sh. Kalmetiev, Yu. N. Orlov, V. Zh. Sakbaev. Averaging of random affine transformations of functions domain// Ufa Math. J., 15:2 (2023), 3-12

- [17] K. A. Dragunova, N. Nikbakht, I. D. Remizov. Numerical Study of the Rate of Convergence of Chernoff Approximations to Solutions of the Heat Equation.// Zhurnal Srednevolzhskogo matematicheskogo obshchestva. 25:4(2023), 255-272
- [18] Galkin, O.E., Remizov, I.D. Upper and lower estimates for rate of convergence in the Chernoff product formula for semigroups of operators. Israel Journal of Mathematics (2024), pp. 1-15. <https://doi.org/10.1007/s11856-024-2678-x>
- [19] O.E. Galkin, I.D. Remizov. Rate of Convergence of Chernoff Approximations of operator C_0 -semigroups.// Mathematical Notes, Math. Notes, 111:2 (2022), 305-307
- [20] A. Gomilko, S. Kosowicz, Yu. Tomilov. A general approach to approximation theory of operator semigroups. // Journal de Mathématiques Pures et Appliquées. 127 (2019), 216–267.
- [21] A.V. Vedenin, V.S. Voevodkin, V.D. Galkin, E.Yu. Karatetskaya, I.D. Remizov. Speed of Convergence of Chernoff Approximations to Solutions of Evolution Equations. // Math. Notes, 108:3 (2020), 451–456.
- [22] V.A. Zagrebnov. Notes on the Chernoff product formula. // J. Funct. Anal. 279:7 (2020), 108696.
- [23] V.A. Zagrebnov. Comments on the Chernoff estimate// Nanosystems: Physics, Chemistry, Mathematics, 13:1 (2022), 17–23.
- [24] R. Sh. Kalmetev. Approximate solution of multidimensional Kolmogorov equation using Feynman-Chernoff iterations// Keldysh Institute preprints, 2023, 021, 15 pp.
- [25] R. Sh. Kalmetiev, Yu. N. Orlov, V. Zh. Sakbaev. Chernoff iterations as an averaging method for random affine transformations// Comput. Math. Math. Phys., 62:6 (2022), 996-1006
- [26] R.P. Feynman. Space-time approach to nonrelativistic quantum mechanics. — Rev. Mod. Phys., 20 (1948), 367-387.
- [27] R.P. Feynman. An operation calculus having applications in quantum electrodynamics. — Phys. Rev. 84 (1951), 108-128.
- [28] S. Mazzucchi. Mathematical Feynman path integrals and their applications, World Scientific, 2009.
- [29] G.W. Johnson, M.L. Lapidus. The Feynman Integral and Feynman's Operational Calculus. Clarendon Press, Oxford, 2000.
- [30] Ya.A. Butko, M. Grothaus, O.G. Smolyanov. Lagrangian Feynman formulas for second-order parabolic equations in bounded and unbounded domains. — Infinite Dimansional Analysis, Quantum Probability and Related Topics, vol. 13, No. 3 (2010), 377-392.
- [31] A. Yu. Neklyudov. Inversion of Chernoff's Theorem// Math. Notes, 83:4 (2008), 530-538