

# Cremona transformations, diffeomorphisms of surfaces and approximation by $(-1)$ -curves

Frédéric Mangolte (Angers)  
joint work with  
János Kollár (Princeton)

Москва, 5 декабря 2012 г.

# Approximating by algebraic maps

Weierstrass (1885)

Every  $C^\infty$ -map  $f: \mathbb{R} \rightarrow \mathbb{R}$  is approximated by polynomials.

# Approximating by algebraic maps

Weierstrass (1885)

Every  $C^\infty$ -map  $f: \mathbb{R} \rightarrow \mathbb{R}$  is approximated by polynomials.

$$S^1 := \{(x, y) \in \mathbb{R}^2, x^2 + y^2 = 1\}$$

Every  $C^\infty$ -map  $f: S^1 \rightarrow S^1$  is approximated by rational maps

$$\Phi: (x, y) \mapsto \left( \frac{p_1(x, y)}{q_1(x, y)}, \frac{p_2(x, y)}{q_2(x, y)} \right)$$

# Approximating by algebraic maps

Weierstrass (1885)

Every  $C^\infty$ -map  $f: \mathbb{R} \rightarrow \mathbb{R}$  is approximated by polynomials.

$S^1 := \{(x, y) \in \mathbb{R}^2, x^2 + y^2 = 1\}$

Every  $C^\infty$ -map  $f: S^1 \rightarrow S^1$  is approximated by rational maps

$$\Phi: (x, y) \mapsto \left( \frac{p_1(x, y)}{q_1(x, y)}, \frac{p_2(x, y)}{q_2(x, y)} \right)$$

$X$  real algebraic variety

Is a given  $C^\infty$ -map  $f: S^1 \rightarrow X$  approximated by rational curves?

[Recall:  $\mathbb{P}^1(\mathbb{R}) \sim S^1$ .]

# Rational curves


## Exemple

$$\begin{aligned} f: \mathbb{R} &\longrightarrow \mathbb{R}^2 \\ t &\longmapsto (t^2 + 1, t(t^2 + 1)) \end{aligned}$$

# Rational curves

## Exemple

$$\begin{aligned} f: \mathbb{R} &\longrightarrow \mathbb{R}^2 \\ t &\longmapsto (t^2 + 1, t(t^2 + 1)) \end{aligned}$$



# Rational curves


## Exemple

$$\begin{aligned} f: \mathbb{R} &\longrightarrow \mathbb{R}^2 \\ t &\longmapsto (t^2 + 1, t(t^2 + 1)) \end{aligned}$$

Compactification

$$\mathbb{R} \hookrightarrow \mathbb{P}^1(\mathbb{R}) \xrightarrow{\hat{f}} X \xleftarrow{\text{bir}} \mathbb{R}^2$$

$X$  rational surface



# Approximating by rational curves

$X$  nonsingular real algebraic variety

$\mathcal{C}^\infty(S^1, X) :=$  space of maps endowed with the  $\mathcal{C}^\infty$ -topology

$\mathcal{A}_X \subset \mathcal{C}^\infty(S^1, X) :=$  subset of rational curves  $\mathbb{P}^1(\mathbb{R}) \rightarrow X$

## Definition

Let  $f \in \mathcal{C}^\infty(S^1, X)$  be a  $\mathcal{C}^\infty$ -map

$f$  is approximated by rational curves

$\Leftrightarrow$

$f \in \overline{\mathcal{A}_X}$ .

## Theorem (Bochnak, Kucharz, 1999)


Let  $X$  be a nonsingular real *rational* variety, then any  $\mathcal{C}^\infty$ -map  $\mathbb{P}^1(\mathbb{R}) \rightarrow X$  is approximated by rational curves.



# Smoothness?

## Remark

$$\begin{aligned} f: \mathbb{R} &\longrightarrow \mathbb{R}^2 \\ t &\longmapsto (t^2 + 1, t(t^2 + 1)) \end{aligned}$$




# Smoothness?

## Remark

$$\begin{aligned} f: \mathbb{C} &\longrightarrow \mathbb{C}^2 \\ t &\longmapsto (t^2 + 1, t(t^2 + 1)) \end{aligned}$$

$$f(\mathbb{R}) \subsetneq f(\mathbb{C}) \cap \mathbb{R}^2$$




# Smoothness?

## Remark

$$\begin{aligned} f: \mathbb{C} &\longrightarrow \mathbb{C}^2 \\ t &\longmapsto (t^2 + 1, t(t^2 + 1)) \end{aligned}$$

$$f(\mathbb{R}) \subsetneq f(\mathbb{C}) \cap \mathbb{R}^2$$

$$\text{Indeed: } f(i) = f(-i) = (0, 0)$$



# Smoothness?

## Remark


$$\begin{aligned} f: \mathbb{C} &\longrightarrow \mathbb{C}^2 \\ t &\longmapsto (t^2 + 1, t(t^2 + 1)) \end{aligned}$$

$$y^2 = x^2(x - 1)$$

$$\mathbb{R} \xrightarrow{f} \mathbb{R}^2$$

$$\downarrow$$

$$\mathbb{C} \xrightarrow{f} \mathbb{C}^2$$



# Approximating by smooth rational curves

$\mathcal{B}_X \subset \mathcal{A}_X \subset \mathcal{C}^\infty(S^1, X) :=$  subset of smooth rational curves  $\mathbb{P}^1(\mathbb{R}) \rightarrow X$

## Definition

$f \in \mathcal{C}^\infty(S^1, X)$  is approximated by smooth rational curves

$\Leftrightarrow$

$f \in \overline{\mathcal{B}_X}$ .

# Approximating by smooth rational curves

$\mathcal{B}_X \subset \mathcal{A}_X \subset \mathcal{C}^\infty(S^1, X) :=$  subset of smooth rational curves  $\mathbb{P}^1(\mathbb{R}) \rightarrow X$

## Definition

$f \in \mathcal{C}^\infty(S^1, X)$  is approximated by smooth rational curves

$\Leftrightarrow$

$f \in \overline{\mathcal{B}_X}$ .

## Main Theorem

*Any embedded circle in a nonsingular real rational **surface** admits a  $\mathcal{C}^\infty$ -approximation by smooth rational curves.*

# Approximating by smooth rational curves

$\mathcal{B}_X \subset \mathcal{A}_X \subset \mathcal{C}^\infty(S^1, X) :=$  subset of smooth rational curves  $\mathbb{P}^1(\mathbb{R}) \rightarrow X$

## Definition

$f \in \mathcal{C}^\infty(S^1, X)$  is approximated by smooth rational curves

$\Leftrightarrow$

$f \in \overline{\mathcal{B}_X}$ .

## Main Theorem

*Any embedded circle in a nonsingular real rational **surface** admits a  $\mathcal{C}^\infty$ -approximation by smooth rational curves.*

## Corollary

*Let  $X$  be a nonsingular real rational variety, then any embedded circle is approximated by smooth rational curves.*

# Real rational surfaces

## Theorem (Comessatti, 1914)

$X$  *orientable nonsingular real rational surface*

$\Rightarrow X$  *diffeomorphic to the sphere  $S^2$  or to the torus  $S^1 \times S^1$*



# Real rational surfaces

## Theorem (Comessatti, 1914)

$X$  *orientable nonsingular real rational surface*

$\Rightarrow X$  *diffeomorphic to the sphere  $S^2$  or to the torus  $S^1 \times S^1$*

Conversely:

$S^2 \sim$  rational model  $\{x^2 + y^2 + z^2 = 1\} \subset \mathbb{R}^3$

$S^1 \times S^1 \sim$  rational model  $\{x^2 + y^2 = z^2 + t^2 = 1\} \subset \mathbb{R}^4$

$\mathbb{RP}^2 \sim$  rational model  $\mathbb{P}^2(\mathbb{R})$

$\#^h \mathbb{RP}^2 \sim$  rational model  $B_{p_1, p_2, \dots, p_{h-1}} \mathbb{P}^2(\mathbb{R})$  (blow-up at  $h - 1$  points)

# Classification of rational models

$$S^1 := \{(x, y) \in \mathbb{R}^2, x^2 + y^2 = 1\}$$

Real algebraic manifold := compact connected submanifold of  $\mathbb{R}^n$  defined by real polynomial equations, for some  $n$ .

$X, Y$  real algebraic manifolds,  $f: X \rightarrow Y$  map

$f$  **algebraic** := (i) real rational (ii) defined  $\forall x \in X$

$f$  **isomorphism** := (i) algebraic, (ii)  $f^{-1}$  exists (iii)  $f^{-1}$  algebraic

# Classification of rational models

$$S^1 := \{(x, y) \in \mathbb{R}^2, x^2 + y^2 = 1\}$$

Real algebraic manifold := compact connected submanifold of  $\mathbb{R}^n$  defined by real polynomial equations, for some  $n$ .

$X, Y$  real algebraic manifolds,  $f: X \rightarrow Y$  map

$f$  **algebraic** := (i) real rational (ii) defined  $\forall x \in X$

$f$  **isomorphism** := (i) algebraic, (ii)  $f^{-1}$  exists (iii)  $f^{-1}$  algebraic



$x \mapsto x + x^3$  algebraic self-diffeomorphism of  $\mathbb{R}$  is not an isomorphism.  
(Implicit function Theorem does not hold in the algebraic setting.)

# Classification of rational models

$$S^1 := \{(x, y) \in \mathbb{R}^2, x^2 + y^2 = 1\}$$

Real algebraic manifold := compact connected submanifold of  $\mathbb{R}^n$  defined by real polynomial equations, for some  $n$ .

$X, Y$  real algebraic manifolds,  $f: X \rightarrow Y$  map

$f$  **algebraic** := (i) real rational (ii) defined  $\forall x \in X$

$f$  **isomorphism** := (i) algebraic, (ii)  $f^{-1}$  exists (iii)  $f^{-1}$  algebraic



$x \mapsto x + x^3$  algebraic self-diffeomorphism of  $\mathbb{R}$  is not an isomorphism.  
(Implicit function Theorem does not hold in the algebraic setting.)

## Theorem (Biswas, Huisman, 2007)

*Two nonsingular real rational surfaces are isomorphic if and only if they are diffeomorphic.*

# Real $(-1)$ -curves

Let  $L \subset X$  be a real algebraic curve on a real algebraic surface

## Definition

$L$  is a  $(-1)$ -curve iff

$\exists$  birational morphism  $\pi: X \rightarrow Y$  such that  $\pi(L)$  is a smooth point on  $Y$  and  $\pi$  restricted to  $X \setminus L \rightarrow Y \setminus \pi(L)$  is an isomorphism.

By Castelnuovo's criterium,  $\exists$  such a birational morphism  $\pi: X \rightarrow Y$  iff there exists a real algebraic surface  $X'$  and a real algebraic isomorphism  $\Phi: X \rightarrow X'$  such that  $L' := \Phi(L)$  is rational, irreducible and nonsingular and  $L' \cdot L' = -1$  (self-intersection over complex points).

# Approximating by $(-1)$ -curves

## Theorem

*$X$  nonsingular real rational surface and  $L \subset X$  a nonsingular curve, the following assertions are equivalent:*

- ❶  *$X$  is nonorientable near  $L$  and one of the following is satisfied:  
     $X \setminus L$  is a punctured sphere, or  
     $X \setminus L$  is a punctured torus, or  
     $X \setminus L$  is nonorientable.*
- ❷  *$L$  is homotopic to a  $(-1)$ -curve*
- ❸  *$L$  admits  $\mathcal{C}^\infty$ -approximation by  $(-1)$ -curves*

# Proof of the approximation by smooth rational curves

- 1 Classify all topological pairs  $(K, S)$  such that  $S$  closed surface either nonorientable or of genus  $\leq 1$  and  $K$  embedded circle in  $S$

# Proof of the approximation by smooth rational curves

- 1 Classify all topological pairs  $(K, S)$  such that  $S$  closed surface either nonorientable or of genus  $\leq 1$  and  $K$  embedded circle in  $S$
- 2 Construct rational models for each topological pair



# Proof of the approximation by smooth rational curves

- 1 Classify all topological pairs  $(K, S)$  such that  $S$  closed surface either nonorientable or of genus  $\leq 1$  and  $K$  embedded circle in  $S$
- 2 Construct rational models for each topological pair
- 3 Get:  $\forall$  pair  $(K, S)$ ,  $\exists X$  nonsingular real rational surface  
 $\exists \varphi: S \xrightarrow{\sim} X$  diffeomorphism  
such that  $L := \varphi(K) \subset X$  nonsingular real rational curve

# Proof of the approximation by smooth rational curves

- 1 Classify all topological pairs  $(K, S)$  such that  $S$  closed surface either nonorientable or of genus  $\leq 1$  and  $K$  embedded circle in  $S$
- 2 Construct rational models for each topological pair
- 3 Get:  $\forall$  pair  $(K, S)$ ,  $\exists X$  nonsingular real rational surface  
 $\exists \varphi: S \xrightarrow{\sim} X$  diffeomorphism  
such that  $L := \varphi(K) \subset X$  nonsingular real rational curve
- 4 The rest of the talk is devoted to deduce the approximation result!

# Density of $\text{Aut}(X)$

Recall:  $f: X \rightarrow X$  automorphism  $\Leftrightarrow$

(i)  $f$  birational map, (ii)  $f$  is a self-diffeomorphism on the real locus

$\text{Aut}(X) :=$  group of real algebraic automorphisms  $X \rightarrow X$

Remark: let  $V|_{\mathbb{R}}$  such that  $V(\mathbb{R}) = X$ , then

$\text{Aut}_{\mathbb{R}}(V) \subset \text{Aut}(X) \subset \text{Bir}_{\mathbb{R}}(V)$

# Density of $\text{Aut}(X)$

Recall:  $f: X \rightarrow X$  automorphism  $\Leftrightarrow$

(i)  $f$  birational map, (ii)  $f$  is a self-diffeomorphism on the real locus

$\text{Aut}(X) :=$  group of real algebraic automorphisms  $X \rightarrow X$

Remark: let  $V|_{\mathbb{R}}$  such that  $V(\mathbb{R}) = X$ , then

$\text{Aut}_{\mathbb{R}}(V) \subset \text{Aut}(X) \subset \text{Bir}_{\mathbb{R}}(V)$

## Theorem (Kollár, M. 2009)

- $S = S^2$ ,  $S^1 \times S^1$ , or any non-orientable surface,  
 $\Rightarrow \exists$  real algebraic model  $X \sim S$  such that  $\overline{\text{Aut}(X)} = \text{Diff}(X)$   
for the  $C^\infty$ -topology.

# Density of $\text{Aut}(X)$

Recall:  $f: X \rightarrow X$  automorphism  $\Leftrightarrow$

(i)  $f$  birational map, (ii)  $f$  is a self-diffeomorphism on the real locus

$\text{Aut}(X) :=$  group of real algebraic automorphisms  $X \rightarrow X$

Remark: let  $V|_{\mathbb{R}}$  such that  $V(\mathbb{R}) = X$ , then

$\text{Aut}_{\mathbb{R}}(V) \subset \text{Aut}(X) \subset \text{Bir}_{\mathbb{R}}(V)$

## Theorem (Kollár, M. 2009)

- $S = S^2$ ,  $S^1 \times S^1$ , or any non-orientable surface,  
 $\Rightarrow \exists$  real algebraic model  $X \sim S$  such that  $\overline{\text{Aut}(X)} = \text{Diff}(X)$   
for the  $C^\infty$ -topology.
- $S$  any orientable surface of genus  $\geq 2$ ,  
 $\Rightarrow \forall$  model  $X \sim S$ ,  $\text{Aut}(X)$  is *not* dense in  $\text{Diff}(X)$ ,  
even for the  $C^0$ -topology.

# Cremona transformation (around 1860)

On  $\mathbb{P}^3$  take  $(x : y : z : t) \mapsto (\frac{1}{x} : \frac{1}{y} : \frac{1}{z} : \frac{1}{t}) = (yzt : ztx : txy : xyz)$

Base locus = 6 edges of a tetraedron  $T$ .

Move vertices to  $(1, \pm i, 0, 0), (0, 0, 1, \pm i)$ , get:

$$\sigma : (x : y : z : t) \mapsto ((x^2 + y^2)z : (x^2 + y^2)t : (z^2 + t^2)x : (z^2 + t^2)y)$$

$\sigma$  diffeomorphism of  $\mathbb{P}^3(\mathbb{C}) \setminus T$

# Cremona transformation (around 1860)

On  $\mathbb{P}^3$  take  $(x : y : z : t) \mapsto (\frac{1}{x} : \frac{1}{y} : \frac{1}{z} : \frac{1}{t}) = (yzt : ztx : txy : xyz)$

Base locus = 6 edges of a tetraedron  $T$ .

Move vertices to  $(1, \pm i, 0, 0), (0, 0, 1, \pm i)$ , get:

$$\sigma : (x : y : z : t) \mapsto ((x^2 + y^2)z : (x^2 + y^2)t : (z^2 + t^2)x : (z^2 + t^2)y)$$

$\sigma$  diffeomorphism of  $\mathbb{P}^3(\mathbb{C}) \setminus T$

Each quadric

$$Q_{abcdef} := a(x^2 + y^2) + b(z^2 + t^2) + cxz + dyt + ext + fyz$$

(i) passes through the vertices of  $T$ ,

(ii) has no real points on  $T$ .

# Cremona transformation (around 1860)

On  $\mathbb{P}^3$  take  $(x : y : z : t) \mapsto (\frac{1}{x} : \frac{1}{y} : \frac{1}{z} : \frac{1}{t}) = (yzt : ztx : txy : xyz)$

Base locus = 6 edges of a tetraedron  $T$ .

Move vertices to  $(1, \pm i, 0, 0), (0, 0, 1, \pm i)$ , get:

$$\sigma : (x : y : z : t) \mapsto ((x^2 + y^2)z : (x^2 + y^2)t : (z^2 + t^2)x : (z^2 + t^2)y)$$

$\sigma$  diffeomorphism of  $\mathbb{P}^3(\mathbb{C}) \setminus T$

Each quadric

$$Q_{abcdef} := a(x^2 + y^2) + b(z^2 + t^2) + cxz + dyt + ext + fyz$$

(i) passes through the vertices of  $T$ ,

(ii) has no real points on  $T$ .

$$\sigma : Q_{abcdef}(\mathbb{R}) \xrightarrow{\cong} Q_{abcdfe}(\mathbb{R})$$



# Action on spheres

$$S^2 := \{(x, y, z) \in \mathbb{R}^3, x^2 + y^2 + z^2 = 1\}$$

$$Q_0 := \{(x, y, z, t) \in \mathbb{P}^3, x^2 + y^2 + z^2 - t^2 = 0\}$$

Take  $Q_{abcdef}$  with  $Q_{abcdef}(\mathbb{R}) \sim S^2$ ,  $\Rightarrow Q_{abcdfe}(\mathbb{R}) \sim S^2$ ,  
then both are equivalent to  $Q_0$  up to linear change of coordinates.

Get:  $\sigma_{abcdef}: S^2 \xrightarrow{\cong} S^2$ , well defined up to  $O(3, 1)$ .

## Theorem

*The Cremona transformations with imaginary base points and  $O(3, 1)$  generate  $\text{Aut}(S^2)$  which is dense in  $\text{Diff}(S^2)$ .*

## Theorem (Lukackiř 1977)

*$SO(m + 1, 1)$  is a maximal closed subgroup of  $\text{Diff}_0(S^m)$ .*

Rational models of non-orientable surfaces:  $(\chi(R_g) = 2 - g)$

$R_g \sim B_{p_1, \dots, p_g} S^2$ , the sphere blown-up at  $g$  points

Let  $q_1, \dots, q_n \in R_g$   $n$  distinct points ( $n$  can be zero.)

### Theorem

$\text{Aut}(R_g, q_1, \dots, q_n)$  is dense in  $\text{Diff}(R_g, q_1, \dots, q_n)$  in the  $C^\infty$ -topology on  $R_g$ .

Rational models of non-orientable surfaces:  $(\chi(R_g) = 2 - g)$

$R_g \sim B_{p_1, \dots, p_g} S^2$ , the sphere blown-up at  $g$  points

Let  $q_1, \dots, q_n \in R_g$   $n$  distinct points ( $n$  can be zero.)

## Theorem

$\text{Aut}(R_g, q_1, \dots, q_n)$  is dense in  $\text{Diff}(R_g, q_1, \dots, q_n)$  in the  $C^\infty$ -topology on  $R_g$ .

Steps of the proof:

① Marked points

[Huisman, M. 2007:  $\text{Aut}(S^m)$  acts  $\infty$ -transitively on  $S^m$ ,  $\forall m > 1$ ]  
 $\Rightarrow \text{Aut}(S^2, p_1, \dots, p_{g+n})$  is dense in  $\text{Diff}(S^2, p_1, \dots, p_{g+n})$  for any finite set of distinct points  $p_1, \dots, p_{g+n} \in S^2$ .

② Identity components


[Fragmentation Lemma]

$\Rightarrow \text{Aut}_0(R_g, q_1, \dots, q_n)$  is dense in  $\text{Diff}_0(R_g, q_1, \dots, q_n)$ .

③ Mapping class group

$\text{Aut}(R_g, q_1, \dots, q_n)$  surjects to  $\mathcal{M}(R_g, q_1, \dots, q_n)$ .

# Cremona transformation with real base points



Factored as:

$$S^2 \longleftarrow B_{p_1, \dots, p_4} S^2 \cong B_{q_1, \dots, q_4} S^2 \longrightarrow S^2$$

## Proposition

*Cremona transformations act transitively on isotopy classes of  $g$  disjoint Möbius bands in  $R_g$ .*

Cremona  $\sigma: B_{p_1, \dots, p_4} S^2 \cong B_{q_1, \dots, q_4} S^2$ ,  $\exists \Phi \in \text{Aut}(S^2)$  such that  $\Phi(p_i) = q_i$ ,  
get  $\Phi \circ \sigma$ :

$$B_{p_1, \dots, p_4} S^2 \xrightarrow{\sigma} B_{q_1, \dots, q_4} S^2 \xrightarrow{\Phi} B_{p_1, \dots, p_4} S^2$$


# The mapping class group

$R$  smooth compact surface

$$\mathcal{M}(R, q_1, \dots, q_n) := \pi_0(\text{Diff}(R, q_1, \dots, q_n))$$

## Theorem (Dehn 1938)

*When  $R$  orientable,  $\mathcal{M}$  is generated by Dehn twists around simple closed curves:*



## Theorem

*When  $R$  non-orientable, Dehn twists generate an index 2 subgroup of  $\mathcal{M}$ , need to add cross-cap slides.*

# Reduction of the set of generators

Chillingworth (1969), and Korkmaz (2002) with base points

Recall  $R_g = B_{p_1, \dots, p_g} S^2$

## Theorem


*Dehn twists around lifts of simple closed curves of  $S^2$  passing through an even number of the  $p_i$  (no self-intersection at the  $p_i$ ) suffice.*

With lantern relation  $\Rightarrow$



## Corollary

*Dehn twists around lifts of simple closed curves of  $S^2$  passing through 0, 2 or 4 of the  $p_i$  suffice.*

# Two models of the annulus blown up in 4 points





# The 4 exceptional curves and Dehn twist around $C_R$






# Deformation



## Image of the four exceptional curves



Cremona with 4 real base points represents the Dehn twist around  $C_R$  passing through the 4 base points.

## Generalizations: geometrically rational surfaces

$S$  := degree 2 Del Pezzo surface with  $\rho(S) = 1$

$C \subset S$  a curve  $\Rightarrow C \sim -aK_S$  for some  $a \in \mathbb{N}$

So  $p_a(C) = (C(C + K_S) - 2) / 2 = a(a - 1) - 1$  is odd

$C$  real rational  $\Rightarrow$  odd number of singular points on  $S(\mathbb{C})$ .

can not all be complex conjugate  $\Rightarrow$  no smooth rational curves on  $S$  at all.

### Conjecture

*Let  $S$  be a geometrically rational surface that is not isomorphic to a degree 2 Del Pezzo surface with Picard number 1, then every embedded circle can be approximated by smooth rational curves.*

## Generalizations: rationally connected varieties

We believe that usually not every homotopy class of  $X(\mathbb{R})$  can be represented by rational curves.

Let  $q_1, q_2, q_3$  be quadrics such that  $C := (q_1 = q_2 = q_3 = 0) \subset \mathbb{P}^4$  is a smooth curve with  $C(\mathbb{R}) \neq \emptyset$ . Consider the family of 3-folds

$$X_t := (q_1^2 + q_2^2 + q_3^2 - t(x_0^4 + \cdots + x_4^4) = 0) \subset \mathbb{P}^4$$

For  $0 < t \ll 1$ , the real points  $X_t(\mathbb{R})$  form an  $\mathbb{S}^2$ -bundle over  $C(\mathbb{R})$ .

### Conjecture

*For  $0 < t \ll 1$ , every rational curve  $g: \mathbb{P}^1 \rightarrow X_t$  gives a contractible map  $g: \mathbb{RP}^1 \rightarrow X_t(\mathbb{R})$ .*

### Conjecture

*Let  $X$  be a smooth, rationally connected variety defined over  $\mathbb{R}$ . Then a  $C^\infty$  map  $\mathbb{S}^1 \rightarrow X(\mathbb{R})$  can be approximated by rational curves iff it is homotopic to a rational curve  $\mathbb{RP}^1 \rightarrow X(\mathbb{R})$ .*

Merci

Спасибо

Thank you