



Math-Net.Ru

All Russian mathematical portal

D. A. Bredikhin, On semigroups of relations with the operation of left and right rectangular products, *Izv. Saratov Univ. Math. Mech. Inform.*, 2020, Volume 20, Issue 3, 280–289

DOI: 10.18500/1816-9791-2020-20-3-280-289

Use of the all-Russian mathematical portal Math-Net.Ru implies that you have read and agreed to these terms of use

<http://www.mathnet.ru/eng/agreement>

Download details:

IP: 18.97.9.170

February 17, 2025, 11:04:44





МАТЕМАТИКА

On Semigroups of Relations with the Operation of Left and Right Rectangular Products

D. A. Bredikhin

Dmitry A. Bredikhin, <https://orcid.org/0000-0003-3600-1294>, Yuri Gagarin State Technical University of Saratov, 77 Politechnicheskaya St., Saratov 410054, Russia, bredikhin@mail.ru

A set of binary relations closed with respect to some collection of operations on relations forms an algebra called an algebra of relations. The class of all algebras (partially ordered algebras) isomorphic to algebras (partially ordered by set-theoretic inclusion \subseteq algebras) of relations with operations from Ω is denoted by $R\{\Omega\}$ ($R\{\Omega, \subseteq\}$). An operation on relations is called primitive-positive if it can be defined by a formula of the first-order predicate calculus containing only existential quantifiers and conjunctions in its prenex normal form. We consider algebras of relations with associative primitive-positive operations $*$ and \star , defined by the following formulas $\rho * \sigma = \{(u, v) : (\exists s, t, w)(u, s) \in \rho \wedge (t, w) \in \sigma\}$ and $\rho \star \sigma = \{(u, v) : (\exists s, t, w)(s, t) \in \rho \wedge (w, v) \in \sigma\}$ respectively. The axiom systems for the classes $R\{*\}$, $R\{*, \subseteq\}$, $R\{\star\}$, $R\{\star, \subseteq\}$, and bases of quasi-identities and identities for quasi-varieties and varieties generated by these classes are found.

Keywords: algebra of relations, primitive positive operation, identity, variety, quasi-identity, quasi-variety, semigroup, partially ordered semigroup.

Received: 11.06.2019 / Accepted: 28.06.2019 /


Published: 31.08.2020

This is an open access article distributed under the terms of Creative Commons Attribution License (CC-BY 4.0)


DOI: <https://doi.org/10.18500/1816-9791-2020-20-3-280-289>

INTRODUCTION

Let $\text{Rel}(U)$ be the set of all binary relations on a base set U . A set of binary relations $\Phi \subseteq \text{Rel}(U)$ closed with respect to some collection Ω of operations on relations forms an algebra (Φ, Ω) called an *algebra of relations*. Theory of algebras of relations is an essential part of modern algebraic logic and has numerous applications in theory of semigroups [1].



НАУЧНЫЙ
ОТДЕЛ





Operations on relations are usually determined using first-order predicate calculus formulas. Such operations are called *logical*. It is known that classes of algebras of relations with logical operations are axiomatizable [1]. Logical operations can be classified by the type of formulas defining them. An operation on relations is called *primitive-positive* [2] (in other terminology — Diophantine operations [3,4], generalized superpositions [5]) if it can be defined by a formula of the first-order predicate calculus containing only existential quantifiers and conjunctions in its prenex normal form.

Note that the set-theoretical inclusion \subseteq is compatible with all primitive-positive operations. Thus, any algebra of relations with primitive-positive operations (Φ, Ω) can be considered as partially ordered $(\Phi, \Omega, \subseteq)$. We denote by $R\{\Omega\}$ (respectively, $R\{\Omega, \subseteq\}$) the class of all algebras (partially ordered algebras) isomorphic to the ones whose elements are binary relations and whose operations are members of Ω . Let $V\{\Omega\}$ ($V\{\Omega, \subseteq\}$) be the variety and let $Q\{\Omega\}$ ($Q\{\Omega, \subseteq\}$) be the quasi-variety generated by $R\{\Omega\}$ (generated by $R\{\Omega, \subseteq\}$ in the class of all partially ordered algebras of the corresponding type).

The following problems naturally arise when the class $R\{\Omega\}$ ($R\{\Omega, \subseteq\}$) is considered.

1. Find a system of axioms for the class $R\{\Omega\}$ ($R\{\Omega, \subseteq\}$).
2. Find a basis of quasi-identities for the quasi-variety $Q\{\Omega\}$ ($Q\{\Omega, \subseteq\}$).
3. Find a basis of identities for the variety $V\{\Omega\}$ ($V\{\Omega, \subseteq\}$).
4. Does the class $R\{\Omega\}$ ($R\{\Omega, \subseteq\}$) form a quasi-variety?
5. Does the quasi-variety $Q\{\Omega\}$ ($Q\{\Omega, \subseteq\}$) form a variety?

Numerous studies have been devoted to solving these problems for various classes of algebras of relations. The first mathematician who treated algebras of relations from the point of view of universal algebra was A. Tarski [6]. He considered algebras of relations (Tarski's algebras of relations) with the following operations: Boolean operations $\cup, \cap, -$; operations of relational product \circ and relational inverse $^{-1}$; constant operations Δ (diagonal relation), \emptyset (empty relation), $\nabla = U \times U$ (universal relation). He showed that the class $R\{\circ, ^{-1}, \cup, \cap, -, \Delta, \emptyset, \nabla\}$ is not a quasi-variety and the quasi-variety generated by this class forms a variety [7]. R. Lyndon [8] found the infinite base of identities for this variety.

In this paper, we will consider algebras of relations with one associative primitive-positive operation, i. e., semigroup of relations. Note that the operation \circ, \cap of Tarski's algebras of relations are primitive-positive. These operations are determined by the formulas:

$$\rho \circ \sigma = \{(u, v) : (\exists s)(u, s) \in \rho \wedge (s, v) \in \sigma\}, \quad \rho \cap \sigma = \{(u, v) : (u, v) \in \rho \wedge (u, v) \in \sigma\}.$$

It is well known that the class $R\{\circ\}$ coincides with the class of all semigroups and the class $R\{\cap\}$ coincides with the class of all semilattices. The class $R\{\circ, \subseteq\}$ coincides with the class of all partially ordered semigroups.

The associative operations $\triangleright, \triangleleft$ of restrictive multiplication were introduced and investigated in [9]. These operations are defined as follows:

$$\rho \triangleright \sigma = \{(u, v) : (\exists s)(u, s) \in \rho \wedge (u, v) \in \sigma\}, \quad \rho \triangleleft \sigma = \{(u, v) : (\exists s)(u, v) \in \rho \wedge (s, v) \in \sigma\}.$$

The classes $R\{\triangleright\}$ and $R\{\triangleleft\}$ coincide with the following varieties of semigroups given by the identities $x^2 = x$, $xyz = yxz$ and $x^2 = x$, $xyz = xzy$ respectively. Characterization of the corresponding partially ordered semigroups requires the additional identities $xy \leq y$ and $xy \leq x$ respectively.



There are other associative primitive-positive operations on relations [5]. It is interesting to consider problems 1–5 for classes of semigroups of relations with these operations. We concentrate our attention on the following associative binary primitive-positive operations $*$ and \star on $\text{Rel}(U)$ that is defined in the following way:

$$\begin{aligned} \rho * \sigma &= \{(u, v) : (\exists s, t, w)(u, s) \in \rho \wedge (t, w) \in \sigma\}, \\ \rho \star \sigma &= \{(u, v) : (\exists s, t, w)(s, t) \in \rho \wedge (w, v) \in \sigma\}. \end{aligned}$$

Since relations $\rho * \sigma$ and $\rho \star \sigma$ are rectangular, we will treat these operations as the operations of left and right *rectangular products* respectively. Note that the mapping $f(\rho) = \rho^{-1}$ is an antiisomorphism of the partially ordered semigroups of relations $(\text{Rel}(U), *, \subseteq)$ and $(\text{Rel}(U), \star, \subseteq)$. Therefore, it will be sufficient to study only one of these operations.

1. MAIN RESULTS

A *partially ordered semigroup* is an algebraic system (A, \cdot, \leq) , where (A, \cdot) is a semigroup and \leq is a partial order relation on A that is compatible with multiplication, i. e., $x \leq y$ implies $xz \leq yz$ and $zx \leq zy$ for all $x, y, z \in A$.

The main results are formulated in the following theorems and corollaries. These results provide a solution to problems 1–5 for classes $\text{R}\{*\}$ and $\text{R}\{*, \subseteq\}$. Their proofs are based on the description of quasi-equational theories of algebras of relations with primitive-positive operations [3].

Theorem 1. *The quasi-variety $\text{Q}\{*, \subseteq\}$ forms a variety in the class of all partially ordered semigroups. A partially ordered semigroup (A, \cdot, \leq) belongs to the quasi-variety $\text{Q}\{*, \subseteq\}$ if and only if it satisfies the identities:*

$$x^2y = xy, \tag{1}$$

$$xy^2 = xy, \tag{2}$$

$$xyz = xzy, \tag{3}$$

$$x \leq x^2, \tag{4}$$

$$xy \leq x^2. \tag{5}$$

Corollary 1. *The quasi-variety $\text{Q}\{*\}$ forms a variety. A semigroup (A, \cdot) belongs to the quasi-variety $\text{Q}\{*\}$ if and only if it satisfies the identities (1)–(3).*

Theorem 2. *The class $\text{R}\{*, \subseteq\}$ does not form a quasi-variety. For a partially ordered semigroup (A, \cdot, \leq) the following three conditions are equivalent.*

1. (A, \cdot, \leq) belongs to the class $\text{R}\{*, \subseteq\}$.

2. One of the following conditions holds:

a) (A, \cdot, \leq) satisfies identity (4) and the identity

$$xy = x^2; \tag{6}$$

b) (A, \cdot, \leq) contains the zero element o , satisfies identity (4) and the axioms:

$$y \neq o \Rightarrow xy = x^2, \tag{7}$$

$$o \leq x. \tag{8}$$



3. (A, \cdot, \leq) satisfies identity (4) and the axioms:

$$xy = x^2 \vee yz = zy = y, \tag{9}$$

$$xy = yx = x \Rightarrow x \leq z. \tag{10}$$

Corollary 2. *The class $R\{*\}$ does not form a quasi-variety. For a semigroup (A, \cdot) the following three conditions are equivalent.*

1. (A, \cdot) belongs to the class $R\{*\}$.
2. One of the following conditions holds:
 - a) (A, \cdot) satisfies identity (6);
 - b) (A, \cdot) contains the zero element and satisfies axiom (7).
3. (A, \cdot) satisfies axiom (9).

2. PROOFS OF RESULTS

Step 1. First of all, let us prove the necessity of the conditions of Theorems and Corollaries. Let $(\Phi, *, \subseteq)$ be the partially ordered semigroup of relations on U with the operation of left rectangular product and let $pr_1\rho = \{x : (\exists y)(x, y) \in \rho\}$ be the first projection of the relation $\rho \in \Phi$. It is clear that $\rho * \sigma = pr_1\rho \times U = \rho * \rho$ if $\sigma \neq \emptyset$, and $\rho * \sigma = \emptyset$ otherwise. It easily follows that that the identities (1)–(5) hold. It also follows that if $\emptyset \notin \Phi$, then $(\Phi, *)$ satisfies identity (6), and if $\emptyset \in \Phi$, then \emptyset is a zero element and axioms (7) and (8) hold.

Conditions 2 and 3 of Theorem 2 and Corollary 2 are equivalent. Indeed, axiom (6) can be represented as $(\neg(\forall w) yw = wy = y) \Rightarrow xy = x^2$. It follows that this axiom is equivalent to identity (6) if A does not contain a zero element, and it is equivalent to axiom (7) otherwise. Axiom (10) just expresses axiom (8) as a universal formula of the first-order language.

Step 2. The proof of sufficiency of conditions of Theorem 1 is based on the result of [3]. Let us give some definitions and notations to formulate this result.

For any formula $\varphi(z_0, z_1, r_1, \dots, r_m)$ of the first-order predicate calculus with the set of free variables included in $\{z_0, z_1\}$ and having m binary predicate symbols r_1, \dots, r_m , we can associate an m -ary operation F_φ on $\text{Rel}(U)$ defined in the following way:

$$F_\varphi(\rho_1, \dots, \rho_m) = \{(u, v) \in U \times U : \varphi(u, v, \rho_1, \dots, \rho_m)\},$$

where $\varphi(u, v, \rho_1, \dots, \rho_m)$ means that the formula φ holds whenever z_0, z_1 are interpreted as u, v , and r_1, \dots, r_m are interpreted as relations ρ_1, \dots, ρ_m from $\text{Rel}(U)$. As we stated earlier an operation on relations is called primitive-positive if it can be defined by a first-order formula containing only existential quantifiers and conjunctions in its prenex normal form. Let us describe primitive-positive operations by using graphs [2].

Let \mathbb{N} be the set of all natural numbers and $\mathbb{N}^0 = \mathbb{N} \cup \{0\}$. A *labeled graph* is a pair $G = (V(G), E(G))$, where $V(G)$ is a finite set (called a vertex set) and $E(G) \subseteq V(G) \times \mathbb{N} \times V(G)$ is a ternary relation. A triple $(u, k, v) \in E(G)$ is called an edge from u to v labeled by k , and it will be graphically represented by $u \cdot \xrightarrow{k} \cdot v$. An *input-output-pointed labeled graph* is a structure $G = (V(G), E(G), \text{in}(G), \text{out}(G))$, where $(V(G), E(G))$ is a labeled graph, $\text{in}(G)$ and $\text{out}(G)$ are two distinguished vertices called input and output vertices respectively. The input-output-pointed labeled graph G with $\text{in}(G) = i$ and $\text{out}(G) = o$ is also denoted by $G^{i,o}$. So, we will just call them graphs as it does not lead to confusion. The concept of graph isomorphism is defined



in a natural way. All graphs will be considered up to isomorphism. Moreover, we will identify graphs differing only in the number of isolated vertices that are distinct from *in* and *out*. These isolated vertices will be omitted.

For a given $u \in V(G)$, the number of edges of the form (u, k, v) [respectively, (v, k, u)] we denote by $deg^+(u)$ [respectively, $deg^-(u)$]. Given two input-output-pointed labeled graphs $G_1 = (V_1, E_1, in_1, out_1)$ and $G_2 = (V_2, E_2, in_2, out_2)$, a mapping $f: V_2 \rightarrow V_1$ is called a homomorphism from G_2 to G_1 if $f(in_2) = in_1$, $f(out_2) = out_1$, and $(f(u), k, f(v)) \in E_1$ whenever $(u, k, v) \in E_2$. We write $G_1 \prec G_2$ if there exists a homomorphism from G_2 to G_1 .

Let $F = F_\varphi$ be a primitive positive operation determined by a formula φ . Then the input-output-pointed labeled graph $G = G_F = G_\varphi$ associated with F is defined as follows (see [2]). Let $\{0, 1, \dots, n\}$ be the set of all subscripts of individual variables of φ . We put $G = (V, E, in, out)$, where $V = \{v_0, v_1, \dots, v_n\}$; $in = v_0$, $out = v_1$; $(i, k, j) \in E$ if and only if the atomic formula $r_k(z_i, z_j)$ occurs in φ .

Note that the graph $G_* = (V, E, in, out)$ corresponding to the considered operation $*$ of rectangular product can be described in the following way:

$$V = \{v_0, v_1, v_2, v_3, v_4\}, \quad E = \{(v_0, 1, v_3), (v_2, 2, v_4)\}, \quad in = v_0, \quad out = v_1,$$

$$in = v_0 \cdot \xrightarrow{1} \cdot v_3 \quad v_2 \cdot \xrightarrow{2} \cdot v_4 \quad \cdot v_1 = out.$$

Let $G = (V, E, in, out)$ and $G_k = (V_k, E_k, in_k, out_k)$ ($k = 1, 2, \dots, n$) be graphs with pairwise disjoint vertex sets. The composition $G(G_1, G_2, \dots, G_n)$ is the graph constructed as follows [3]: take G and substitute every edge $(u, k, v) \in E$ by the graph G_k identifying the input vertex in_k with u and the output vertex out_k with v .

Let us consider the set $\Lambda = \{x_1, \dots, x_n, \dots\}$ of individual variables that are interpreted as elements of a semigroup. A semigroup term p is a word in the alphabet Λ , i.e., an expression of the form $x_{j_1}x_{j_2} \dots x_{j_{m-1}}x_{j_m}$. For convenience, we will also use other letters of the Latin alphabet as variables.

Suppose that $p = x_{j_1}x_{j_2} \dots x_{j_{m-1}}x_{j_m}$ be the term of a semigroup that satisfies identities (1)–(3). Then without loss of generality, we can assume that all variables x_{j_1}, \dots, x_{j_m} are different. Moreover, we can also presume that variables x_{j_2}, \dots, x_{j_m} can be arbitrarily permuted. In what follows, we will use these properties without special mentions.

For any semigroup term p we define the graph $G(p) = (V(p), E(p), in(p), out(p))$ in the following inductive way:

- 1) if $p = x_k$, then $G(p)$ is the following graph: $in \cdot \xrightarrow{k} \cdot out$;
- 2) if $p = p_1p_2$, then $G(p) = G_*(G(p_1), G(p_2))$.

For any term $p = x_{j_1}x_{j_2} \dots x_{j_{m-1}}x_{j_m}$ the graph $G(p)$ has the following form:

$$in = v_0 \cdot \xrightarrow{j_1} \cdot \quad \cdot \xrightarrow{j_2} \cdot \quad \dots \quad \cdot \xrightarrow{j_{m-1}} \cdot \quad \cdot \xrightarrow{j_m} \cdot \quad \cdot v_1 = out.$$

Let G be a labeled graph, $u, v \in V(G) = \{v_0, v_1, \dots, v_n\}$, and Q be an input-output-pointed labeled graph. Without loss of generality, we can suppose that $V(Q) = \{w_0, w_1, \dots, w_m\}$, $in(Q) = w_0 = u$, $out(Q) = w_1 = v$, and $V(G) \cap V(Q) = \{u, v\}$. The labeled graph $(V(G) \cup V(Q), E(G) \cup E(Q))$ we denote by $G[u, v, Q]$. Note that the edges set of $G[u, v, Q]$ can be represented as $\{v_0, v_1, \dots, v_n, v_{n+1}, \dots, v_{n+m-1}\}$, where $v_{n+1} = w_2, \dots, v_{n+m-1} = w_m$. Virtually, the graph $G[u, v, Q]$ is obtained from the graph G by “gluing” the graph Q to the vertices u and v .



Define an n -system to be a pair $\omega = (\alpha, \beta)$, where $\alpha, \beta : \{1, \dots, n\} \rightarrow \mathbb{N}^0$ are mappings, $\alpha(k), \beta(k) < 2 + (k - 1)(m - 2)$ for all $k = 1, \dots, n$, and m is the number of vertices of the graph that determines the considered operation on relations (for the operation $*$ we have $m = 5$ and $\alpha(k), \beta(k) < 3k - 1$).

Given an n -system $\omega = (\alpha, \beta)$, we construct by induction the sequence of graphs $G_0 \subseteq \dots \subseteq G_n = G_\omega$. We put $G_0: v_0 \xrightarrow{1} \cdot v_1$, and for $k = 1, \dots, n$ we put:

$$G_k = G_{k-1}[v_{\alpha(k)}, v_{\beta(k)}, G(x_{2k}x_{2k+1})].$$

The following proposition presents the result of [3] formulated for the class $R\{*, \subseteq\}$.

Proposition 1. *A partially ordered semigroup (A, \cdot, \leq) belongs to the quasi-variety $Q\{*, \subseteq\}$ if and only if it satisfies the quasi-identity*

$$\left(\bigwedge_{k=1}^n p_k \leq x_{2k}x_{2k+1} \right) \Rightarrow x_1 \leq p_0 \tag{11}$$

for every n -system $\omega = (\alpha, \beta)$ and arbitrary terms p_0, \dots, p_n such that $G_\omega^{v_0, v_1} \prec G(p_0)$ and $G_{k-1}^{v_{\alpha(k)}, v_{\beta(k)}} \prec G(p_k)$.

Step 3. We are ready to prove the sufficiency of the conditions of Theorem 1. Let $\omega = (\alpha, \beta)$ be the n -system and p_0, p_1, \dots, p_n be the terms such that $G_\omega^{v_0, v_1} \prec G(p_0)$ and $G_{k-1}^{v_{\alpha(k)}, v_{\beta(k)}} \prec G(p_k)$ for $k = 1, \dots, n$. This system corresponds to the sequence of graphs $G_0 \subseteq G_1 \subseteq \dots \subseteq G_n = G_\omega$, where $G_k = (V_k, E_k)$ for $k = 0, \dots, n$. For any $k \leq n$ we have that $V_k = \{v_0, v_1, \dots, v_{3k-1}, v_{3k}, v_{3k+1}\}$ and

$$E_k = \{(v_0, 1, v_1)\} \cup \{(v_{\alpha(i)}, 2i, v_{3i-1}), (v_{3i}, 2i + 1, v_{3i+1}) : i = 1, \dots, k\}.$$

Let us prove by induction on k that $\alpha(k) = 3j$ for some $j = 0, \dots, k$; $\text{deg}^- v_{3i} = 0$, $\text{deg}^+ v_{3i} > 0$ for any $i = 0, \dots, k$, and $\text{deg}^- v = 1$, $\text{deg}^+ v = 0$ for all other vertices v of the graph G_k .

Let $k = 1$. Since $G_0^{v_{\alpha(1)}, v_{\beta(1)}} \prec G(p_1)$, we have $p_1 = x_1$ or $p_1 = x_1^2$, and $\alpha(1) = 0$, $\beta(1) = 1$; $\text{deg}^- v_0 = 0$, $\text{deg}^+ v_0 = 2$; $\text{deg}^- v_1 = 1$, $\text{deg}^+ v_1 = 0$; $\text{deg}^- v_2 = 1$, $\text{deg}^+ v_2 = 0$; $\text{deg}^- v_3 = 0$, $\text{deg}^+ v_3 = 1$; $\text{deg}^- v_4 = 1$, $\text{deg}^+ v_4 = 0$.

Suppose now that it holds for $k - 1$, and let us show that this is true for k . Since $G_{k-1}^{v_{\alpha(k)}, v_{\beta(k)}} \prec G(p_k)$, according to the definition of a graph homomorphism, we get $\text{deg}^+ v_{\alpha(k)} > 0$. Then according to the induction assumption we get $\alpha(k) = 3j$ for some $j = 0, \dots, k$; $\text{deg}^- v_{3i} = 0$, $\text{deg}^+ v_{3i} > 0$ for any $i = 0, \dots, k$, and $\text{deg}^- v = 1$, $\text{deg}^+ v = 0$ for all other vertices v of the graph G_k .

Let (A, \cdot, \leq) be a partially ordered semigroup satisfying identities (1)–(5). Suppose that the premise of quasi-identity (11) holds for some values of the variables $x_1 = a_1$, $x_2 = a_2$, $x_3 = a_3, \dots, x_{2n} = a_{2n}, x_{2n+1} = a_{2n+1}$, i. e., $p_k(\vec{a}) \leq a_{2k}a_{2k+1}$ for all $k = 1, \dots, n$, where $\vec{a} = (a_1, a_2, \dots, a_{2n+1})$. Let $p_0 = x_{j_1}x_{j_2} \dots x_{j_{m-1}}x_{j_m}$. Note that $G_k^{v_0, v_1} \prec G(p_0)$ if and only if $\{x_{j_1}, x_{j_2}, \dots, x_{j_{m-1}}, x_{j_m}\} \subseteq \{x_1, x_2, \dots, x_{2k}, x_{2k+1}\}$, $x_{j_1} = x_1$ or $x_{j_1} = x_{3i}$ for some $0 < i \leq k$ such that $\alpha(i) = 0$.

Let $\max(p_0)$ be the greatest k such that at least one of the variables x_{2k} or x_{2k+1} is included in the term p_0 . Let us prove by induction on $\max(p_0)$ that $a_1 \leq p_0(\vec{a})$. If $\max(p_0) = 0$, then $p = x_1$ or $p_0 = x_1^2$. Thus, using identity (4) we obtain $a_1 \leq p_0(\vec{a})$.



Suppose now that $a_1 \leq p_0(\vec{a})$ holds for $\max(p_0) = k - 1$, and let us show that this is true for $\max(p_0) = k$.

If both variables x_{2k}, x_{2k+1} are included in p_0 , then the following cases are possible:

1) $p_0 = x_{2k}x_{2k+1}x_{j_3} \dots x_{j_{m-1}}x_{j_m}$, then $\alpha(k) = 0$. It follows that $G_k^{v_0, v_1} \prec G(p_k x_{j_3} \dots x_{j_m})$, and by the induction assumption we get

$$a_0 \leq p_k(\vec{a})a_{j_3} \dots a_{j_{m-1}}a_{j_m} \leq a_{2k}a_{2k+1}a_{j_3} \dots a_{j_{m-1}}a_{j_m} = p_0(\vec{a});$$

2) $p_0 = x_{j_1}x_{2k}x_{2k+1}x_{j_4} \dots x_{j_{m-1}}x_{j_m}$, then using the induction assumption we get

$$a_0 \leq a_{j_1}p_k(\vec{a})a_{j_4} \dots a_{j_{m-1}}a_{j_m} \leq a_{j_1}a_{2k}a_{2k+1}a_{j_4} \dots a_{j_{m-1}}a_{j_m} = p_0(\vec{a}).$$

If only one of the variables x_{2k} or x_{2k+1} is included in p_0 , then the following cases are possible:

3) $p_0 = x_{2k}x_{j_2} \dots x_{j_{m-1}}x_{j_m}$, then $\alpha(k) = 0$. It follows that $G_k^{v_0, v_1} \prec G(p_k x_{j_2} \dots x_{j_m})$, and by the induction assumption we get

$$\begin{aligned} a_0 &\leq p_k(\vec{a})a_{j_2} \dots a_{j_{m-1}}a_{j_m} \leq a_{2k}a_{2k+1}a_{j_3} \dots a_{j_{m-1}}a_{j_m} \leq \\ &\stackrel{(5)}{\leq} a_{2k}^2 a_{j_2} \dots a_{j_{m-1}}a_{j_m} \stackrel{(1)}{=} a_{2k}a_{j_2} \dots a_{j_{m-1}}a_{j_m} = p_0(\vec{a}); \end{aligned}$$

4) $p_0 = x_{j_1}x_{2k}x_{j_3} \dots x_{j_{m-1}}x_{j_m}$, then using the induction assumption we get

$$\begin{aligned} a_0 &\leq a_{j_1}p_k(\vec{a})a_{j_3} \dots a_{j_{m-1}}a_{j_m} \leq a_{j_1}a_{2k}a_{2k+1}a_{j_3} \dots a_{j_{m-1}}a_{j_m} \leq \\ &\stackrel{(5)}{\leq} a_{j_1}a_{2k}^2 a_{j_3} \dots a_{j_{m-1}}a_{j_m} \stackrel{(2)}{=} a_{j_1}a_{2k}a_{j_3} \dots a_{j_{m-1}}a_{j_m} = p_0(\vec{a}); \end{aligned}$$

5) $p_0 = x_{j_1}x_{2k+1}x_{j_3} \dots x_{j_{m-1}}x_{j_m}$, then using the induction assumption we get

$$\begin{aligned} a_0 &\leq a_{j_1}p_k(\vec{a})a_{j_4} \dots a_{j_{m-1}}a_{j_m} \leq a_{j_1}a_{2k}a_{2k+1}a_{j_4} \dots a_{j_{m-1}}a_{j_m} \leq \\ &\stackrel{(5)}{\leq} a_{j_1}a_{2k+1}^2 a_{j_4} \dots a_{j_{m-1}}a_{j_m} \stackrel{(2)}{=} a_{j_1}a_{2k+1}a_{j_4} \dots a_{j_{m-1}}a_{j_m} = p_0(\vec{a}). \end{aligned}$$

Thus, we have proved that the partially ordered semigroup (A, \cdot, \leq) satisfies quasi-identity (11). Therefore, according to Proposition 1 we have $(A, \cdot, \leq) \in \mathcal{Q}\{*, \subseteq\}$. This completes the proof of Theorem 1.

Step 4. Let us prove Corollary 1. Suppose that a semigroup (A, \cdot) satisfies identities (1)–(3) and $A^2 = \{a^2 : a \in A\}$. We define the relation \leq on the set A by setting

$$\leq = \{(x, y) \in A \times A^2 : x^2 = yx\} \cup \{(x, x) \in A \times A : x \in A\}.$$

Let us show that (A, \cdot, \leq) is the partially ordered semigroup satisfying identities (4) and (5). The reflexivity of the relation \leq follows directly from the definition.

To prove the transitivity assume that $x \leq y$ and $y \leq z$. Without loss of generality, we can suppose that $x \neq y$ and $y \neq z$. Then $x^2 = yx$, $y^2 = zy$ and $y^2 = y$, $z^2 = z$, hence $x^2 = yx = zy x = z x^2 \stackrel{(2)}{=} z x$, i. e., $x \leq z$. Thus, \leq is transitive.

Assume that $x \leq y$, $y \leq x$ and $x \neq y$. Then $x^2 = yx$, $y^2 = xy$ and $x^2 = x$, $y^2 = y$, hence $x = x^2 = yx \stackrel{(1)}{=} y^2 x = xyx \stackrel{(3)}{=} x^2 y \stackrel{(1)}{=} xy = y^2 = y$. This contradicts the assumption $x \neq y$. Thus, \leq is a partially order relation.



Let us show that the relation \leq is compatible with multiplication. Suppose that $x \leq y$ and $x \neq y$. Then $x^2 = yx$ and $y^2 = y$, hence $(xz)^2 = xzxz \stackrel{(3)}{=} x^2z^2 = yxz^2 \stackrel{(3)}{=} (yz)(xz)$ and $(yz)^2 = yzyz \stackrel{(3)}{=} y^2z^2 \stackrel{(1)}{=} yz^2 \stackrel{(2)}{=} yz$. Thus, $xz \leq yz$. Further, $(zx)^2 = zxzx \stackrel{(3)}{=} z^2x^2 = z^2yx \stackrel{(3)}{=} (zy)(zx)$ and $(zy)^2 = zyzzy \stackrel{(3)}{=} z^2y^2 \stackrel{(1)}{=} zy^2 \stackrel{(2)}{=} zy$. Thus, $zx \leq zy$.

Since $x^2 \stackrel{(3)}{=} x^3 = x^2x$ and $(x^2)^2 = x^4 \stackrel{(1)}{=} x^2$, we have $x \leq x^2$. Since $(xy)^2 = xyxy \stackrel{(3)}{=} xxy^2 \stackrel{(1)}{=} x^2xy^2 \stackrel{(2)}{=} x^2xy$ and $(x^2)^2 = x^4 \stackrel{(1)}{=} x^2$, we have $xy \leq x^2$. Therefore, (A, \cdot, \leq) satisfies identities (1) and (2), hence $(A, \cdot, \leq) \in Q\{*, \subseteq\}$ and $(A, \cdot) \in Q\{*\}$. This completes the proof of Corollary 1.

Step 5. Let us prove Theorem 2 and Corollary 2. First of all, it is easy to see that for $U \neq \emptyset$, the Cartesian square of the semigroup $(\text{Rel}(U), *)$ of relations contains the zero element (\emptyset, \emptyset) and does not satisfy axiom (7). It follows that the classes $R\{*\}$ and $R\{*, \subseteq\}$ do not form quasi-varieties.

Lemma 1. *Let $\{U_j : j \in J\}$ be a family of pairwise non-intersecting sets and $U = \bigcup\{U_j : j \in J\}$. If a partially ordered semigroup (A, \cdot, \leq) is a subdirect product of a family $\{(\Phi_j, *, \subseteq) : j \in J\}$ of partially ordered semigroups of relations on U_j and satisfies identity (6), then (A, \cdot, \leq) is isomorphically embedded in $(\text{Rel}(U), *, \subseteq)$.*

Proof. Let $\varphi_j : A \rightarrow \Phi_j$ be the corresponding surjective homomorphisms from A on the components of the direct product $\prod\{\Phi_j : j \in J\}$. According to the properties of homomorphic images, we have that all components $(\Phi_j, *, \subseteq)$ satisfy identity (6). Hence, for all $j \in J$ we have $\emptyset \notin \Phi_j$ or $\Phi_j = \{\emptyset\}$. It follows that (A, \cdot, \leq) is a subdirect product of the family $\{(\Phi_j, *, \subseteq) : j \in J_0\}$, where $J_0 = \{j \in J : \emptyset \notin \Phi_j\}$.

For a given $a \in A$, we put $\rho_j^a = \varphi_j(a)$. We define a mapping $\varphi : A \rightarrow \text{Rel}(U)$ in the following way. We put $\varphi(a) = \bigcup\{pr_1\rho_j^a : j \in J_0\} \times U$, if $a^2 = a$, and $\varphi(a) = \bigcup\{\rho_j^a : j \in J_0\} \cup \bigcup\{\varphi(b) : b^2 = b \leq a\}$ otherwise. Let us show that φ is an isomorphic embedding (A, \cdot, \leq) in $(\text{Rel}(U), *, \subseteq)$.

Note that $\varphi(a) \cap U_j \times U_j = \rho_j^a$ for all $a \in A$. It follows that $\varphi(a) \subseteq \varphi(b)$ if and only if $a \leq b$. Furthermore, since $\emptyset \notin \Phi_j$ and $(ab)^2 = abab \stackrel{(3)}{=} aabb \stackrel{(2)}{=} aab \stackrel{(1)}{=} ab$, we have

$$\begin{aligned} \varphi(ab) &= \bigcup\{pr_1\rho_j^{ab} : j \in J_0\} \times U = \left(\bigcup\{pr_1\rho_j^a : j \in J_0\} \times U\right) * \left(\bigcup\{pr_1\rho_j^b : j \in J_0\} \times U\right) \\ &= pr_1\varphi(a) \times U * \varphi(b) = \varphi(a) * \varphi(b). \end{aligned} \quad \square$$

Lemma 2. *Suppose that (A, \cdot, \leq) satisfies identities (4) and (6). Then (A, \cdot, \leq) belongs to $R\{*, \subseteq\}$.*

Proof. It is easy to see that identity (6) implies identities (1)–(3) and (5). Hence, if (A, \cdot, \leq) satisfies identities (4) and (6), then according to Theorem 1 we have that $(A, \cdot, \leq) \in Q\{*, \subseteq\}$. In respect that the class $R\{*, \subseteq\}$ is axiomatizable [1], we obtain that (A, \cdot, \leq) is a subdirect product of a family of partially ordered semigroups from $R\{*, \subseteq\}$. Hence, according to Lemma 1, we obtain that (A, \cdot, \leq) belongs to $R\{*, \subseteq\}$. \square

Lemma 3. *Suppose that (A, \cdot) satisfies identity (6). Then (A, \cdot) belongs to $R\{*\}$.*

Proof. If (A, \cdot) satisfies identity (6), then it satisfies identities (1)–(3). Let \leq be the partial order relation constructed in the proof of Corollary 1. Then by Lemma 2 we have $(A, \cdot, \leq) \in R\{*, \subseteq\}$. Therefore, $(A, \cdot) \in R\{*\}$. \square



Lemma 4. Suppose that (A, \cdot) contains the zero element o and satisfies axiom (7). Then (A, \cdot) satisfies identity (6) or $ab \neq o$ for all $a, b \neq o$.

Proof. If there exist elements $a \neq o$ and $b \neq o$ such that $ab = o$, then for all $x, y \neq o$ we have $xa \stackrel{(7)}{=} x^2$, $yb \stackrel{(7)}{=} y^2$, and $xab = o$, hence $xy \stackrel{(1)}{=} x^2y \stackrel{(2)}{=} x^2y^2 = xayb \stackrel{(3)}{=} xyab = xyoy = o$. It follows that $xy = x^2$ for all $x, y, z \in A$, i. e., (A, \cdot) satisfies identity (6). \square

Suppose that (A, \cdot, \leq) contains the zero element o and satisfies identity (4) and axioms (7) and (8). We put $B = A \setminus \{o\}$. According to Lemmas 3 and 4 we can suppose that $xy \in B$ for all $x, y \in B$, and (B, \cdot, \leq) satisfies identities (4) and (6), hence (B, \cdot, \leq) belongs to $R\{*, \subseteq\}$. It means that there exists an isomorphism F from the partially ordered semigroup (B, \cdot, \leq) to some partially ordered semigroup of relations $(\Phi, *, \subseteq)$ and $\emptyset \notin \Phi$. Putting $F(o) = \emptyset$, we get the isomorphism from (A, \cdot, \leq) to $(\Phi \cup \{\emptyset\}, *, \subseteq)$. Therefore, (A, \cdot, \leq) belongs to $R\{*, \subseteq\}$. This completes the proof of Theorem 2.

Suppose that (A, \cdot) contains the zero element o and satisfies axiom (7), $B = A \setminus \{o\}$, and let \leq be the partial order relation on B constructed in the proof of Theorem 2. Extend the relation \leq on A by putting $o \leq a$ for all $a \in A$. Then (A, \cdot, \leq) satisfies the conditions of Theorem 2, hence $(A, \cdot, \leq) \in R\{*, \subseteq\}$. Therefore, (A, \cdot) belongs to $R\{*\}$. This completes the proof of Corollary 2.

CONCLUSION

The results of the article show that the classes $R\{*\}$, $R\{*, \subseteq\}$, $R\{\star\}$, $R\{\star, \subseteq\}$ are finitely axiomatizable and are not quasi-varieties; the quasivarieties generated by these classes are finitely based varieties, and also that $R\{\triangleleft\} \subset V\{*\}$, $R\{\triangleleft, \subseteq\} \subset V\{*, \subseteq\}$, $R\{\triangleright\} \subset V\{\star\}$, $R\{\triangleright, \subseteq\} \subset V\{\star, \subseteq\}$.

References

1. Schein B. M. Relation algebras and function semigroups. *Semigroup Forum*, 1970, vol. 1, pp. 1–62.
2. Böner P., Pöschel F. R. Clones of operations on binary relations. *Contributions to general algebra*, 1991, vol. 7, pp. 50–70.
3. Bredikhin D. A. On quasi-identities of algebras of relations with Diophantine operations. *Sib. Math. J.*, 1997, vol. 38, pp. 23–33. DOI: <https://doi.org/10.1007/BF02674896>
4. Bredikhin D. A. On algebras of relations with Diophantine operations. *Dokl. Math.*, 1998, vol. 57, no. 3, pp. 435–436.
5. Bredikhin D. A. On relation algebras with general superpositions. *Algebraic Logic*, 1998, vol. 54, pp. 111–124.
6. Tarski A. On the calculus of relations. *J. Symbolic Logic*, 1941, vol. 6, pp. 73–89.
7. Tarski A. Contributions to the theory of models, III. *Proc. Koninkl. Nederl. Akad. Wet.*, 1956, vol. 58, pp. 56–64.
8. Lyndon R. C. The representation of relation algebras, II. *Ann. Math.*, 1956, vol. 63, no. 2, pp. 294–307. DOI: <https://doi.org/10.2307/1969611>
9. Wagner V. V. Restrictiv semigroups. *Izv. Vyssh. Uchebn. Zaved. Mat.*, 1962, no. 6, pp. 19–27 (in Russian).

Cite this article as:

Bredikhin D. A. On Semigroups of Relations with the Operation of Left and Right Rectangular Products. *Izv. Saratov Univ. (N. S.), Ser. Math. Mech. Inform.*, 2020, vol. 20, iss. 3, pp. 280–289. DOI: <https://doi.org/10.18500/1816-9791-2020-20-3-280-289>



УДК 501.1

О полугруппах отношений с операцией левого и правого прямоугольного произведения

Д. А. Бредихин

Бредихин Дмитрий Александрович, доктор физико-математических наук, профессор кафедры математики и моделирования, Саратовский государственный технический университет имени Гагарина Ю. А., Россия, 410054, г. Саратов, ул. Политехническая, д. 77, bredikhin@mail.ru

Множество бинарных отношений, замкнутое относительно некоторой совокупности операций над ними, образует алгебру, называемую алгеброй отношений. Класс всех алгебр (частично упорядоченных алгебр), изоморфных алгебрам (частично упорядоченным теоретико-множественным включениям \subseteq алгебрам) отношений с операциями из Ω , обозначим $R\{\Omega\}$ ($R\{\Omega, \subseteq\}$). Операция над бинарными отношениями называется примитивно-позитивной, если она может быть определена формулой, содержащей в своей префиксной нормальной форме лишь кванторы существования и операцию конъюнкции. В работе рассматриваются алгебры отношений с ассоциативными примитивно-позитивными операциями $*$ и \star , определяемыми следующими формулами: $\rho * \sigma = \{(u, v) : (\exists s, t, w)(u, s) \in \rho \wedge (t, w) \in \sigma\}$ и $\rho \star \sigma = \{(u, v) : (\exists s, t, w)(s, t) \in \rho \wedge (w, v) \in \sigma\}$ соответственно. Найдены системы аксиом для классов $R\{*\}$, $R\{*, \subseteq\}$, $R\{\star\}$, $R\{\star, \subseteq\}$ и базисы тождеств для порожденных этими классами квазимногообразий и многообразий.

Ключевые слова: алгебра отношений, примитивно-позитивная операция, тождество, многообразие, квазитожество, квазимногообразие, полугруппа, частично упорядоченная полугруппа.

Поступила в редакцию: 11.06.2019 / Принята: 28.06.2019 / Опубликовано: 31.08.2020

Статья опубликована на условиях лицензии Creative Commons Attribution License (CC-BY 4.0)

Образец для цитирования:

Bredikhin D. A. On Semigroups of Relations with the Operation of Left and Right Rectangular Products [Бредихин Д. А. О полугруппах отношений с операцией левого и правого прямоугольного произведения] // Изв. Саратов. ун-та. Нов. сер. Сер. Математика. Механика. Информатика. 2020. Т. 20, вып. 3. С. 280–289. DOI: <https://doi.org/10.18500/1816-9791-2020-20-3-280-289>
