

Общероссийский математический портал

У. А. Ровба, К. А. Смотритский, Е. В. Дирвук, О константе Лебега интерполяционного рационального процесса с узлами Чебышева – Маркова,
Журн. Белорус. гос. ун-та. Матем. Инф., 2018, том 3, 12–20

<https://www.mathnet.ru/bgumi115>

Использование Общероссийского математического портала Math-Net.Ru подразумевает, что вы прочитали и согласны с пользовательским соглашением

<https://www.mathnet.ru/rus/agreement>

Параметры загрузки:

IP: 18.97.9.175

14 мая 2025 г., 01:38:12



УДК 513.5

О КОНСТАНТЕ ЛЕБЕГА ИНТЕРПОЛЯЦИОННОГО РАЦИОНАЛЬНОГО ПРОЦЕССА С УЗЛАМИ ЧЕБЫШЕВА – МАРКОВА

Е. А. РОВБА¹⁾, К. А. СМОТРИЦКИЙ¹⁾, Е. В. ДИРВУК¹⁾

¹⁾Гродненский государственный университет им. Янки Купалы,
ул. Ожешко, 22, 230023, г. Гродно, Беларусь

Рассматривается оценка константы Лебега интерполяционного рационального процесса Лагранжа на отрезке $[-1, 1]$ с узлами в нулях косинус-дробей Чебышева – Маркова. Показано, что в случае двух действительных геометрически различных полюсов аппроксимирующих функций нормы фундаментальных многочленов Лагранжа ограничены. На основании этого результата доказано, что в рассматриваемом случае оценка сверху константы Лебега не зависит от расположения полюсов и последовательность констант Лебега растет с логарифмической скоростью. В предыдущих работах оценки констант Лебега были получены только для конкретных наборов полюсов или зависели от расположения полюсов.

Ключевые слова: рациональные приближения; интерполирование; дробь Чебышева – Маркова; константа Лебега.

ON A LEBESGUE CONSTANT OF INTERPOLATION RATIONAL PROCESS AT THE CHEBYSHEV – MARKOV NODES

Y. A. ROUBA^a, K. A. SMATRYTSKI^a, Y. V. DIRVUK^a

^aYanka Kupala State University of Grodno, 22 Ažėška Street, Grodno 230023, Belarus
Corresponding author: K. A. Smatrytski (k_smotritski@mail.ru)

In the present paper estimate of a Lebesgue constant of the interpolation rational Lagrange process on the segment $[-1, 1]$ at the Chebyshev – Markov cosine fractions nodes is considered. It is shown that in the case of two real geometrically distinct poles of approximating functions, the norms of the Lagrange fundamental polynomials are bounded. Based on this result, it is proved that in the case under consideration the upper estimate of the Lebesgue constant does not depend on the arrangement of the poles and the sequence of the Lebesgue constant grows with logarithmic rate. Note, that in previous works the estimates of Lebesgue constants were obtained only for particular choices of poles or depended on the arrangement of poles.

Key words: rational approximation; interpolation; Chebyshev – Markov fraction; Lebesgue constant.

Образец цитирования:

Ровба ЕА, Смотрицкий КА, Дирвук ЕВ. О константе Лебега интерполяционного рационального процесса с узлами Чебышева – Маркова. *Журнал Белорусского государственного университета. Математика. Информатика.* 2018;3:12–20.

For citation:

Rouba YA, Smatrytski KA, Dirvuk YV. On a Lebesgue constant of interpolation rational process at the Chebyshev – Markov nodes. *Journal of the Belarusian State University. Mathematics and Informatics.* 2018;3:12–20. Russian.

Авторы:

Евгений Алексеевич Ровба – доктор физико-математических наук, профессор; заведующий кафедрой фундаментальной и прикладной математики факультета математики и информатики.

Константин Анатольевич Смотрицкий – кандидат физико-математических наук, доцент; доцент кафедры фундаментальной и прикладной математики факультета математики и информатики.

Евгений Владимирович Дирвук – кандидат физико-математических наук; доцент кафедры системного программирования и компьютерной безопасности факультета математики и информатики.

Authors:

Yauheni A. Rouba, doctor of science (physics and mathematics), full professor; head of the department of fundamental and applied mathematics, faculty of mathematics and informatics. rovba.ea@gmail.com

Kanstantin A. Smatrytski, PhD (physics and mathematics), doцент; associate professor at the department of fundamental and applied mathematics, faculty of mathematics and informatics. k_smotritski@mail.ru

Yauheni V. Dirvuk, PhD (physics and mathematics); associate professor at the department of system programming and computer security, faculty of mathematics and informatics. dirvuk@gmail.com

Introduction

Behaviour of the Lebesgue constant of interpolating polynomial processes was always of great interest of the researchers (see, for example, [1]). In this direction very deep and profound results were obtained [2].

In the case of interpolation by rational functions, this problem is less studied. Beginning of the investigation of the behavior of the Lebesgue constants of interpolation rational Lagrange processes at the Chebyshev – Markov nodes on a segment was made by V. N. Rusak [3], continued by A. P. Starovoitov [4]. However, in these papers estimates of the Lebesgue constants were obtained under certain conditions on the poles of interpolation rational functions. The main result of [5] is of a different nature. Let us quote it.

Let $\{a_k\}_{k=0}^n$ be a sequence of real numbers such that $a_0 = 0$, $a_k \in (-1, 1)$, $k = 1, 2, \dots, n$.

Chebyshev – Markov rational fraction can be defined as follows

$$M_n(x) = \cos \mu_n(x),$$

where

$$\mu_n(x) = \sum_{k=0}^n \arccos \frac{x + a_k}{1 + a_k x},$$

and

$$\mu'_n(x) = -\frac{\lambda_n(x)}{\sqrt{1-x^2}}, \quad \lambda_n(x) = \sum_{k=0}^n \frac{\sqrt{1-a_k^2}}{1+a_k x}, \quad x \in (-1, 1).$$

The function $M_n(x)$ has $n + 1$ zeroes x_k , $k = 0, 1, \dots, n$, on the interval $(-1, 1)$. Then for any function f defined on $[-1, 1]$ we construct the interpolating rational Lagrange function

$$L_n(x; f) = \sum_{k=0}^n f(x_k) l_k(x), \tag{1}$$

where

$$l_k(x) = \frac{M_n(x)}{(x - x_k) M'_n(x_k)}, \quad k = 0, 1, \dots, n.$$

Theorem 1. *For the Lebesgue function*

$$L_n(x) = \sum_{k=0}^n |l_k(x)|$$

of the interpolating Lagrange process (1) the following estimate holds

$$L_n(x) < 1 + Cd_n \ln \sum_{k=0}^n \left(\sqrt{\frac{1-a_k}{1+a_k}} + \sqrt{\frac{1+a_k}{1-a_k}} \right), \quad n = 1, 2, \dots, \tag{2}$$

where C is a positive constant,

$$d_n = \max_{k=1, 2, \dots, n} \left\{ \frac{\lambda_n(x_{k-1})}{\lambda_n(x_k)}, \frac{\lambda_n(x_k)}{\lambda_n(x_{k-1})} \right\}.$$

Note, that estimate (2) was obtained for any real poles, without any restrictions on their location.

Let us pay some attention to the right-hand side of the estimate (2). Like in previous studies it is a function of poles. However in [6] it was proved that in one particular case $n = 2$ there is no such dependence. In the present work we continue such study.

Basic definitions and auxiliary results

Let $a \in [0, 1)$. Chebyshev – Markov rational fraction with two geometrically distinct real poles can be written as follows:

$$M_{2n}(x) = \cos \mu_{2n}(x), \quad x \in [-1, 1], \tag{3}$$

$$\mu_{2n}(x) = n \left(\arccos \frac{x+a}{1+ax} + \arccos \frac{x-a}{1-ax} \right).$$

Remark 1. If

$$x = \frac{\cos \theta}{\sqrt{1 - a^2 \sin^2 \theta}}, \quad \theta \in [0, \pi],$$

then the rational fraction (1) can be written in the following form

$$M_{2n} \left(\frac{\cos \theta}{\sqrt{1 - a^2 \sin^2 \theta}} \right) = \cos 2n\theta, \quad \theta \in [0, \pi].$$

Using the definition, we obtain directly the following lemma.

Lemma 1. *The function $M_{2n}(x)$ has $2n$ simple zeroes, x_k , $k = 1, 2, \dots, 2n$ on the interval $(-1, 1)$, that are symmetric with respect to the imaginary axis:*

$$-1 < x_{2n} < x_{2n-1} < \dots < x_{n+1} < 0 < x_n < x_{n-1} < \dots < x_1 < 1;$$

$$x_k = \frac{\cos \theta_k}{\sqrt{1 - a^2 \sin^2 \theta_k}}, \quad \theta_k = \frac{2k-1}{4n} \pi, \quad k = 1, 2, \dots, 2n;$$

$$x_{n+k} = -x_{n-k+1}, \quad k = 1, 2, \dots, n.$$

For the further research we need the following property of the zeroes of Chebyshev – Markov rational fraction (3).

Lemma 2. *The following inequalities hold:*

$$\frac{1 - a^2 \sin^2 \theta_{k-1}}{1 - a^2 \sin^2 \theta_k} < 9, \quad a \in [0, 1), \quad \theta_k = \frac{2k-1}{4n} \pi, \quad k = 2, 3, \dots, n+1.$$

Proof. Let us consider the functions

$$\varphi_k(u) = \frac{1 - u \sin^2 \theta_{k-1}}{1 - u \sin^2 \theta_k}, \quad u \in [0, 1), \quad k = 2, 3, \dots, n.$$

Since

$$\varphi'_k(u) = \frac{\sin^2 \theta_k - \sin^2 \theta_{k-1}}{(1 - u \sin^2 \theta_k)^2} > 0,$$

then

$$\frac{1 - a^2 \sin^2 \theta_{k-1}}{1 - a^2 \sin^2 \theta_k} < \frac{1 - \sin^2 \theta_{k-1}}{1 - \sin^2 \theta_k} = \left(\frac{\cos \theta_{k-1}}{\cos \theta_k} \right)^2.$$

Now,

$$\left(\frac{\cos \left(\theta - \frac{\pi}{2n} \right)}{\cos \theta} \right)' = \frac{\sin \left(\frac{\pi}{2n} \right)}{\cos^2 \theta} > 0, \quad \theta \in \left[0, \frac{\pi}{2} \right),$$

and

$$\left(\frac{\cos \theta_{k-1}}{\cos \theta_k} \right)^2 < \left(\frac{\cos \theta_{n-1}}{\cos \theta_n} \right)^2 = \left(\frac{\sin \left(\frac{3\pi}{4n} \right)}{\sin \left(\frac{\pi}{4n} \right)} \right)^2 < 9.$$

We have $\theta_{n+1} + \theta_n = \pi$ for $k = n + 1$. It means

$$\frac{1 - a^2 \sin^2 \theta_n}{1 - a^2 \sin^2 \theta_{n+1}} = 1.$$

Lemma 2 is proved.

Now for any function $f(x)$, defined on the segment $[-1, 1]$, we construct the interpolating rational Lagrange function

$$L_{2n}(x; f) = \sum_{k=1}^{2n} f(x_k) l_k(x), \quad (4)$$

where

$$l_k(x) = \frac{M_{2n}(x)}{(x - x_k) M'_{2n}(x_k)}, \quad k = 1, 2, \dots, 2n. \quad (5)$$

It turns out that in this case the uniform norms of the Lagrange's fundamental polynomials are bounded.

Lemma 3. *The following inequalities hold:*

$$\max_{x \in [-1, 1]} |l_k(x)| < 6, \quad k = 1, 2, \dots, n + 1.$$

Proof. Since

$$M'_{2n}(x) = \sin \mu_{2n}(x) \frac{2n\sqrt{1-a^2}}{(1-a^2x^2)\sqrt{1-x^2}},$$

then from (5) it follows that

$$|l_k(x)| = \frac{|M_{2n}(x)|(1-a^2x_k^2)\sqrt{1-x_k^2}}{2n\sqrt{1-a^2}|x-x_k|}. \quad (6)$$

Using lemma 2, it is not difficult to check, that

$$1 - a^2 x_k^2 = 1 - \frac{a^2 \cos^2 \theta_k}{1 - a^2 \sin^2 \theta_k} = \frac{1 - a^2}{1 - a^2 \sin^2 \theta_k}, \quad (7)$$

$$\sqrt{1 - x_k^2} = \sqrt{1 - \frac{\cos^2 \theta_k}{1 - a^2 \sin^2 \theta_k}} = \frac{\sin \theta_k \sqrt{1 - a^2}}{1 - a^2 \sin^2 \theta_k}. \quad (8)$$

Besides, assuming

$$x = \frac{\cos \theta}{\sqrt{1 - a^2 \sin^2 \theta}}, \quad \theta \in [0, \pi],$$

we obtain

$$\begin{aligned} x - x_k &= \frac{\cos \theta}{\sqrt{1 - a^2 \sin^2 \theta}} - \frac{\cos \theta_k}{\sqrt{1 - a^2 \sin^2 \theta_k}} = \frac{\cos \theta \sqrt{1 - a^2 \sin^2 \theta_k} - \cos \theta_k \sqrt{1 - a^2 \sin^2 \theta}}{\sqrt{1 - a^2 \sin^2 \theta} \sqrt{1 - a^2 \sin^2 \theta_k}} = \\ &= \frac{\cos^2 \theta (1 - a^2 \sin^2 \theta_k) - \cos^2 \theta_k (1 - a^2 \sin^2 \theta)}{\sqrt{1 - a^2 \sin^2 \theta} \sqrt{1 - a^2 \sin^2 \theta_k} (\cos \theta \sqrt{1 - a^2 \sin^2 \theta_k} + \cos \theta_k \sqrt{1 - a^2 \sin^2 \theta})}. \end{aligned}$$

Then

$$\begin{aligned} \cos^2 \theta (1 - a^2 \sin^2 \theta_k) - \cos^2 \theta_k (1 - a^2 \sin^2 \theta) &= \cos^2 \theta - a^2 \cos^2 \theta (1 - a^2 \cos^2 \theta_k) - \\ &- \cos^2 \theta_k + a^2 \cos^2 \theta_k (1 - a^2 \cos^2 \theta) = (1 - a^2) (\cos^2 \theta - \cos^2 \theta_k). \end{aligned}$$

Therefore,

$$x - x_k = \frac{(1 - a^2)(\cos^2 \theta - \cos^2 \theta_k)}{\sqrt{1 - a^2 \sin^2 \theta} \sqrt{1 - a^2 \sin^2 \theta_k} (\cos \theta \sqrt{1 - a^2 \sin^2 \theta_k} + \cos \theta_k \sqrt{1 - a^2 \sin^2 \theta})}. \quad (9)$$

It is clear, that

$$|l_k(x)| = \left| l_k \left(\frac{\cos \theta}{\sqrt{1 - a^2 \sin^2 \theta}} \right) \right|.$$

Then, plugging (7)–(9) into (6), using remark 1, we obtain

$$|l_k(x)| = \frac{|\cos 2n\theta| \sin \theta_k \sqrt{1 - a^2 \sin^2 \theta} |\cos \theta \sqrt{1 - a^2 \sin^2 \theta_k} + \cos \theta_k \sqrt{1 - a^2 \sin^2 \theta}|}{2n(1 - a^2 \sin^2 \theta_k) \sqrt{1 - a^2 \sin^2 \theta} |\cos^2 \theta - \cos^2 \theta_k|}. \quad (10)$$

Now, let $x \in [0, 1]$ and $j, 2 \leq j \leq n$, be such an index that $x \in (x_j, x_{j-1})$. Let us pay attention to the equality (10). If $k \leq j - 1$, then

$$|\cos \theta \sqrt{1 - a^2 \sin^2 \theta_k} + \cos \theta_k \sqrt{1 - a^2 \sin^2 \theta}| < \sqrt{1 - a^2 \sin^2 \theta_k} (\cos \theta + \cos \theta_k).$$

Therefore,

$$|l_k(x)| < \frac{|\cos 2n\theta| \sin \theta_k \sqrt{1 - a^2 \sin^2 \theta}}{2n \sqrt{1 - a^2 \sin^2 \theta_k} (\cos \theta_k - \cos \theta)} < \frac{|\cos 2n\theta| \sin \theta_k}{2n(\cos \theta_k - \cos \theta)} \sqrt{\frac{1 - a^2 \sin^2 \theta_j}{1 - a^2 \sin^2 \theta_k}}.$$

The estimate for the first factor in the right-hand side of the last inequality can be found, for example in [7, p. 528]. For the second factor we use lemma 3. Then

$$\frac{1 - a^2 \sin^2 \theta_j}{1 - a^2 \sin^2 \theta_k} \leq \frac{1 - a^2 \sin^2 \theta_j}{1 - a^2 \sin^2 \theta_{j-1}} < 9.$$

Finally,

$$|l_k(x)| < 9, \quad k \leq j - 1.$$

Then let $j \leq k \leq n$. In this case the factor in the numerator of (10) can be estimated as follows

$$|\cos \theta \sqrt{1 - a^2 \sin^2 \theta_k} + \cos \theta_k \sqrt{1 - a^2 \sin^2 \theta}| < \sqrt{1 - a^2 \sin^2 \theta} (\cos \theta + \cos \theta_k).$$

Thus,

$$|l_k(x)| < \frac{|\cos 2n\theta| \sin \theta_k}{2n(\cos \theta_k - \cos \theta)} \frac{1 - a^2 \sin^2 \theta}{1 - a^2 \sin^2 \theta_k}.$$

It is no difficult to check that the second factor in the right-hand side of the last inequality is not greater than 1 and therefore we obtain

$$|l_k(x)| < 2, \quad j \leq k \leq n.$$

So, we proved lemma when $x \in (x_n, x_1)$. If $x \in [0, x_n] \cup [x_1, 1]$ then in the scheme of the proof of the lemma, there will be obvious changes. Besides, it is clear that

$$\max_{x \in [0, 1]} |l_k(x)| > \max_{x \in [-1, 0]} |l_k(x)|.$$

This concludes the proof.

The main result

According to the definition Lebesgue function of an interpolating process is as follows

$$\Lambda_{2n}(x) = \sum_{k=1}^{2n} |l_k(x)|,$$

and Lebesgue constant is defined as

$$\Lambda_{2n} = \max_{x \in [-1, 1]} \Lambda_{2n}(x), \quad k = 1, 2, \dots$$

Theorem 2. For the Lebesgue constant of the interpolating rational Lagrange process (4) the following estimate holds

$$\Lambda_{2n} \leq C_1(1 + \ln n), \quad n = 1, 2, \dots,$$

where C_1 is a positive constant.

Proof. Here we use modification of the method, applied by S. N. Bernstein to estimate the Lebesgue constant in the polynomial case (see, for example, [7, p. 539]). We are going to check the inequality $\Lambda_{2n}(x) \leq C_1(1 + \ln n)$ for arbitrary $x \in [-1, 1]$. First, it is clear, that if $x = x_k, k = 1, 2, \dots, 2n$, then $\Lambda_{2n}(x_k) = 1$.

Now, let $x \in [0, 1]$. Since the Chebyshev – Markov rational fraction (3) is even,

$$\Lambda_{2n}(x) \leq 2 \sum_{k=1}^n |l_k(x)|.$$

Let $j, j = 1, 2, \dots, n$, be a number such that $x_j < x < x_{j-1}$ (assume that $x_0 = 1$). We represent the sum in the right-hand side of the last equality in the form

$$\Lambda_{2n}(x) \leq 2 \left(S_1(x) + \sum_{k=j-2}^{j+1} |l_k(x)| + S_2(x) \right),$$

where

$$S_1(x) = \sum_{k=1}^{j-3} |l_k(x)|, \quad S_2(x) = \sum_{k=j+2}^n |l_k(x)|.$$

Note that the $S_1(x)$ vanishes for $j = 1, 2, 3$, and the sum $S_2(x)$ vanishes for $j = n - 1, n$. Applying lemma 3 we obtain

$$\Lambda_{2n}(x) \leq 2(S_1(x) + S_2(x) + 24). \tag{11}$$

Let us consider the sum $S_1(x)$. Using definitions of Chebyshev – Markov rational fractions (3) and Lagrange's fundamental polynomials (5), it is easy to see that for $k = 1, 2, \dots, j - 3$

$$|l_k(x)| = \frac{(1 - a^2 x_k^2) \sqrt{1 - x_k^2}}{2n \sqrt{1 - a^2} (x_k - x)}.$$

Assuming $x = \cos \theta$ and applying lemma 2, we obtain

$$|l_k(x)| \leq \frac{\sin \theta_k}{2n(1 - a^2 \sin^2 \theta_k)} \frac{\sqrt{1 - a^2 \sin^2 \theta} (\cos \theta \sqrt{1 - a^2 \sin^2 \theta_k} + \cos \theta_k \sqrt{1 - a^2 \sin^2 \theta})}{\cos^2 \theta_k - \cos^2 \theta}.$$

Given the location of the points $\theta_k, k = 1, 2, \dots, j - 3$, the last estimate can be transformed as follows:

$$|l_k(x)| \leq \frac{\sin \theta_k}{2n(1 - a^2 \sin^2 \theta_k)} \frac{\sqrt{1 - a^2 \sin^2 \theta} \sqrt{1 - a^2 \sin^2 \theta_k} (\cos \theta + \cos \theta_k)}{\cos^2 \theta_k - \cos^2 \theta} \leq \frac{\sin \theta_k}{2n(\cos \theta_k - \cos \theta)}.$$

Therefore,

$$S_1(x) \leq \frac{1}{2n} \sum_{k=1}^{j-3} \frac{\sin \theta_k}{\cos \theta_k - \cos \theta}.$$

Then, since the function $\frac{\sin x}{\cos x - \cos \theta}$ increases for $x \in [0, \theta)$,

$$\frac{\sin \theta_k}{\cos \theta_k - \cos \theta} \leq \frac{\sin x}{\cos x - \cos \theta}, \quad x \in [\theta_k, \theta_{k+1}].$$

Integrating the last inequality with respect to x from θ_k to θ_{k+1} , we get

$$\frac{\pi}{2n} \frac{\sin \theta_k}{\cos \theta_k - \cos \theta} \leq \int_{\theta_k}^{\theta_{k+1}} \frac{\sin x \, dx}{\cos x - \cos \theta}.$$

Then

$$S_1(x) \leq \frac{1}{\pi} \int_{\theta_1}^{\theta_{j-2}} \frac{\sin x \, dx}{\cos x - \cos \theta} < \frac{1}{\pi} \int_0^{\theta_{j-2}} \frac{\sin x \, dx}{\cos x - \cos \theta} = \frac{1}{\pi} \ln \frac{1 - \cos \theta}{\cos \theta_{j-2} - \cos \theta}.$$

Further, we note that

$$1 - \cos \theta \leq 2, \quad \cos \theta_{j-2} - \cos \theta > \cos \theta_{j-2} - \cos \theta_{j-1}.$$

From here,

$$S_1(x) \leq -\frac{1}{\pi} \left(\ln \sin \frac{\theta_{j-2} + \theta_{j-1}}{2} + \ln \sin \frac{\pi}{4n} \right).$$

It remains to note that, $\theta_{j-1} > \theta_{j-2} > \theta_1$ and therefore

$$\ln \sin \frac{\theta_{j-2} + \theta_{j-1}}{2} > \ln \sin \theta_1 = \ln \sin \frac{\pi}{4n}.$$

Finally for the sum $S_1(x)$ we obtain,

$$S_1(x) \leq -\frac{2}{\pi} \ln \sin \frac{\pi}{4n} \leq -\frac{2}{\pi} \ln \frac{1}{2n} = \frac{2}{\pi} \ln n + \frac{2 \ln 2}{\pi}.$$

A similar estimate also holds for the sum $S_2(x)$. Plugging the obtained results in the inequality (11), we conclude that

$$\Lambda_{2n}(x) \leq \frac{8}{\pi} \ln n + \frac{8 \ln 2}{\pi} + 48.$$

The case $x \in [-1, 0]$ can be considered in the same way. Theorem 2 is proved.

Numerical examples

The result of theorem 2 shows that in the case of two geometrically distinct real poles the behavior of the Lebesgue constant does not depend on the choice of the parameters $\{a_k\}_{k=0}^n$. In this section we show results of some numerical experiments for calculation of the Lebesgue function and the Lebesgue constant in the considered case of the choice of the parameters as well as some other cases. Numerical experiment was carried out using programming language *Python*, version 3.6, with the help of libraries *NumPy*, *SciPy* and *Matplotlib*.

Two real poles

Let n be even number, $a_0 = 0$, $a_k = \frac{1}{2}$, $k = 1, 2, \dots, \frac{n}{2}$, $a_k = -\frac{1}{2}$, $k = \frac{n}{2} + 1, \frac{n}{2} + 2, \dots, n$. Then the graph of the function $\frac{\Lambda_{2n}(x)}{1 + \ln n}$ for $n = 30$ is presented on the fig. 1 (horizontal axis – values of x , vertical axis – values of the function $\frac{\Lambda_{2n}(x)}{1 + \ln n}$).

Behavior of the Lebesgue constant in this case is described by the fig. 2 (horizontal axis – values of n , vertical axis – values of the ratio $\frac{\Lambda_{2n}}{1 + \ln n}$).




Fig. 1. Graph of $\frac{\Lambda_{2n}(x)}{1 + \ln n}$ for the case of two real poles and $n = 30$




Fig. 2. Lebesgue constant for the case of two real poles

The results of the experiment support the conclusions of theorem 2 and even allow us to make an assumption about the value of the constant in the statement of this theorem.

Also in the experiment we considered many other cases. Here the results of some of them.

Real poles

Let n be even number, $a_0 = 0$, $a_k = e^{-1/n}$, $k = 1, 2, \dots, \frac{n}{2}$, $a_k = -e^{-1/n}$, $k = \frac{n}{2} + 1, \frac{n}{2} + 2, \dots, n$. Then the graph of the function $\frac{\Lambda_{2n}(x)}{1 + \ln n}$ for $n = 30$ is presented on the fig. 3 (axes mean the same as in the subsection «Two real poles»).

Behavior of the Lebesgue constant in this case is described by the fig. 4.




Fig. 3. Graph of $\frac{\Lambda_{2n}(x)}{1 + \ln n}$ for the case of real poles and $n = 30$




Fig. 4. Lebesgue constant for the case of real poles

Complex conjugate poles

Let n be even number, $a_0 = 0$, $a_k = i e^{-1/\sqrt{n}}$, $k = 1, 2, \dots, \frac{n}{2}$, $a_k = -i e^{-1/\sqrt{n}}$, $k = \frac{n}{2} + 1, \frac{n}{2} + 2, \dots, n$. Then the graph of the function $\frac{\Lambda_{2n}(x)}{1 + \ln n}$ for $n = 30$ is presented on the fig. 5 (axes mean the same as in the subsection «Two real poles»).

Behavior of the Lebesgue constant in this case is described by the fig. 6.




Fig. 5. Graph of $\frac{\Lambda_{2n}(x)}{1 + \ln n}$ for the case of complex conjugate poles and $n = 30$




Fig. 6. Lebesgue constant for the case of complex conjugate poles

The results of all conducted experiments show, that the Lebesgue constant grows with logarithmic rate and allow us to make an assumption of independence of the estimation of the Lebesgue constant from the choice of zeroes of Chebyshev – Markov rational cosine fractions.

Библиографические ссылки

1. Привалов АА. *Теория интерполирования функций*. Саратов: Издательство Саратовского университета; 1990. 229 с.
2. Дзядык ВК. *Введение в теорию равномерного приближения функций полиномами*. Москва: Наука; 1977. 511 с.
3. Русак ВН. Об интерполировании рациональными функциями с фиксированными полюсами. *Доклады Академии наук Белорусской ССР*. 1962;4(9):548–550.
4. Старовойтов АП. О рациональной интерполяции с фиксированными полюсами. *Известия АН БССР. Серия физико-математических наук*. 1983;6:105–106.
5. Ровба ЕА, Дирвук ЕВ. О константе Лебега интерполяционных рациональных процессов Лагранжа по узлам Чебышева – Маркова. *Известия НАН Беларуси. Серия физико-математических наук*. 2015;4:32–38.
6. Ровба ЕА, Козловская НЮ. К вопросу об оценке константы Лебега интерполяционного рационального процесса с узлами Чебышева – Маркова. *Вестник Гродзенскага дзяржаўнага ўніверсітэта імя Янкі Купалы. Серыя 2*. 2016;6(3):6–11.
7. Натансон ИП. *Конструктивная теория функций*. Москва, Ленинград: Гостехиздат; 1949. 688 с.

References

1. Privalov AA. *Teoriya interpolirovaniya funktsii* [Theory of interpolation of functions]. Saratov: Izdatel'stvo Saratovskogo universiteta; 1990. 229 p. Russian.
2. Dzyadyk VK. *Vvedenie v teoriyu ravnomernogo priblizheniya funktsii polinomami* [An Introduction to the theory of uniform approximation of functions by polynomials]. Moscow: Nauka; 1977. 511 p. Russian.
3. Rusak VN. [On interpolation of rational functions with fixed poles]. *Doklady Akademii nauk Belorusskoi SSR*. 1962;4(9):548–550. Russian.
4. Starovoitov AP. [On rational interpolation with fixed poles]. *Izvestiya AN BSSR. Seriya fiziko-matematicheskikh nauk*. 1983; 6:105–106. Russian.
5. Rouba YA, Dirvuk YU. [On a Lebesgue constant of interpolation rational Lagrange processes at the Chebyshev – Markov nodes]. *Izvestiya NAN Belarusi. Seriya fiziko-matematicheskikh nauk*. 2015;4:32–38. Russian.
6. Rouba YA, Kazlouskaya NY. [On a problem of estimation of a Lebesgue constant of interpolation rational processes at the Chebyshev – Markov nodes]. *Vesnik Hrodzenskaha dzjarzhaunaha universiteta imia Ianki Kupaly. Seriya 2*. 2016;6(3):6–11. Russian.
7. Natanson IP. *Konstruktivnaya teoriya funktsii* [Constructive function theory]. Moscow, Leningrad: Gostekhizdat; 1949. 688 p. Russian.

Received by editorial board 28.06.2018.