

Math-Net.Ru

All Russian mathematical portal

A. Shumakovich, Casson invariant of knots associated with
divides,
Algebra i Analiz, 2003, Volume 15, Issue 4, 236–252

<https://www.mathnet.ru/eng/aa816>

Use of the all-Russian mathematical portal Math-Net.Ru implies that you have
read and agreed to these terms of use

<https://www.mathnet.ru/eng/agreement>

Download details:

IP: 18.97.9.170

May 20, 2025, 04:02:04



CASSON INVARIANT OF KNOTS ASSOCIATED WITH DIVIDES

© A. Shumakovich

Abstract. A formula for the Casson invariant of knots associated with divides is presented. The formula is written in terms of Arnold's invariants of pieces of the divide. Various corollaries are discussed.

Introduction

A *divide* P is the image of a generic relative immersion of a compact (not necessarily connected) 1-manifold into the standard unit disk

$$D = \{(x, y) \in \mathbb{R}^2 \mid x^2 + y^2 \leq 1\}$$

in the plane \mathbb{R}^2 . The relativity of the immersion implies that the boundary of the manifold is mapped to the boundary of D . The genericity condition means that a divide has only transversal double points as singularities and is transversal to ∂D at its boundary points. Divides are considered modulo ambient isotopy in D .

With every divide P , a link $L(P) \subset S^3$ can be associated by using the following construction. Consider a standard projection π of

$$S^3 = \{(x, y, u, v) \in \mathbb{R}^4 \mid x^2 + y^2 + u^2 + v^2 = 1\}$$

onto D , i.e., $\pi(x, y, u, v) = (x, y)$. The preimage of a point $p \in D$ under π is either a circle if p belongs to the interior of D , or a single point.

Now $L(P)$ consists of all points $(x, y, u, v) \in \pi^{-1}(P) \subset S^3$ such that either $(x, y) \in \partial P \subset \partial D$, or (u, v) is tangent to P at (x, y) . It is easily seen that for any $p \in P$ the preimage $\pi^{-1}(p) \cap L(P)$ in $L(P)$ consists of either 1, 4, or 2 points, depending on whether p is a boundary, double, or generic point of P . $L(P)$ is indeed a link with $2c + i$ components, where c and i are the numbers of closed and

Key words and phrases. Divide, Casson invariant, Arnold's invariants of plane curves.
Partially supported by the Swiss National Science Foundation.

nonclosed components of P , respectively. Obviously, the ambiently isotopic divides give rise to ambiently isotopic links.

Divides and associated links were originally considered by N. A'Campo [1] and are closely related to the real morsifications of isolated complex plane curve singularities (see [2, 3]).

In this paper I present a formula for the Casson invariant of the knot associated with a given divide with no closed components and only one nonclosed component. Such divides are called the *I-divides* in this paper. The Casson invariant of a link L can be defined as $\frac{1}{2}\Delta_L''(1)$, where $\Delta_L(t)$ is the Alexander polynomial of L and $\Delta_L''(1)$ is the value of its second derivative at 1. It is also a unique Vassiliev invariant of degree 2 that takes the value 0 on the unknot and the value 1 on a trefoil. Since the Casson invariant of a link L is equal to the sum of its values on the components of L , only the case of knots is of interest.

Amazingly enough, the formula is written in terms of Arnold's invariants J^\pm and St of pieces of the divide (see Subsection 1.2 or [6] for the definitions).

In [7], S. Chmutov constructed a second order J^\pm -type invariant of long curves, called J_2^\pm , and proved that its value on an I-divide is equal to the Casson invariant of the associated knot. Since the invariant was defined by an actuality table only, the computation of its values on a curve with even as few as 2 double points was rather involved. Also, the original definition did not provide a mean to compute changes of J_2^\pm under self-tangency *perestroikas*. My formula fills these gaps (see Lemmas 3.2.A and 3.2.C and Subsection 3.3).

Acknowledgments. I am deeply grateful to Norbert A'Campo for valuable remarks and encouragement.

§1. Main ingredients and definitions

1.1. Diagrams of divide links. Let $P \subset D$ be a divide. In this section, I present a way to draw a usual diagram with over- and under-crossings of the link $L(P)$. The following construction is due to M. Hirasawa [10] with some minor modifications (cf. [9]).

First, we perturb P slightly without changing its ambient isotopy type so that the following two conditions be met:

- at every double point the two branches of P are parallel to the main diagonals $y = \pm x$ of the plane;
- P has only finitely many points where the tangent vector is parallel to the y -axis, and at every such point the projection of P to the x -axis has a local extremum.

The next step is to double the divide, i.e., to draw two parallel strands along the divide and to join them at the endpoints (see Figure 1.a). The resulting closed

curves have a natural orientation in accordance with the “*keeping right*” rule. One puts the over- and under-crossings near the double points in accordance with the rule depicted in Figure 1.b. And finally, at every point of P with tangent vector parallel to the y -axis, the strand pointing downwards should make a “*jump through infinity*”. This means to go far right (left) above (below) the rest of the diagram and to return back below (above) it, depending on whether the projection of P to the x -axis has a local maximum or minimum at that point (see Figure 1. c, d). We denote the diagram obtained by $D(P)$. For example, the divide depicted in Figure 1.a gives rise to a (right) trefoil (see Figure 2).




Figure 1. Drawing a diagram of the divide link.




Figure 2. Example of a divide link.

1.1.A. Theorem (Hirasawa [10]). *The diagram $D(P)$ represents the link $L(P)$ in S^3 .*

1.2. Arnold's invariants of plane curves. Let C be a generic plane curve, i.e., a (C^1 -smooth) immersion of the circle into the plane that has only transversal double points as singularities. For any such a curve, its *Whitney index* or simply *index* can be defined as the total rotation number of the vector tangent to the curve. Clearly, this number is the degree of the map that takes every point of the circle to the direction of the tangent vector. The index of C is denoted by $w(C)$. It is easily seen that the index does not change under a *regular homotopy* (i.e., a C^1 -smooth homotopy in the class of C^1 -immersions). Moreover, two plane curves are regularly homotopic if and only if their Whitney indices are the same.




Figure 3. Three types of singularities that may appear in a generic regular homotopy.

A generic regular homotopy connecting two generic curves can contain only a finite number of nongeneric ones. Each of these curves differs from a generic one either at exactly one point of triple transversal self-intersection or at exactly one point of self-tangency. At a point of self-tangency, the velocity vectors of the tangent branches have either the same directions or the opposite ones. In the first case the self-tangency is said to be *direct* and in the second *inverse* (see Fig. 3). The type of self-tangency does not change under reversing of orientation.

Hence there are three types of singular curves that may appear in a generic regular homotopy. Passages through these curves correspond to three *perestroikas* of generic curves.

Consider the triple point *perestroika* more carefully. Just before and just after the passage through a singular curve with a triple point, there is a small *vanishing* triangle close to the place of *perestroika*, which is formed by three branches of the curve. The orientation of the curve determines a cyclic order of sides of the triangle. This is the order in which one meets the sides while traveling along the curve. This cyclic order determines an orientation of the triangle and, therefore, an orientation of its sides. We denote by q the number of sides of the vanishing

triangle for which the orientation obtained coincides with the orientation of the curve (obviously, $q \in \{0, 1, 2, 3\}$).

We define the *sign* of a vanishing triangle as $(-1)^q$. This sign does not change under reversing of orientation of the curve. It is easily seen that before and after the *perestroika* the signs of the vanishing triangles are different.

1.2.A. Definitions (Arnold [6]). 1. A triple point *perestroika* is *positive* if the newborn vanishing triangle is positive.

2. A self-tangency *perestroika* is *positive* if it increases (by 2) the number of self-intersection points of the curve.




Figure 4. The standard curves with Whitney indices $0, \pm 1, \pm \omega$.

The following theorem provides a definition of invariants of generic (plane) curves.

Theorem (Arnold [6]). *There exist three integers $St(C)$, $J^+(C)$, and $J^-(C)$ assigned to an arbitrary generic plane curve C that are uniquely defined by the following properties:*

- (i) St , J^+ , and J^- are invariant under a regular homotopy in the class of generic curves.
- (ii) St does not change under self-tangency *perestroikas* and increases by 1 under a positive triple point *perestroika*.
- (iii) J^+ does not change under triple point and inverse self-tangency *perestroikas* and increases by 2 under a positive direct self-tangency *perestroika*.
- (iv) J^- does not change under triple point and direct self-tangency *perestroikas* and decreases by 2 under a positive inverse self-tangency *perestroika*.
- (v) St , J^+ , and J^- take the following values on the standard curves K_ω shown in Figure 4:

$$\begin{aligned} St(K_0) &= 0, & St(K_{\omega+1}) &= \omega & (\omega = 0, 1, 2, \dots); \\ J^+(K_0) &= 0, & J^+(K_{\omega+1}) &= -2\omega & (\omega = 0, 1, 2, \dots); \\ J^-(K_0) &= -1, & J^-(K_{\omega+1}) &= -3\omega & (\omega = 0, 1, 2, \dots). \end{aligned}$$

Remark. The normalization of St and J^\pm , which is fixed by the last property, makes them additive with respect to the connected sum of curves. It is easily seen that the invariants are independent of the orientation of a curve.

1.3. Formulas for Arnold's invariants. It is very complicated to use the definition given above to compute Arnold's invariants for a given curve with many double points. The two theorems stated below provide explicit formulas, which make such computations easy and straightforward.

Consider a generic plane curve C . For every region r of C , the *index* of r with respect to C is the total rotation number of the radius vector that connects an arbitrary interior point of r with a point traveling along C . Clearly, this number is independent of the choice of the point in r and is denoted by $ind_C(r)$. The *index* $ind_C(e)$ of an edge e of C is the half-sum of the indices of the two regions adjacent to e . The *index* $ind_C(v)$ of a double point v of C is the quarter-sum of the indices of the four regions adjacent to v .




Figure 5. The curve smoothing at a self-intersection point.

1.3.A. Theorem (Viro [13]). Let C be a generic plane curve and \tilde{C} a family of embedded circles obtained as a result of smoothing C at each double point with respect to the orientation (see Figure 5). Then

$$J^+(C) = 1 - \sum_{r \in \mathcal{R}_{\tilde{C}}} ind_C^2(r) \chi(r) + n, \tag{1.1}$$

$$J^-(C) = 1 - \sum_{r \in \mathcal{R}_{\tilde{C}}} ind_C^2(r) \chi(r), \tag{1.2}$$

where $\mathcal{R}_{\tilde{C}}$ is the set of all regions of \tilde{C} , χ is the Euler characteristic, and n is the number of double points of C .

Now, on a generic plane curve C we fix a base point f that is not a double point. One can enumerate all edges by numbers from 1 to $2n$ (where n is again the number of double points of C) following the orientation and assigning 1 to the edge containing f .

Consider an arbitrary double point v of C . There are two edges pointing to v . Let them have numbers i and j such that the tangent vector of the edge i and the tangent vector of the edge j give a positive orientation of the plane. We define the *sign* $s(v)$ of v to be $\text{sign}(i - j)$.

1.3.B. Theorem [12]. *Let C be a generic plane curve. Then*

$$\text{St}(C) = \sum_{v \in \mathcal{V}_C} \text{ind}_C(v)s(v) + \text{ind}_C^2(f) - \frac{1}{4}, \tag{1.3}$$

where \mathcal{V}_C is the set of all double points of C .

Remark. If the point f is chosen on an exterior edge (i.e., an edge that bounds the exterior region), then $\text{ind}_C(f) = \pm \frac{1}{2}$, and

$$\text{St}(C) = \sum_{v \in \mathcal{V}_C} \text{ind}_C(v)s(v). \tag{1.4}$$

Remark. Other formulas for Arnold's invariants can be found in [8, 11, 12].

§2. Formula for the Casson invariant

2.1. The main results. Let $P \subset D$ be an I-divide. Choose an arbitrary orientation on P . For any double point v of P we denote by O_v and I_v the closed and the nonclosed curves¹ obtained by smoothing P at v with regard to the orientation (see Figure 5). The endpoints of P split the boundary ∂D into two arcs. We complete P with one of those arcs so that the orientation of P induce the counter-clockwise orientation on the arc. Let \bar{P} be the resulting closed curve. Finally, for a given plane curve C , we denote $1 - J^-(C)$ by $\tilde{J}(C)$.




Figure 6. Closure of a divide depending on the orientation.

¹Notation rationale: the letters O and I were chosen because they conveniently represent the topological type of the corresponding curves.

Remark. Clearly, the closure \bar{P} of P depends on the orientation of P (see Figure 6).

Remark. $1 - J^-(C)$ can be expressed via a very simple formula $1 - J^-(C) = \sum_{r \in \mathcal{R}_C^-} \text{ind}_C^2(r) \chi(r)$ and, therefore, deserves a special notation.

2.1.A. Theorem. *Let P be an I-divide (i.e., a divide with only one nonclosed component). Then the Casson invariant v_2 of the knot $L(P)$ is given by*

$$v_2(L(P)) = \sum_{v \in \mathcal{V}_P} (\tilde{J}(O_v) + \frac{1}{4} \#(O_v \cap I_v)) + \frac{J^+(\bar{P}) + 2\text{St}(\bar{P})}{4}, \tag{2.1}$$

where \mathcal{V}_P is the set of all double points of P and $\#(O_v \cap I_v)$ is the number of intersection points of O_v and I_v .

Remark. $J^+(\bar{P})$ and $\text{St}(\bar{P})$ obviously depend on the orientation of P , but the sum $J^+(\bar{P}) + 2\text{St}(\bar{P})$ does not, since all the other ingredients of (2.1) are independent of the orientation. This fact can also be verified directly.

2.2. Special cases. It is well known [5] that for a tree-like generic plane curve C the sum $J^+(C) + 2\text{St}(C)$ is always 0. Here a curve C is said to be *tree-like* if smoothing at every double point of C produces two disjoint curves. A *tree-like divide* can be defined in a similar way. From the definition it follows that $\#(O_v \cap I_v) = 0$ for every double point v of a tree-like divide P . Hence, formula (2.1) can be simplified.

2.2.A. *Let P be a tree-like divide. Then*

$$v_2(L(P)) = \sum_{v \in \mathcal{V}_P} \tilde{J}(O_v). \tag{2.2}$$

A slalom divide (see [4] for the definitions) is a special case of tree-like divides. In this case the numbers $\tilde{J}(O_v)$ can be calculated directly.

2.2.B. *Let P be a slalom divide. Then*

$$v_2(L(P)) = \sum_{v \in \mathcal{V}_P} (1 + n(O_v)), \tag{2.3}$$

where $n(O_v)$ is the number of the double points of O_v . Since this number is always nonnegative, the Casson invariant of a slalom knot is always strictly positive.

Remark. Formula 2.1 admits an easy rewriting in terms of the Gauss diagrams of divides (see [11] for the definitions and examples of the technique). In particular, the term $\sum_{v \in \mathcal{V}_P} \frac{1}{4} \#(O_v \cap I_v)$ is half the number of the generic intersections of chords in the Gauss diagram of a divide P .

§3. Proof of Theorem 2.1.A

3.1. Normalization. We denote the right-hand side of (2.1) by $\mathcal{X}(P)$. It is easy to check that the required identity $v_2(L(P)) = \mathcal{X}(P)$ is true for the standard divides D_0, D_1 , etc. Here a standard divide D_k looks like a standard curve K_{k+1} cut at the external edge and appropriately immersed into the unit disk. Indeed, $v_2(L(D_0)) = 0 = \mathcal{X}(D_0)$ and $v_2(L(D_1)) = 1 = \mathcal{X}(D_1)$, because $L(D_1)$ is a trefoil knot (see Fig. 2). Furthermore, $L(D_n)$ is the connected sum of n copies of $L(D_1)$ for $n > 1$, whence $v_2(L(D_n)) = v_2(\#nL(D_1)) = nv_2(L(D_1)) = n = n\tilde{J}(K_1) = \sum_{v \in \mathcal{V}_{D_n}} \tilde{J}(O_v) = \mathcal{X}(D_n)$.

3.2. Invariance under perestroikas. Since any divide can be transformed into a standard one with a finite sequence of Arnold's *perestroikas*, it suffices to check that $v_2(L(P))$ and $\mathcal{X}(P)$ change in the same way under the *perestroikas*. This is proved in the lemmas that follow.

Lemma. *Let P' be the result of a positive inverse self-tangency perestroika of P . We define $\Delta\mathcal{X} = \mathcal{X}(P') - \mathcal{X}(P)$ and $\Delta v_2 = v_2(L(P')) - v_2(L(P))$. Then $\Delta\mathcal{X} = \Delta v_2$.*

Proof. We recall that the divides under consideration are additionally equipped with an orientation. Without loss of generality, we assume that at every double point the two branches of P are oriented upwards. It is always possible to perturb P slightly without changing its isotopy type in order to satisfy this condition, as it is shown in Fig. 7. We also assume that the inverse self-tangency *perestroika* is happening "horizontally", i.e., the two new double points are created side by side as depicted in Figure 8. Denote these two points by a and b .

The common part of P and P' consists of 3 nonclosed curves. Two of them contain the initial and final points of P (or P') and we denote these curves by α and β , respectively. The third curve is denoted by γ . Without loss of generality, we may assume that α, β , and γ are placed as in Fig. 8. Of course, they may intersect each other and may have multiple self-intersections.

Now we observe that there are two crossing points of $L(P')$ close to a and b and such that changing the over- and under-crossings at them allows one to pull 4 strands of $L(P')$ placed between a and b away from each other and to obtain $L(P)$. Those points are marked with small circles in Figure 8 and are denoted by ε_a and ε_b , respectively. More precisely, let K_a be the result of the over- and under-crossing change of $L(P')$ at ε_a , and let K_b be the result of such a change of K_a at ε_b . Then K_b and $L(P)$ are ambiently isotopic, whence $v_2(L(P)) = v_2(K_b)$. It is well known how the value of v_2 behaves under these crossing changes. Namely, the following skein relation is true:

$$v_2 \left(\begin{array}{c} \nearrow \\ \searrow \end{array} \right) - v_2 \left(\begin{array}{c} \nwarrow \\ \swarrow \end{array} \right) = lk \left(\begin{array}{c} \curvearrowright \\ \curvearrowleft \end{array} \right), \tag{3.1}$$




Figure 7. Small perturbation of a divide near a double point.




Fig. 8. Positive inverse self-tangency *perestroika* of a divide.

where lk is the linking number of a two-component link.

Now let L_a be the result of smoothing of $L(P')$ at ϵ_a with regard for the orientation and let L_b be the result of smoothing of K_a at ϵ_b . Both L_a and L_b are links with two components L'_a, L''_a and L'_b, L''_b , respectively. Then (3.1) implies that $v_2(L(P')) - v_2(K_a) = lk(L'_a, L''_a)$ and $v_2(K_a) - v_2(K_b) = lk(L'_b, L''_b)$.

Thus, $\Delta v_2 = lk(L'_a, L''_a) + lk(L'_b, L''_b)$. We compute the first linking number. Denote the diagrams of L'_a and L''_a by D'_a and D''_a , respectively. Then the linking number in question is half the sum of signs of crossings in $D'_a \cap D''_a$. Here the *sign of a crossing* is $+1$ or -1 if it looks like the first or the second summand in (3.1), respectively. In Figure 8 D''_a is depicted with a dotted line.



Figure 9. Contribution of crossing points to $2lk(L'_a, L''_a)$.

D'_a and D''_a may intersect each other either near a double point of the divide, or where a branch of $D(P')$ jumps through infinity. The self-intersections of α and β do not contribute to twice the linking number, because the corresponding pieces




Figure 10. Contribution of "jumps through infinity" to $2lk(L'_a, L''_a)$.

of $D(P)$ belong entirely to D'_a and D''_a , respectively. Contribution of other double points is shown in Figure 9. It is 2 for every intersection point of β with α and γ and every self-intersection of γ . The contribution is 0 for every intersection of α with γ . The points a and b both contribute 2. Hence, the part of $lk(L'_a, L''_a)$ coming from the crossings located near self-intersections of P' is $\#(\alpha \cap \beta) + \#(\beta \cap \gamma) + n(\gamma) + 2$.

Let p be a point of local extremum on P' with respect to the projection to the x -axis. Denote by F_p the corresponding piece of $D(P')$ that makes a jump through the infinity. We may assume that all these pieces are pairwise disjoint and intersect P' transversely at finitely many points. The intersections of F_p with the branches of $D(P')$ corresponding to α and β do not contribute to twice the linking number (see the two leftmost pictures in Figure 10). The contribution from the intersections with γ is 2 if F_p belongs to D'_a (D''_a) and the intersected branch of γ is oriented upwards (downwards), and -2 otherwise (see Figure 10). Moreover, we should add 2 if $p \in \gamma$ and is a local maximum. The total contribution of F_p to twice the linking number is summarized in the following table

Location of p	Contribution of F_p	Location of p	Contribution of F_p
$p \in \overset{\alpha}{\curvearrowright}$	$2 \text{ind}_{\bar{\gamma}}(p)$	$p \in \overset{\gamma}{\curvearrowright}$	$-2 \text{ind}_{\bar{\gamma}}(p) - 1$
$p \in \overset{\alpha}{\curvearrowleft}$	$-2 \text{ind}_{\bar{\gamma}}(p)$	$p \in \overset{\gamma}{\curvearrowleft}$	$2 \text{ind}_{\bar{\gamma}}(p) - 1$
$p \in \overset{\beta}{\curvearrowright}$	$-2 \text{ind}_{\bar{\gamma}}(p)$	$p \in \overset{\gamma}{\curvearrowright}$	$2 \text{ind}_{\bar{\gamma}}(p) + 1$
$p \in \overset{\beta}{\curvearrowleft}$	$2 \text{ind}_{\bar{\gamma}}(p)$	$p \in \overset{\gamma}{\curvearrowleft}$	$-2 \text{ind}_{\bar{\gamma}}(p) + 1$

Here $\bar{\gamma}$ is the natural closure of γ , and $\overset{C}{\curvearrowright}$ and $\overset{C}{\curvearrowleft}$ denote the sets of all local minima and maxima on the curve C , respectively (with an appropriate orientation

if necessary). We recall that $\text{ind}_\gamma(p)$ is half-integer for $p \in \gamma$. Consequently, the contribution of F_p is always even, as it should be.

It is obvious that $\#(\tilde{\gamma}) - \#(\tilde{\gamma}) = 0$, because $\tilde{\gamma}$ is closed. The difference between γ and $\tilde{\gamma}$ is a local maximum near the point b . Therefore, $\#(\tilde{\gamma}) - \#(\gamma) = -1$. Finally,

$$lk(L'_a, L''_a) = \sum_{p \in \alpha \setminus \{u\}^\beta} \text{ind}_{\tilde{\gamma}}(p) - \sum_{p \in \beta \setminus \{u\}^\alpha} \text{ind}_{\tilde{\gamma}}(p) + \sum_{p \in \gamma \setminus \{u\}^\gamma} \text{ind}_{\tilde{\gamma}}(p) - \sum_{p \in \gamma \setminus \{u\}^\gamma} \text{ind}_{\tilde{\gamma}}(p) + \#(\alpha \cap \beta) + \#(\beta \cap \gamma) + n(\gamma) + 3/2.$$

Computation of $lk(L'_b, L''_b)$ is almost the same. We only exchange the roles of α and β throughout and take into account that a does not contribute to the linking number anymore. The contribution of b remains 2. Therefore,

$$\Delta v_2 = 2 \sum_{p \in \gamma \setminus \{u\}^\gamma} \text{ind}_{\tilde{\gamma}}(p) - 2 \sum_{p \in \gamma \setminus \{u\}^\gamma} \text{ind}_{\tilde{\gamma}}(p) + 2\#(\alpha \cap \beta) + \#(\alpha \cap \gamma) + \#(\beta \cap \gamma) + 2n(\gamma) + 2. \tag{3.2}$$

It is simpler to compute $\Delta \mathcal{X}$. First, $J^+(\bar{P}) = J^+(\bar{P}')$ and $\text{St}(\bar{P}) = \text{St}(\bar{P}')$, because neither J^+ nor St changes under an inverse self-tangency *perestroika*. Let v be a double point of P . If v is a self-intersection point of either α , β , or γ , then neither $\tilde{J}(O_v)$ nor $\#(O_v \cap I_v)$ changes. If $v \in \alpha \cap \gamma$ or $v \in \beta \cap \gamma$, then $\tilde{J}(O_v)$ does not change, but $\#(O_v \cap I_v)$ increases by 2 under the *perestroika*. Finally, if $v \in \alpha \cap \beta$, then $\#(O_v \cap I_v)$ does not change, but O_v experiences a positive inverse self-tangency *perestroika* and, hence, $\tilde{J}(O_v)$ increases by 2. Moreover,

$$\#(O_a \cap I_a) = \#(O_b \cap I_b) = \#(\alpha \cap \gamma) + \#(\beta \cap \gamma).$$

Collecting these facts, we obtain

$$\Delta \mathcal{X} = 2\#(\alpha \cap \beta) + \#(\alpha \cap \gamma) + \#(\beta \cap \gamma) + \tilde{J}(O_a) + \tilde{J}(O_b). \tag{3.3}$$

Subtracting (3.3) from (3.2), we get

$$\Delta v_2 - \Delta \mathcal{X} = 2 \sum_{p \in \gamma \setminus \{U\}} \text{ind}_{\tilde{\gamma}}(p) - 2 \sum_{p \in \gamma \setminus \{U\}} \text{ind}_{\tilde{\gamma}}(p) + 2n(\gamma) + 2 - \tilde{J}(O_a) - \tilde{J}(O_b).$$

Lemma 3.2.B below easily implies that this expression is always zero. Indeed,

$$\tilde{J}(O_a) = \tilde{J}(\gamma) + 1 - (\text{ind}_{\tilde{\gamma}}(a) - 1/2) \quad \text{and} \quad \tilde{J}(O_b) = \tilde{J}(\gamma) + \text{ind}_{\tilde{\gamma}}(b) + 1/2$$

with $\text{ind}_{\tilde{\gamma}}(a) = \text{ind}_{\tilde{\gamma}}(b)$, where

$$\tilde{J}(\gamma) := \sum_{p \in \gamma \setminus \{U\}} \text{ind}_{\tilde{\gamma}}(p) - \sum_{p \in \gamma \setminus \{U\}} \text{ind}_{\tilde{\gamma}}(p) + n(\gamma). \quad \bullet$$

3.2.B. Lemma. *Let C be an oriented closed plane curve such that*

- *at every double point, the two branches of C are parallel to the main diagonals $y = \pm x$ of the plane;*
- *C has only finitely many points where the tangent vector is parallel to the y -axis, and at every such point the projection of C to the x -axis has either a local minimum or a local maximum.*

Then

$$\tilde{J}(C) = \#(\text{X}) + \#(\text{X}) + \sum_{p \in C \setminus \{U\}} \text{ind}_C(p) - \sum_{p \in C \setminus \{U\}} \text{ind}_C(p).$$

The proof is elementary and is left to the reader. Hint: smoothing at a double point of C with regard for the orientation does not change the above expression.




Figure 11. Positive direct self-tangency perestroika of a divide.

3.2.C. Lemma. *Let P' be the result of a positive direct self-tangency perestroika of P . We put $\Delta\mathcal{X} = \mathcal{X}(P') - \mathcal{X}(P)$ and $\Delta v_2 = v_2(L(P')) - v_2(L(P))$. Then $\Delta\mathcal{X} = \Delta v_2$.*

The proof is similar to that of Lemma 3.2.A. The corresponding picture of a divide and its link close to the place of perestroika is shown in Figure 11. In this case,

$$\begin{aligned} \Delta v_2 &= 2 \sum_{p \in \gamma \setminus \{u\}} \text{ind}_\gamma(p) - 2 \sum_{p \in \gamma \setminus \{u\}} \text{ind}_\gamma(p) + \#(\alpha \cap \gamma) + \#(\beta \cap \gamma) + 2n(\gamma) + 1 \\ &= \#(\alpha \cap \gamma) + \#(\beta \cap \gamma) + \tilde{J}(O_a) + \tilde{J}(O_b) + 1 = \Delta\mathcal{X}. \end{aligned} \tag{3.4}$$

Now we turn to triple point perestroikas. The definition implies that if a divide P experiences a triple point perestroika, then the ambient isotopy type of the corresponding link does not change.




Figure 12. Different types of triple points.




Figure 13. Positive triple point perestroika of type A_+ .

3.2.D. Lemma. *Let P' be the result of a positive triple point perestroika of P . We put $\Delta\mathcal{X} = \mathcal{X}(P') - \mathcal{X}(P)$. Then $\Delta\mathcal{X} = 0$.*

Proof. We distinguish two kinds of triple points, depending on whether the three branches of a curve (or a divide) at the point are directed into the same half-plane or not. A triple point is said to be of *type B* in the former case and of *type A* in the latter one (see Figure 12). Accordingly, there are two kinds of triple point *perestroikas*. It is easy to check that a triple point *perestroika* of type B is equivalent to a sequence of triple point *perestroikas* of type A and several self-tangency *perestroikas*. Therefore, it suffices to consider only type A *perestroikas* in the proof.

Triple points of type A can further be classified into types A_+ or A_- , depending of whether the cyclic order of the branches determines the positive or the negative orientation of the plane (see Figure 12). Without loss of generality, we shall restrict our consideration to the *perestroikas* of type A_+ only. The corresponding picture of divides before and after a *perestroika* is shown in Figure 13.

Now, it is straightforward to check that

$$\text{St}(\overline{P'}) = \text{St}(\overline{P}) + 1, \tilde{J}(O_{a'}) = \tilde{J}(O_a) - 2, \tilde{J}(O_{b'}) = \tilde{J}(O_b), \tilde{J}(O_{c'}) = \tilde{J}(O_c),$$

and

$$\#(O_{v'} \cap I_{v'}) = \#(O_v \cap I_v) + 2 \quad \text{for } v \in \{a, b, c\}.$$

Hence, $\Delta\mathcal{X} = 1/2 - 2 + 6/4 = 0$. •

Applications to Chmutov's J_2^\pm invariant. J_2^\pm is a second order J^\pm -type invariant of long curves (or I-divides). Chmutov [7] defined it by explicitly specifying its actuality table (i.e., the values of the invariant on all chord diagrams with two chords) and its values on the standard divides with at most one self-tangency point. He also proved that $v_2(L(P)) = J_2^\pm(P)$ for any I-divide P . The definition of J_2^\pm did not allow one to integrate this invariant, i.e., to compute its values on chord diagrams with one chord. Since these values are none other than the changes of J_2^\pm under self-tangency *perestroikas*, formulas (3.3) and (3.4) provide an answer to this question.

3.3.A. Corollary. *Suppose that P is an I-divide and P' is the result of a positive inverse self-tangency perestroika of P . Let $\alpha, \beta, a,$ and b be as in the proof of Lemma 3.2.A (see Figure 8). Then*

$$\begin{aligned} J_2^\pm(P') - J_2^\pm(P) &= 2\tilde{J}(O_a) + 2 \text{ind}_{O_a}(b) + \#(O_a \cap I_a) + 2\#(\alpha \cap \beta) - 1 \\ &= 2\tilde{J}(O_b) - 2 \text{ind}_{O_b}(a) + \#(O_b \cap I_b) + 2\#(\alpha \cap \beta) + 1. \end{aligned} \tag{3.5}$$

The proof follows from (3.3) and the fact that

$$\#(O_a \cap I_a) = \#(O_b \cap I_b) = \#(\alpha \cap \gamma) + \#(\beta \cap \gamma), \tilde{J}(O_a) = \tilde{J}(O_b) - 2 \operatorname{ind}_{O_a}(b) + 1$$

and $\operatorname{ind}_{O_a}(b) = \operatorname{ind}_{O_b}(a)$.

3.3.B. Corollary. *Suppose that P is an I-divide and P' is the result of a positive direct self-tangency perestroika of P . Let a and b be as in the proof of Lemma 3.2.C (see Figure 11). Then*

$$J_2^\pm(P') - J_2^\pm(P) = 2\tilde{J}(O_a) + \#(O_a \cap I_a) = 2\tilde{J}(O_b) + \#(O_b \cap I_b).$$

The proof follows from (3.4) and the fact that

$$\#(O_a \cap I_a) = \#(O_b \cap I_b) = \#(\alpha \cap \gamma) + \#(\beta \cap \gamma) + 1$$

and $\tilde{J}(O_a) = \tilde{J}(O_b)$.

References

- [1] A'Campo N., *Le groupe de monodromie du déploiement des singularités isolées de courbes planes. II*, Proceedings of the International Congress of Mathematicians (Vancouver, BC, 1974), Vol. 1, Canad. Math. Congress, Montreal, Que., 1975, pp. 395-404.
- [2] A'Campo N., *Real deformations and complex topology of plane curve singularities*, Ann. Fac. Sci. Toulouse Math. (6) 8 (1999), 5-23; arXiv: alg-geom/9710023.
- [3] A'Campo N., *Generic immersions of curves, knots, monodromy and Gordian number*, Inst. Hautes Études Sci. Publ. Math. No. 88 (1998), 151-169 (1999); arXiv: math.GT/9803081.
- [4] A'Campo N., *Planar trees, slalom curves and hyperbolic knots*, Inst. Hautes Études Sci. Publ. Math. No. 88 (1998), 171-180 (1999); arXiv: math.GT/9906087.
- [5] Aicardi F., *Tree-like curves*, Singularities and Bifurcations, Adv. Soviet Math., vol. 21, Amer. Math. Soc., Providence, RI, 1994, pp. 1-31.
- [6] Arnol'd V. I., *Plane curves, their invariants, perestroikas and classifications*, Singularities and Bifurcations, Adv. Soviet Math., vol. 21, Amer. Math. Soc., Providence, RI, 1994, pp. 33-91.
- [7] Chmutov S., *An integral generalization of the Gusein-Zade-Natanzon theorem*, arXiv: math.GT/0207238.
- [8] Chmutov S., Duzhin S., *Explicit formulas for Arnold's generic curve invariants*, The Arnold-Gelfand Mathematical Seminars, Birkhäuser Boston, Boston, MA, 1997, pp. 123-138.
- [9] Gibson W., Ishikawa M., *Links of oriented divides and fibrations in link exteriors*, Osaka J. Math. (в печати).
- [10] Hirasawa M., *Visualization of A'Campo's fibered links and unknotting operation*, Proceedings of the First Joint Japan-Mexico Meeting in Topology (Morelia, 1999), Topology Appl. 121 (2002), no. 1-2, 287-304.
- [11] Polyak M., *Invariants of curves and fronts via Gauss diagrams*, Topology 37 (1998), no. 5, 989-1009.

- [12] Шумакович А., *Явные формулы для странности плоской кривой*, Алгебра и анализ 7 (1995), №3, 165–199.
- [13] Viro O. Ya., *First degree invariants of generic curves on surfaces*, Preprint no. 21, Dep. Math. Uppsala Univ., Uppsala, 1994.

Mathematisches Institut
Universität Basel
Rheinsprung 21, CH-4051, Basel
Switzerland

E-mail: Shurik@math.unibas.ch

Поступило 1 октября 2002 г.