

Math-Net.Ru

Общероссийский математический портал

Andrey Kostin, Cohomology of the categorical at zero semigroups,
Algebra Discrete Math., 2006, выпуск 1, 51–66

<https://www.mathnet.ru/adm248>

Использование Общероссийского математического портала Math-Net.Ru подразумевает,
что вы прочитали и согласны с пользовательским соглашением
<https://www.mathnet.ru/rus/agreement>

Параметры загрузки:

IP: 18.97.14.85

18 мая 2025 г., 18:55:34



Cohomology of the categorical at zero semigroups

Andrey Kostin

Communicated by B. V. Novikov

ABSTRACT. In this article we consider the relation between 0-cohomology and extended Eilenberg-MacLane cohomology of categorical at zero semigroups.

1. Introduction

The 0-cohomology theory of semigroups with zero element was introduced in the work [7] as a result of investigation devoted to projective representations of semigroups. This theory was applied also to studying Brauer monoid [9], matrix algebras [11], calculation of the Eilenberg-MacLane cohomology (EM-cohomology) of semigroups (see survey [10] and references there). Unfortunately the 0-cohomology functor is not a derived functor for all semigroups.

Nevertheless the 0-cohomology functor becomes a derived functor if we extend the category of coefficients up to the category of covariant functors from the small category to the category of Abelian groups [5].

On the other hand the 0-cohomology functor is a derived functor for a certain class of semigroups. B. V. Novikov showed that the categorical at zero semigroups belong to this class.

More precisely, [7]: let S be a categorical at zero semigroup, then there is an isomorphism $H_0^n(S, A) \cong H^n(\bar{S}, A)$ for all 0-modules under semigroup S and $n \geq 0$. Here H_0^* denotes the 0-cohomology functor, and \bar{S} is the *gown* of the semigroup S (see sec.2). This theorem provides a relation between 0-cohomology of semigroups and EM-cohomology and takes an important role in the cohomology theory of semigroups.

2000 Mathematics Subject Classification: 20M50.

Key words and phrases: 0-cohomology of semigroups, cohomology of semigroups, cohomology of small categories, categorical at zero semigroups.

The aim of this article is the generalization of this theorem for the category of all natural systems.

The paper consists of five sections. The necessary definitions and theorems are given in second section. The third section is devoted to the proof of the main result of the article, see theorem 3. The examples of applications of the main theorem are considered in section 4. The relation between the theory of Baues's-Wirshing's cohomology for small categories in case of categories without inverse morphisms and 0-cohomology of semigroups is considered in section 5.

I would like to thank Prof. Boris Novikov for his kind assistance and encouragement during the work on this paper.

2. Preliminaries

By S^1 we will denote a semigroup S with an adjoint identity. A semigroup S with zero element is called *categorical at zero* [3] if $abc = 0$ implies $ab = 0$ or $bc = 0$.

A small category \mathbf{C} is called *connected* if for all objects a and b there is a sequence of morphisms $a \longrightarrow c_1 \longleftarrow c_2 \longrightarrow \dots \longrightarrow c_n \longleftarrow b$ of the category \mathbf{C} .

Let $F : \mathbf{C} \longrightarrow \mathbf{D}$ be a covariant functor of small categories and $d \in \text{Ob}\mathbf{D}$. The *comma category* $(F \downarrow d)$ is the category which objects are morphisms $Fc \xrightarrow{\alpha} d$, and a morphism from $Fk \xrightarrow{\alpha} d$ to $Fl \xrightarrow{\beta} d$ is a morphism $\xi : k \longrightarrow l$ such that $\beta \circ F\xi = \alpha$.

Let \mathbf{B} be a category, \mathbf{A} is a subcategory of \mathbf{B} . A functor $R : \mathbf{B} \longrightarrow \mathbf{A}$ is said to be a *reflector* [2], if for all objects $b \in \mathbf{B}$ there is a morphism $\eta_b : b \longrightarrow Rb$ such that each arrow $g : b \longrightarrow a \in \mathbf{A}$ can be represented as $g = f\eta_b$ for unique morphism $f \in \text{Mor}_{\mathbf{A}}(Rb, a)$.

Let us consider the category \mathbf{Sem}_0 whose objects are semigroups with zero and morphisms are mappings $f : S \setminus 0 \longrightarrow T$ such that $f(xy) = f(x)f(y)$ for $xy \neq 0$. As it was shown in [8] there is a reflector $R : \mathbf{Sem}_0 \longrightarrow \mathbf{Sem}$ where \mathbf{Sem} is the category of all semigroups.

The reflector's value RS is called the *gown* of a semigroup S and is denoted by \bar{S} [7].

The gown \bar{S} consists of tuples $\langle s_1, \dots, s_n \rangle$ with $s_i \neq 0$, $s_i s_{i+1} = 0$. In case if S is a categorical at zero semigroup, the product of elements $s = \langle s_1, \dots, s_n \rangle$ and $t = \langle t_1, \dots, t_m \rangle$ is defined by the formula:

$$st = \begin{cases} \langle s_1, \dots, s_n t_1, \dots, t_m \rangle, & \text{if } s_n t_1 \neq 0, \\ \langle s_1, \dots, s_n, t_1, \dots, t_m \rangle, & \text{if } s_n t_1 = 0. \end{cases}$$

For simplicity we sometime will omit brackets in products like $\alpha < u >$ or $< u > \alpha$, where $\alpha \in \bar{S}$ and $u \in S$.

By the symbol $|s|$ we will denote the number of elements in a tuple $s = \langle s_1, \dots, s_n \rangle$.

By a *nerve* of a category \mathbf{C} we will call the set $N_n(\mathbf{C})$ of all tuples $(\alpha_1, \dots, \alpha_n)$ which components are morphisms of the category \mathbf{C} such that the composition $\alpha_{i+1}\alpha_i$ exists for all $1 \leq i \leq n-1$ and $n \geq 1$. The nerve $N_0(\mathbf{C})$ consists of all objects of the category \mathbf{C} .

Let $C_n(\mathbf{C})$ denote a free Abelian group with the set of generators $N_n(\mathbf{C})$. We define a coboundary homomorphism $d_n : C_{n+1}(\mathbf{C}) \rightarrow C_n(\mathbf{C})$ on the generating set by the formula:

$$\begin{aligned} d_n(\alpha_1, \dots, \alpha_{n+1}) &= (\alpha_2, \dots, \alpha_{n+1}) + \\ &\sum_{i=1}^n (-1)^i (\alpha_1, \dots, \alpha_i \alpha_{i+1}, \dots, \alpha_{n+1}) + \\ &(-1)^{n+1} (\alpha_1, \dots, \alpha_n). \end{aligned}$$

The homology of the complex $\{C_n(\mathbf{C}), d_n\}_{n=0}^\infty$ is called an *integral homology of the nerve of the category \mathbf{C}* and is denoted by $H_n(\mathbf{C})$ (see, for example, [12]).

By the symbol $\Delta : \mathbf{Ab} \rightarrow \mathbf{Ab}^{\mathbf{C}}$ we denote the *diagonal functor*, which maps an Abelian group A to the constant functor $\Delta A : \mathbf{C} \rightarrow \mathbf{Ab}$. For an object $c \in \text{Ob}\mathbf{C}$ the value of the functor ΔA is the Abelian group A and $(\Delta A)(f) = \text{id}_A$ for all morphisms f of the category \mathbf{C} .

The right adjoint functor to the functor Δ is called an *inverse limit* $\varprojlim_{\mathbf{C}} : \mathbf{Ab}^{\mathbf{C}} \rightarrow \mathbf{Ab}$.

Theorem 1. [6] *Let $\tau : \mathbf{C} \rightarrow \mathbf{E}$ be a functor of small categories. Then the following conditions are equivalent:*

- The category $(\tau \downarrow e)$ is connected and $H_n(\tau \downarrow e) = 0$ for all $n > 0$, $e \in \mathbf{E}$;*
- For every functor $F : \mathbf{E} \rightarrow \mathbf{Ab}$ the canonical morphism:*

$$\varprojlim_{\mathbf{E}}^n F \rightarrow \varprojlim_{\mathbf{C}}^n F\tau$$

is an isomorphism for all $n > 0$.

Let \mathbf{C} be a small category. The *category of factorization* $\mathbb{F}\mathbf{C}$ [4] is the category whose objects are all morphisms of \mathbf{C} and the set $\text{Mor}_{\mathbf{C}}(f, g)$ consists of three-tuples (α, f, β) such that $\alpha f \beta = g$. A covariant functor $D : \mathbb{F}\mathbf{C} \rightarrow \mathbf{Ab}$ is called a *natural system on category \mathbf{C}* .

If we consider semigroup S^1 as a category with a single object, we obtain the correspondent definitions for the *category of factorizations* $\mathbb{F}S^1$ in semigroup S and *natural system* $\mathbb{F}S^1 \rightarrow \mathbf{Ab}$ on S .

The cohomology of a category \mathbf{C} with coefficients in the natural system D [4] is the Abelian groups $H^n(\mathbf{C}, D) = \text{Ext}^n(Z, D)$ for all $n \geq 0$ where symbol Ext denotes the derived functor of Hom-functor and $Z : \mathbb{F}\mathbf{C} \longrightarrow \mathbf{Ab}$ is the constant functor: $Z(c) \cong \mathbb{Z}$.

Analogously to [13], [4] we define the category of 0-factorizations in the semigroup S with zero \mathbb{F}_0S^1 , whose objects are elements from $S \setminus 0$, and morphism's set $\text{Mor}(a, b)$ consists of three-tuples (α, a, β) where α, β are elements from S , such that $\alpha a \beta = b$ [5].

A 0-natural system on semigroup S with zero is a covariant functor $D : \mathbb{F}_0S^1 \longrightarrow \mathbf{Ab}$. For simplicity we will denote the value of a functor D on an object $a \in \text{Ob}\mathbb{F}_0S^1$ by D_a . Let us denote $\alpha_* = D(\alpha, a, 1)$ and $\beta^* = D(1, a, \beta)$, then $D(\alpha, a, \beta) = \alpha_*\beta^*$ for each morphism (α, a, β) .

For given natural number $n \geq 1$ let us denote by N_nS^1 the set of all n -tuples (a_1, \dots, a_n) of elements from S^1 such that $a_1 \cdots a_n \neq 0$ (the nerve of semigroup S^1). In case $n = 0$ let $N_0S^1 = \{1\}$. The map with the domain on the nerve of S^1 that sends each $a = (a_1, \dots, a_n)$ to an element from $D_{a_1 \cdots a_n}$ is called a n -cochain. The set of all n -cochains is an Abelian group $C_0^n(S^1, D)$ with respect to pointwise addition. For $n = 0$ let $C_0^0(S^1, D) = D_1$.

Let us define the coboundary homomorphism $\delta^n : C_0^n(S^1, D) \longrightarrow C_0^{n+1}(S^1, D)$ by the formula ($n \geq 1$)

$$(\delta f)(a_1, \dots, a_{n+1}) = a_1 * f(a_2, \dots, a_{n+1}) + \sum_{i=1}^n (-1)^i f(a_1, \dots, a_i a_{i+1}, \dots, a_{n+1}) + (-1)^{n+1} a_{n+1}^* f(a_1, \dots, a_n).$$

For the case $n = 0$ we set $(\delta f)(x) = x_* f - x^* f$ for $f \in D_1, x \in S^1 \setminus 0$. It is simple to prove that $\delta^n \delta^{n-1} = 0$. The group of cocycles $Z_0^n(S^1, D)$ is the kernel of the coboundary δ^n . Let $B_0^n(S^1, D)$ denote the group of coboundaries which is the image of δ^{n-1} . The 0-cohomology of a semigroup S with coefficients in 0-natural system D is the Abelian groups

$$H_0^n(S^1, D) = Z_0^n(S^1, D) / B_0^n(S^1, D).$$

Theorem 2. [5] Let S be a semigroup with zero, $D : \mathbb{F}_0S^1 \longrightarrow \mathbf{Ab}$ is a 0-natural system. Then there is an isomorphism

$$H_0^n(S^1, D) \cong \text{Ext}^n(Z, D),$$

which is natural in D , where Z is the constant 0-natural system $Z(s) \cong \mathbb{Z}$.

3. Main theorem

Let S be a categorical at zero semigroup and \bar{S} be the grown of S . Consider the embedding functor $i : \mathbb{F}_0 S^1 \longrightarrow \mathbb{F} \bar{S}^1$ which is defined by the formula:

$$\begin{aligned} i(l) &= \langle l \rangle, \\ i(s, l, p) &= (\langle s \rangle, \langle l \rangle, \langle p \rangle), \end{aligned} \tag{1}$$

for all objects $l \in \text{Ob} \mathbb{F}_0 S^1$ and morphisms $(s, l, p) \in \text{Mor} \mathbb{F}_0 S^1$.

Lemma 1. *The embedding i is full.*

Proof. Let us consider the arrow $\langle l \rangle \xrightarrow{(\sigma, \tau)} \langle t \rangle$ for some $\sigma, \tau \in \bar{S}^1$ and $l, t \in \text{Ob} \mathbb{F}_0 S^1$. Then $\langle t \rangle = \sigma \langle l \rangle \tau$, what implies the following equality $|\sigma \langle l \rangle \tau| = 1$. This is possible only if $\sigma, \tau \in S^1$ and $\sigma l \tau \neq 0$, i.e. $(\sigma, l, \tau) \in \text{Mor} \mathbb{F}_0 S^1$. \square

Let us formulate the main result of the article.

Theorem 3. *Let S be a categorical at zero semigroup, $i : \mathbb{F}_0 S^1 \longrightarrow \mathbb{F} \bar{S}^1$ is the embedding functor which was defined in (1), $D : \mathbb{F} \bar{S}^1 \longrightarrow \mathbf{Ab}$ is a natural system on \bar{S}^1 . Then the functor i induces an isomorphism of cohomology*

$$H^n(\bar{S}^1, D) \cong H^n(S^1, Di)$$

for all $n \geq 0$.

Let us sketch the main steps of the proof. In the first step we consider cohomology functor of semigroup as a derived functor of the limit functor under some factorization category. Further, we explore the integral complex of the nerve of comma category $(i \downarrow s)$ for $s \in \bar{S}^1$. For completion of the proof, due to the theorem 1, it is sufficient to show that the comma category $(i \downarrow s)$ is connected for all $s \in \bar{S}^1$ and that the integral homology of nerve $(i \downarrow s)$ is acyclic. The lemmas 2 and 3 are devoted to this aim.

Proof. In [4] it was shown that the cohomology of a small category \mathbf{C} with coefficients in natural systems is the derived functor of the inverse limit functor:

$$H^n(\mathbf{C}, E) = \varprojlim_{\mathbb{F} \mathbf{C}}^n E, \quad n \geq 0,$$

where $E : \mathbb{F} \mathbf{C} \longrightarrow \mathbf{Ab}$ is a natural system. If we consider the monoid \bar{S}^1 as a category with a single object, then we have the following isomorphism: $H^n(\bar{S}^1, D) \cong \varprojlim_{\mathbb{F} \bar{S}^1}^n D$ for all $n \geq 0$.

Since the 0-cohomology functor is a derived functor (see theorem 2), it is easy to show that $H_0^n(S^1, D) \cong \text{Ext}^n(Z, D) \cong \varprojlim_{\mathbb{F}_0 S^1}^n D$ for all $n \geq 0$.

The embedding functor $i : \mathbb{F}_0 S^1 \rightarrow \mathbb{F} \bar{S}^1$ induces the natural homomorphism:

$$\Psi : \varprojlim_{\mathbb{F} \bar{S}^1}^n D \rightarrow \varprojlim_{\mathbb{F}_0 S^1}^n Di, \quad n \geq 0.$$

By the theorem 1 let us show that under conditions of the theorem 3 the homomorphism Ψ is an isomorphism of cohomologies.

Let us adduce auxiliary notations which will be useful in future. Let $s = \langle s_1, \dots, s_m \rangle$ be an element of the gown \bar{S} . Then define


$$e_k = (\langle s_1, \dots, s_k \rangle, 1, \langle s_{k+1}, \dots, s_m \rangle) \in (i \downarrow s), \quad 0 \leq k \leq m,$$

$$\bar{s}_k = (\langle s_1, \dots, s_{k-1} \rangle, \langle s_k \rangle, \langle s_{k+1}, \dots, s_m \rangle) \in (i \downarrow s), \quad 1 \leq k \leq m.$$

Lemma 2. *Let S be a categorical at zero semigroup, \bar{S} is the gown of S , $i : \mathbb{F}_0 S^1 \rightarrow \mathbb{F} \bar{S}^1$ is the embedding functor. Then the comma category $(i \downarrow s)$ is connected for each $s \in \bar{S}^1$.*

Proof. Consider the elements $(\alpha, l, \beta), (\gamma, t, \delta) \in \text{Ob}(i \downarrow s)$. We define the path which connects these objects. Let $s = \langle s_1, \dots, s_n \rangle$.

a) If $n = 1$, then $(1, s, 1) \in \text{Ob}(i \downarrow s)$ and the commutative diagram



gives the required path.

b) Let $n \geq 2$. It is obvious that the diagram

$$\bar{s}_p \xleftarrow{(s_p, 1)} e_p \xrightarrow{(1, s_{p+1})} \bar{s}_{p+1}$$

shows us that the set $\{e_k\} \cup \{\bar{s}_p\} \subset \text{Ob}(i \downarrow s)$ is connected.

Suppose $(\alpha, l, \beta) = (\langle s_1, \dots, s_{k-1}, \alpha_k \rangle, l, \langle \beta_1, s_{k+1}, \dots, s_n \rangle) \in (i \downarrow s)$ does not belong to the set $\{e_k\} \cup \{\bar{s}_p\}$. It is easy to check that the morphism $(u, l, v) : (\alpha, l, \beta) \rightarrow \bar{s}_k$ of the comma category $(i \downarrow s)$ which is defined by the equation

$$(u, l, v) = \begin{cases} (\alpha_k, l, \beta_1), & \text{if } \alpha_k l \beta_1 \neq 0, \\ (\alpha_k, l, 1), & \text{if } \alpha_k l \neq 0 \text{ and } l \beta_1 = 0, \\ (1, l, \beta_1), & \text{if } \alpha_k l = 0 \text{ and } l \beta_1 \neq 0, \end{cases}$$

for some $1 \leq k \leq n$, defines the path between objects (α, l, β) and \bar{s}_k . \square

The group \hat{C}_0 coincides with C_0 by the definition.

Lemma 3. *The complex $\{\hat{C}_n(i \downarrow s), d_n\}_{n=0}^\infty$ is acyclic.*

Proof. Let us construct a contracting homotopy $\varepsilon_{n+1} : \hat{C}_n \longrightarrow \hat{C}_{n+1}$, $n \geq 1$. Let $\Omega = [\alpha, u_n, \dots, u_1, l, v_1, \dots, v_n, \beta] \in \hat{N}_n(i \downarrow s)$ for elements $\alpha = \langle \alpha_1, \dots, \alpha_k \rangle$, $\beta = \langle \beta_1, \dots, \beta_p \rangle$. For all elements $s = \langle s_1, \dots, s_m \rangle \in \bar{S}$ let us denote $l(s) = \langle s_2, \dots, s_m \rangle$ and $r(s) = \langle s_1, \dots, s_{m-1} \rangle$. Denote by ω the product $u_n \cdots u_1 l v_1 \cdots v_n$. Define the homotopy by the formula:

$$\varepsilon_{n+1}\Omega = \begin{cases} 0, & \text{if } \alpha_k \omega = \omega \beta_1 = 0, \\ (-1)^{n+1} [r(\alpha), \alpha_k, u_n, \dots, v_n, \beta_1, l(\beta)], & \text{if } \alpha_k \omega \beta_1 \neq 0, \\ (-1)^{n+1} [r(\alpha), \alpha_k, u_n, \dots, v_n, 1, \beta], & \text{if } \alpha_k \omega \neq 0, \omega \beta_1 = 0, \\ (-1)^{n+1} [\alpha, 1, u_n, \dots, v_n, \beta_1, l(\beta)], & \text{if } \alpha_k \omega = 0, \omega \beta_1 \neq 0, \end{cases}$$

for all $n \geq 1$.

Let us prove that in the case $n \geq 2$ the diagram

$$\begin{array}{ccccc} \hat{C}_{n+1} & \xrightleftharpoons[d_n]{d_{n+1}} & \hat{C}_n & \xrightleftharpoons[\varepsilon_n]{d_{n-1}} & \hat{C}_{n-1} \end{array}$$

yields the following equation:

$$\varepsilon_n d_{n-1} + d_n \varepsilon_{n+1} = \text{id}_{\hat{C}_n}. \tag{2}$$

Consider three possible cases which correspond to the definition of ε_{n+1} .

Case 1. $\alpha_k \omega = \omega \beta_1 = 0$. Then $d_n \varepsilon_{n+1} \Omega = 0$ and

$$\begin{aligned} \varepsilon_n d_{n-1} \Omega &= \varepsilon_n [\alpha, u_n, \dots, u_1 l v_1, \dots, v_n, \beta] + \\ &\sum_{i=1}^{n-1} (-1)^i \varepsilon_n [\alpha, u_n, \dots, u_{i+1} u_i, \dots, u_1, l, v_1, \dots, v_i v_{i+1}, \dots, v_n, \beta] + \\ &(-1)^n \varepsilon_n [\alpha u_n, \dots, u_1, l, v_1, \dots, v_n \beta]. \end{aligned} \tag{3}$$

By associativity of S^1 all the summands of equality (3) except the last one are equal to zero.

Let $u_n = 1$ then $v_n \neq 1$. Since S is categorial at zero we conclude that $v_n \beta_1 = 0$. Thus we have

$$\begin{aligned} \varepsilon_n [\alpha u_n, u_{n-1}, \dots, v_{n-1}, v_n \beta] &= \varepsilon_n [\alpha, u_{n-1}, \dots, v_{n-1}, \langle v_n, \beta_1 \rangle] = \\ &(-1)^n [\alpha, 1, u_{n-1}, \dots, v_{n-1}, v_n, \beta] = (-1)^n \Omega \end{aligned}$$

The proof of the equation (2) in the case when $u_n \neq 1$ and $v_n = 1$ is dual with the given one. Now let $u_n \neq 1$ and $v_n \neq 1$. Since S is categorical at zero we have $\alpha_k u_n = v_n \beta_1 = 0$. Hence

$$\begin{aligned} \varepsilon_n[\alpha u_n, u_{n-1}, \dots, v_{n-1}, v_n \beta] = \\ \varepsilon_n[\langle \dots, \alpha_k, u_n \rangle, u_{n-1}, \dots, v_{n-1}, \langle v_n, \beta_1, \dots \rangle] = (-1)^n \Omega. \end{aligned}$$

So the proof of the equation (2) is finished for the first case.

Case 2. Let $\Omega \in \hat{N}_n(i \downarrow s)$ such that $\alpha_k \omega \beta_1 \neq 0$. Then

$$\begin{aligned} d_n \varepsilon_{n+1} \Omega = (-1)^{n+1} d_n[r(\alpha), \alpha_k, u_n, \dots, v_n, \beta_1, l(\beta)] = \\ (-1)^{n+1}([r(\alpha), \alpha_k, u_n, \dots, u_1 l v_1, \dots, v_n, \beta_1, l(\beta)] + \\ \sum_{i=1}^{n-1} (-1)^i [r(\alpha), \alpha_k, \dots, u_{i+1} u_i, \dots, v_i v_{i+1}, \dots, \beta_1, l(\beta)] + \\ (-1)^n [r(\alpha), \alpha_k u_n, u_{n-1}, \dots, v_{n-1}, v_n \beta_1, l(\beta)] + (-1)^{n+1} \Omega) \end{aligned} \quad (4)$$

Evidently the sum of the first n summands of (3) equals to the correspondent sum of (4) with the opposite sign. By the definition of the product in \bar{S}^1 the last summand of the equality (3) has the form:

$$\begin{aligned} (-1)^n \varepsilon_n[\alpha u_n, u_{n-1}, \dots, v_{n-1}, v_n \beta] = \\ (-1)^n \varepsilon_n[\langle \dots, \alpha_{k-1}, \alpha_k u_n \rangle, u_{n-1}, \dots, v_{n-1}, \langle v_n \beta_1, \beta_2, \dots \rangle] = \\ (-1)^n [r(\alpha), \alpha_k u_n, \dots, v_n \beta_1, l(\beta)]. \end{aligned}$$

This statement ends the proof of the second case.

Case 3. In case when $\alpha_k \omega = 0$ and $\omega \beta_1 \neq 0$ (the case when $\alpha_k \omega \neq 0$ and $\omega \beta_1 = 0$ is dual) the proof of (2) is similar with the second case.

Now we start the proof of equation (2) for $n = 1$. Let $s = \langle s_1, \dots, s_m \rangle$, $m \geq 2$ (the case when $m = 1$ is evident).

Define the homomorphism $\varepsilon_1 : C_0(i \downarrow s) \longrightarrow \hat{C}_1(i \downarrow s)$, $m \geq 2$. Introduce the following auxiliary notations:

$$\begin{aligned} s_k^l = [\langle s_1, \dots, s_{k-1} \rangle, s_k, 1, 1, \langle s_{k+1}, \dots, s_m \rangle] \in \hat{C}_1(i \downarrow s), \quad 1 \leq k \leq m, \\ s_k^r = [\langle s_1, \dots, s_{k-1} \rangle, 1, 1, s_k, \langle s_{k+1}, \dots, s_m \rangle] \in \hat{C}_1(i \downarrow s), \quad 1 \leq k \leq m. \end{aligned}$$

We give ε_1 by the formulas:

$$\begin{aligned} \varepsilon_1 e_k = \sum_{i=1}^k s_i^r + \sum_{i=k+1}^m s_i^l, \quad 0 \leq k \leq m, \\ \varepsilon_1 \bar{s}_k = \sum_{i=1}^k s_i^r + \sum_{i=k}^m s_i^l, \quad 1 \leq k \leq m. \end{aligned}$$

Now consider the case when the element $\gamma = [\alpha, u, \beta] \in C_0(i \downarrow s)$ does not belong to the set $\{e_k, \bar{s}_l\}_{k,l}$. Then define

$$\varepsilon_1(\gamma) = \begin{cases} -[r(\alpha), \alpha_k, u, 1, 1, \beta] + \varepsilon_1 \bar{s}_k, & \text{if } u \neq 1, \alpha_k u \neq 0, u\beta_1 = 0, \\ -[\alpha, 1, u, \beta_1, l(\beta)] + \varepsilon_1 \bar{s}_{k+1}, & \text{if } u \neq 1, \alpha_k u = 0, u\beta_1 \neq 0, \\ -[r(\alpha), \alpha_k, u, \beta_1, l(\beta)] + \varepsilon_1 \bar{s}_k, & \text{if } \alpha_k u \beta_1 \neq 0. \end{cases}$$

Let $\Omega = [\alpha, u_1, l, v_1, \beta] \in \hat{N}_1(i \downarrow s)$. Denote $\omega = u_1 l v_1$. Consider three cases.

Case 1: $\alpha_k \omega \beta_1 \neq 0$. Then we have:

$$\varepsilon_1 d_0 \Omega = \varepsilon_1 [\alpha, u_1 l v_1, \beta] - \varepsilon_1 [\alpha u_1, l, v_1 \beta] = -[r(\alpha), \alpha_k, u_1 l v_1, \beta_1, l(\beta)] + \varepsilon_1 \bar{s}_k + [r(\alpha), \alpha_k u_1, l, v_1 \beta_1, l(\beta)] - \varepsilon_1 \bar{s}_k.$$

In the same time

$$d_1 \varepsilon_2 \Omega = d_1 [r(\alpha), \alpha_k, u_1, l, v_1, \beta_1, l(\beta)] = [r(\alpha), \alpha_k, u_1 l v_1, \beta_1, l(\beta)] - [r(\alpha), \alpha_k u_1, l, v_1 \beta_1, l(\beta)] + \Omega.$$

This ends the exploration of the first case.

Case 2: $\alpha_k \omega = 0$ and $\omega \beta_1 \neq 0$.

a) $l \neq 1$. Then

$$\begin{aligned} \varepsilon_1 d_0 \Omega &= -[\alpha, 1, u_1 l v_1, \beta_1, l(\beta)] + \varepsilon_1 \bar{s}_{k+1} + [\alpha, u_1, l, v_1 \beta_1, l(\beta)] - \varepsilon_1 \bar{s}_{k+1}, \\ d_1 \varepsilon_2 \Omega &= d_1 [\alpha, 1, u_1, l, v_1, \beta_1, l(\beta)] = [\alpha, 1, u_1 l v_1, \beta_1, l(\beta)] - [\alpha, u_1, l, v_1 \beta_1, l(\beta)] + \Omega. \end{aligned}$$

b) $l = 1, u_1 = 1, v_1 \neq 1$. In this case we have:

$$\begin{aligned} \varepsilon_1 d_0 \Omega &= \varepsilon_1 [\alpha, v_1, \beta] - \varepsilon_1 [\alpha, 1, \langle v_1 \beta_1, \dots \rangle] = -[\alpha, 1, v_1, \beta_1, l(\beta)] + \varepsilon_1 \bar{s}_{k+1} - \varepsilon_1 e_k = -[\alpha, 1, v_1, \beta_1, l(\beta)] + s_{k+1}^r, \\ d_1 \varepsilon_2 \Omega &= d_1 [\alpha, 1, 1, 1, v_1, \beta_1, l(\beta)] = [\alpha, 1, v_1, \beta_1, l(\beta)] - s_{k+1}^r + \Omega. \end{aligned}$$

c) $l = 1, u_1 \neq 1, v_1 = 1$. Then we obtain

$$\begin{aligned} \varepsilon_1 d_0 \Omega &= \varepsilon_1 [\alpha, u_1, \beta] - \varepsilon_1 [\langle \dots, \alpha_k, u_1 \rangle, 1, \beta] = -[\alpha, 1, u_1, \beta_1, l(\beta)] + \varepsilon_1 \bar{s}_{k+1} + [\alpha, u_1, 1, \beta_1, l(\beta)] - \varepsilon_1 \bar{s}_{k+1}, \\ d_1 \varepsilon_2 \Omega &= [\alpha, 1, u_1, \beta_1, l(\beta)] - [\alpha, u_1, 1, \beta_1, l(\beta)] + \Omega. \end{aligned}$$

This ends our consideration of the second case. It is clear that the proof in case when $\alpha_k \omega \neq 0$ and $\omega \beta_1 = 0$ is similar to the considered one.

Case 3: $\alpha_k \omega = 0$ and $\omega \beta_1 = 0$. Then $(\alpha, u_1 l v_1, \beta) = \bar{s}_{k+1} \in C_0(i \downarrow s)$, $\varepsilon_2 \Omega = 0$. We have

$$\varepsilon_1 d_0 \Omega = \varepsilon_1 \bar{s}_{k+1} - \varepsilon_1 (\alpha u_1, l, v_1 \beta).$$

a) $l \neq 1$ or $l = 1$ when $u_1 \neq 1$ and $v_1 \neq 1$. Then $\varepsilon_1 d_0 \Omega = \varepsilon_1 \bar{s}_{k+1} + \Omega - \varepsilon_1 \bar{s}_{k+1} = \Omega$.

b) $l = 1$ and $u_1 = 1$. Then $\varepsilon_1 d_0 \Omega = \varepsilon_1 \bar{s}_{k+1} - \varepsilon_1 e_k = s_{k+1}^r = \Omega$.

c) $l = 1$ and $v_1 = 1$. In this case $\varepsilon_1 d_0 \Omega = \varepsilon_1 \bar{s}_{k+1} - \varepsilon_1 e_{k+1} = s_{k+1}^l = \Omega$.

In such a way the equation (2) for the case $n = 1$ is proven. The proof of the lemma as well as the main theorem is completed. \square

\square

Remark. There is another form of the described isomorphism. Let $i^* : \mathbf{Ab}^{\mathbb{F}\bar{S}^1} \longrightarrow \mathbf{Ab}^{\mathbb{F}_0 S^1}$ be the restriction functor which is induced by i . Let us denote by Lan_i left Kan extension along the functor i ($Lan_i \dashv i^*$). Since i is full, we get from [2] (Ch. X)

$$i^*(Lan_i G) = G,$$

for each functor $G : \mathbb{F}_0 S^1 \longrightarrow \mathbf{Ab}$. Replacing D by $Lan_i G$ in the theorem 3, we obtain the other form of the isomorphism:

$$H_0^n(S, G) \cong H^n(\bar{S}, Lan_i G).$$

Corollary. [7]. *Let S be a categorical at zero semigroup, \bar{S} its grown. Then*

$$H^n(\bar{S}, A) \cong H_0^n(S, A), \quad n \geq 0,$$

for each 0–module A over S .

4. Examples

Example 1. Let us compute the cohomology of the semigroup W with the representation: $\langle a_i, b_j, w \mid a_i b_i = w, 1 \leq i, j \leq n \rangle$. Let the ideal $I = W \setminus \{a_i, b_j, w \mid 1 \leq i, j \leq n\}$. It is simple to check that Rees factor semigroup $S = W/I$ is categorical at zero and the grown $\bar{S} = W$.

Using theorem 3 for computing the cohomology of W , it is sufficient to calculate 0–cohomology of the semigroup S .

Let $\mathbb{F}_0 S^1$ be the category of 0–factorizations of S^1 and $D : \mathbb{F}_0 S^1 \longrightarrow \mathbf{Ab}$ is a natural system. Consider the normalized complex:

$$\hat{C}_0^0(S^1, D) \longrightarrow \hat{C}_0^1(S^1, D) \longrightarrow \hat{C}_0^2(S^1, D) \longrightarrow \dots \longrightarrow \hat{C}_0^n(S^1, D) \longrightarrow \dots$$

It is obvious that $\hat{C}_0^p(S^1, D) = 0$ if $p \geq 3$. Let $f \in \hat{Z}_0^2(S^1, D) = \hat{C}_0^2(S^1, D)$. Then f is defined by its values on the elements of the set $\{(a_k, b_k)\}_{k=1}^n \subset S \times S$. In that way we have: $\hat{Z}_0^2(S^1, D) \cong \bigoplus_{i=1}^n D_w$.

Now let $f \in \hat{B}_0^2(S^1, D)$. It means that there is a function $h \in \hat{C}_0^1(S^1, D)$ such that

$$f(a_i, b_i) = D(a_i, 1)h(b_i) - h(w) + D(1, b_i)h(a_i), \quad 1 \leq i \leq n.$$

Denote by M_i an Abelian group which consists of elements $D(a_i, 1)x + D(1, b_i)y$ for all $x \in D_{b_i}, y \in D_{a_i}, 1 \leq i \leq n$.

Let K be a subset of $\bigoplus_{i=1}^n D_w$ of the tuples (m_1, \dots, m_n) where $m_i \in t + M_i$ for some $t \in D_w, 1 \leq i \leq n$. Then $H_0^2(S^1, D) \cong (\bigoplus_{i=1}^n D_w)/K$. Let us compute this factor group.

Consider the map

$$\eta : D_w/M_1 \oplus D_w/M_2 \oplus \dots \oplus D_w/M_n \longrightarrow D_w^n/K, \quad (5)$$

which is given by the rule:

$$\eta(x_1 + M_1, x_2 + M_2, \dots, x_n + M_n) = (x_1, x_2, \dots, x_n) + K.$$

We now verify that η is a correctly defined map. Let $x'_i \in x_i + M_i, 1 \leq i \leq n$. Then $\eta(x_1 + M_1, x_2 + M_2, \dots, x_n + M_n) - \eta(x'_1 + M_1, x'_2 + M_2, \dots, x'_n + M_n) = (x_1 - x'_1, \dots, x_n - x'_n) + K = 0$ since $x_i - x'_i \in M_i$, and the map η is correctly defined.

Obviously η is a surjective map. Let us calculate the kernel L of η . Let $\eta(x_1 + M_1, x_2 + M_2, \dots, x_n + M_n) = (t + m_1, \dots, t + m_n)$ for some $t \in M, m_i \in M_i, 1 \leq i \leq n$. It follows that

$$x_i = t + m_i, \quad 1 \leq i \leq n.$$

Thus L consists of tuples of cosets $(t + M_1, \dots, t + M_n)$ generated by an element $t \in D_w$. By the Noether Theorem we conclude that

$$H^2(S^1, D) \cong \left(\bigoplus_{i=1}^n D_w/M_i \right) / L, \quad L = \{(t + M_1, \dots, t + M_n) \mid t \in D_w\} \quad (6)$$

Example 2. Let us compute the second cohomology group of the semigroup S from the example 1 for more specific case. We will need this example in section 5.

Let $S = W/I$ be the semigroup from example for $n = 3, A$ be an Abelian finite group of odd order, $D : \mathbb{F}_0 S^1 \longrightarrow \mathbf{Ab}$ be a 0-natural system. For each nonzero element $s \in S$ the value of the functor D is the Abelian group $D_s = A \oplus A$. If (s, u, t) be a morphism from $\mathbb{F}_0 S^1, a = (a_1, a_2) \in D_u$ then:

$$D(s, u, t)a = s_* t^* a = \begin{cases} a, & \text{if } s = t = 1, \\ (a_1 + a_2, a_1 + a_2), & \text{if } s = a_1 \text{ and } t = 1, \\ (a_1 - a_2, a_2 - a_1), & \text{if } s \in \{a_2, a_3\} \text{ and } t = 1, \\ 0, & \text{otherwise.} \end{cases}$$

It is easy to check that D is a covariant functor.

Now compute the second 0-cohomology group $H_0^2(S, D)$.

Using notation from example 1 we have:

$$\begin{aligned} M_1 &= a_{1*}D_{b_1} = \{(x, x) \mid x \in A\}, \\ M_i &= a_{i*}D_{b_i} = \{(x, -x) \mid x \in A\}, \quad i = 2, 3, \\ K &= \{(v + m_1, v + m_2, v + m_3) \mid v \in D_w, m_i \in M_i\}. \end{aligned}$$

Define the map $\varphi : \bigoplus_{i=1}^3 D_w/M_i \longrightarrow A^3$ by the formula

$$\varphi(l + M_1, p + M_2, k + M_3) = (l_1 - l_2, p_1 + p_2, k_1 + k_2),$$

where $l = (l_1, l_2)$, $p = (p_1, p_2)$ and $k = (k_1, k_2)$.

Lemma 4. *The map φ is an isomorphism of Abelian groups. The inverse map φ^{-1} is defined by the formula $\varphi^{-1}(a_1, a_2, a_3) = ((a_1, 0) + M_1, (a_2, 0) + M_2, (a_3, 0) + M_3)$.*

Consider the epimorphism $\psi = \eta \circ \varphi^{-1} : A^3 \longrightarrow D_w^3/K$ where

$$\eta : D_w/M_1 \bigoplus D_w/M_2 \bigoplus D_w/M_3 \longrightarrow D_w^3/K$$

is the homomorphism which was defined in (5). Using the formula (6) we obtain $H_0^2(S^1, D) = A^3/Ker\psi$.

Let us compute the kernel ψ . Let $(a_1, a_2, a_3) \in Ker\psi$. Then

$$\begin{aligned} ((a_1, 0), (a_2, 0), (a_3, 0)) \in K &\Leftrightarrow \\ ((a_1, 0), (a_2, 0), (a_3, 0)) &= (t + (l_1, l_2), t + (l_2, -l_2), t + (l_3, -l_3)) \end{aligned}$$

for some $t_i, l_i \in A$. It follows that $a_1 = T - P$ and $a_2 = a_3 = T + P$ where $T, P \in A$. Since A has odd order, $Ker\psi = \{(a, b, b), a, b \in A\}$. Thus we obtain

$$H_0^2(S^1, D) \cong A. \quad (7)$$

5. Cohomology of categories without inverse morphisms

A small category \mathbf{C} is called a *category without inverse morphisms* if for all morphisms f, g from $f \circ g = id_x$ it follows $f = g = id_x$, $x \in Ob\mathbf{C}$.

Let \mathbf{C} be a category without inverse morphisms. Define the semi-group $S_N\mathbf{C}$. The elements of $S_N\mathbf{C}$ are all nonidentical morphisms of \mathbf{C} and a zero element. For all $f, g \in S_N\mathbf{C} \setminus \{0\}$ define the multiplication by the formula:

$$fg = \begin{cases} \bar{f} \circ \bar{g}, & \text{if the composition } \bar{f} \circ \bar{g} \text{ exists,} \\ 0, & \text{otherwise.} \end{cases}$$

Here and further on for the element $f \in S_N\mathbf{C}$ we denote by \bar{f} the correspondent morphism of the category \mathbf{C} .

Let us define the map

$$* : \text{Mor}\mathbf{C} \longrightarrow \text{Mor}\mathbf{C} \cup \{1\}$$

by the rule

$$f^* = \begin{cases} 1, & \text{if } f = \text{id}_x, x \in \text{Ob}\mathbf{C}, \\ f, & \text{otherwise.} \end{cases}$$

Lemma 5. *Let \mathbf{C} be a small category without inverse morphisms. The map $*$ can be extended up to the equivalence of categories $* : \mathbb{F}\mathbf{C} \xrightarrow{\sim} \mathbb{F}_0S_N\mathbf{C}^1$.*

Proof. Let the value of the functor $*$ for object $f \in \text{Ob}\mathbb{F}\mathbf{C}$ be $f^* \in \text{Ob}\mathbb{F}_0S_N\mathbf{C}^1$ and (α^*, f^*, β^*) be the image of the morphism $(\alpha, f, \beta) \in \text{Mor}\mathbb{F}\mathbf{C}$. It is simple to check that $*$ is well-defined covariant functor.

Let us prove that the functor $*$ is an equivalence of categories. Denote by ψ the map $\text{Mor}\mathbb{F}\mathbf{C}(f, g) \longrightarrow \text{Mor}_{\mathbb{F}_0S_N\mathbf{C}^1}(f^*, g^*)$ which is induced by the functor $*$.

Ensure that ψ is injective. Let $(\alpha^*, f^*, \beta^*) = (\alpha^*, k^*, \beta^*)$ for some morphisms $(\alpha, f, \beta), (\alpha, k, \beta) \in \text{Mor}\mathbb{F}\mathbf{C}(f, g)$. Let g be a nonidentical morphism. Consider two cases. If $f^* = 1$ then $f = k = \text{id}_{\text{dom}\alpha}$. In case if $f^* \neq 1$ the morphism $f = k = \bar{f}^*$. If g is the identical morphism, it is obvious that ψ is injective.


Let us check surjectivity. If (s_1, f^*, s_2) be a morphism $\mathbb{F}_0S_N\mathbf{C}^1$ then $s_1f^*s_2 \neq 0$ and the composition $\bar{s}_1 \circ \bar{f}^* \circ \bar{s}_2$ exists. It is obvious that $(\bar{s}_1, \bar{f}^*, \bar{s}_2)$ is the necessary preimage of (s_1, f^*, s_2) . \square

Since the equivalence of the categories implies the isomorphism of cohomologies, we have proved the following

Theorem 4. *Let \mathbf{C} be a small category without inverse morphisms, $D : \mathbb{F}_0S_N\mathbf{C}^1 \longrightarrow \mathbf{Ab}$ is a 0-natural system. Then there is the isomorphism of cohomologies:*

$$H^n(\mathbf{C}, D^*) \cong H_0^n(S_N\mathbf{C}^1, D), \quad n \geq 0.$$

Example 3. Let \mathbf{E} be a small category which is defined by the commutative diagram



with $w = a_i b_i$. Obviously \mathbf{E} is the category without inverse morphisms and $S_N \mathbf{E}$ coincides with the semigroup S from example 1.

Using the result from this example and theorem 4 we get

Proposition. Let $P : \mathbb{F}\mathbf{E} \longrightarrow \mathbf{Ab}$ be a natural system on \mathbf{E} , $\tau : \mathbb{F}_0 S_N \mathbf{C}^1 \xrightarrow{\sim} \mathbb{F}\mathbf{C}$ is an equivalence of categories. Then

$$H^2(\mathbf{E}, P) = \left(\bigoplus_{i=1}^n (P\tau)w / K_i \right) / L, \quad L = \{ (m + K_1, \dots, m + K_n) \mid m \in (P\tau)w \},$$

where K_i is a subgroup of $(P\tau)w$ which consists of the elements $(P\tau)(a_i, 1)x + (P\tau)(1, b_i)y$.

Remark. The cohomology of the category \mathbf{E} from example 3 was explored in [4]. In that work the following result was introduced without proof

$$H^2(\mathbf{E}, P) = P_w / I_2 \bigoplus P_w / I_3, \tag{8}$$

with $I_2 = a_{1*}P_{b_1} + a_{2*}P_{b_2}$ and $I_3 = a_{1*}P_{b_1} + a_{3*}P_{b_3}$. It is erroneous.

Indeed, let us consider the functor $P = D^*$ where D is the 0-natural system which was defined in (7). Then $I_2 = M_1 + M_2 = D_w$, $I_3 = M_1 + M_3 = D_w$ and formula (8) becomes the form $H^2(\mathbf{E}, D^*) = 0$.

From the other hand, using the result that was received in (7) we obtain the formula $H^2(\mathbf{E}, D^*) \cong A$ which contradicts with (8).

References

[1] S. MacLane. *Homology*. Berlin-Goettingen-Heidelberg: Springer, 1963.
 [2] S. MacLane. *Categories for the working mathematician*. Springer, 1971.
 [3] A. H. Clifford, G. B. Preston. *Algebraic Theory of Semigroups*. Amer. Math. Soc., Providence, 1964.

- [4] H.-J. Baues, G. Wirshing. *Cohomology of small categories*. J. Pure Appl. Algebra, **38**(1985), N 2/3, 187-211.
- [5] A. A. Kostin, B. V. Novikov. *Semigroup cohomology as a derived functor*. Filomat (Yugoslavia), v.15 (2001), 17-23.
- [6] U. Oberst. *Basiserweiterung in der Homologie kleiner Kategorien*. Math. Z. **100**, 36-58 (1967).
- [7] B. V. Novikov. *On 0-cohomologies of semigroups*. Teor. Appl. Quest. Diff. Eq. and Algebra, Kiev, 1978, 185-188 (Russian).
- [8] B. V. Novikov. *Defining relations and O-modules over a semigroup*. In: Theory of Semigr. and its Appl. (Saratov), 1983, 94-99 (Russian).
- [9] B. V. Novikov. *On the Brauer monoid*. Matem. zametki, **57** (1995), No. 4, 633-636 (Russian).
- [10] B. V. Novikov. *Semigroup cohomology and applications*. Algebra — Representation Theory (ed. K. W. Roggenkamp and M. Ştefănescu), Kluwer, 2001, 311-318.
- [11] W. E. Clark. *Cohomology of semigroups via topology with an application to semigroup algebras*. Commun. Algebra **4** (1976), 979-997.
- [12] Gabriel P., Zisman M. *Calculus of fractions and homotopy theory*. Berlin-Heidelberg-New York: Springer-Verlag, 1967.
- [13] Leech J. *Cohomology theory for monoid congruences*. Houston J. Math., **11** (1985), No. 2, 207-223.

CONTACT INFORMATION

Andrey Kostin

ul. Luk'yanovskaya 21, kv. 99, 04071,
Kiev, Ukraine
E-Mail: kostin_a@inbox.ru